

Functional brain organization of preparatory attentional control in visual search

Patrick Bourke<sup>1</sup>, Steven Brown<sup>2</sup>, Elton Ngan<sup>3</sup> and Mario Liotti<sup>2</sup>

<sup>1</sup>Department of Psychology, University of Lincoln, UK;

<sup>2</sup>Department of Psychology, Simon Fraser University, Burnaby, Canada;

<sup>3</sup>Department of Psychology, University of British Columbia, Vancouver, Canada

Corresponding Author: Patrick A. Bourke, Department of Psychology, Brayford Pool, Lincoln LN6 7TS, United Kingdom. e-mail: [pbourke@lincoln.ac.uk](mailto:pbourke@lincoln.ac.uk); Tel : +44 1522 886180; Fax: +44 1522 886026.

### **Abstract**

Looking for an object that may be present in a cluttered visual display requires that an advanced specification of that object be created and then matched against the incoming visual input. Here, fast event-related fMRI was used to identify the brain networks that are active when preparing to search for a visual target. By isolating the preparation phase of the task it has been possible to show that for an identical stimulus, different patterns of cortical activation occur depending on whether participants anticipate a 'feature' or a 'conjunction' search task. When anticipating a conjunction search task, there was more robust activation in ventral occipital areas, new activity in the transverse occipital sulci and right posterior intraparietal sulcus. In addition, preparing for either type of search activated ventral striatum and lateral cerebellum. These results suggest that when participants anticipate a demanding search task, they develop a different advanced representation of a visually identical target stimulus compared to when they anticipate a non demanding search.

Key words: visual-search, template, preparatory, representation, fMRI, sensory-motor.

## 1 Introduction

In a complex visual scene the object to which we attend is not always the most intrinsically salient e.g. the brightest or largest. Rather the things to which we attend are more often those that are relevant to our current goals and interests. For example, we can find our keys on the desk amongst the clutter or our car amongst many others in a large car park. These illustrate the general case of having something of current importance 'in mind' and seeking for that precise visual information in a cluttered visual world. This ability has been empirically studied with 'Visual search', an experimental paradigm that simulates these conditions (e.g. Treisman and Gelade 1980). In this, participants are asked to decide whether a 'target' such as a specific colored letter is present or not among a display of many similar items. Despite aspects of visual search being studied for over 30 years, the cognitive neuroscience of the formation of the advanced representation of the target is poorly understood.

The global network of areas involved in visual search tasks has been well documented in functional imaging studies (e.g. Anderson et al. 2010; Donner et al. 2000; Kim et al. 2012; Leonard et al. 2000; Nobre et al. 2003). The most consistently activated areas include superior parietal cortex, intraparietal sulcus, and occipital cortex along with various parts of frontal cortex (Anderson et al. 2007). However, understanding what aspect of visual search is performed by which part of this network of regions remains undetermined. Specifically, the design of these earlier studies have not allowed the 'prepare to search' and the 'search' element of the task to be separated. In the first, people have to

develop and maintain some adequate representation of the item to be found and in the second they have to match incoming stimuli against this.

From other lines of work, reasonable expectations can be formed as to the brain areas that may be involved in preparing to search for a target. A short term description of currently relevant visual information is often thought to be implemented by biasing feature maps in extra-striate regions of occipital cortex (Desimone and Duncan 1995). Consistent with this, sustained activation over the posterior scalp has been shown as people hold representations of targets for which they are about to search (Carlisle et al, 2011). In the human brain, feature maps for shapes and colors seem to exist in ventral occipital cortex (Beauchamp et al. 1999; Corbetta et al. 1990, 1991; Shulman et al. 1999, 2003). Furthermore, it has been demonstrated that such maps are activated when people are preparing to process visual input to find a target (Chawla et al. 1999, Giesbrecht et al. 2003). For example, Giesbrecht et al. (2003) showed increased bilateral activation in the fusiform region when people are waiting to make an orientation decision about a soon to be displayed colored, rectangle. It is clear that in visual search some similar advanced specification of the target must be formed and 'held in mind' prior to any search. The task must proceed by comparing multiple items in a visual display against this representation. It seems likely therefore that in visual search, the advanced specification of the target will also be found to be implemented here.

Importantly, as human cognition is highly flexible it seems likely that the advanced specification of target identity will vary with the current task demands. A target could be identical in two search tasks, but its advanced representation is predicted to be simpler when the upcoming task is expected to be undemanding compared to when it is expected to be demanding. For example the pre-biasing that might occur when preparing to find a red X is likely to be different when the task is expected to be a 'feature' search i.e. all distracters will

be green Os, relative to when the task is expected to be a 'conjunction' search i.e. distracters will be green Xs and red Os. In the first, a simple representation will suffice to perform the task. This could be implemented neurologically by the detection of any activation in feature maps coding other than green or O or by activation in red or diagonal feature maps. In contrast when the distracters will be green Xs and red Os, a more elaborated representation of the target including its relationship with the distracters must be formed (Duncan and Humphreys, 1989). If so, it is likely that for the identical stimulus, when preparing for such an undemanding feature search there will be less neural activation than when preparing for a demanding conjunction search. This may be detected as a smaller fMRI signal.

In addition to the variable activity in feature maps that might be expected to be seen in occipital cortex, the 'preparing to search' phase of a visual search task is likely to include other regions that are involved in modulating this sensory activity. A frontal-parietal control system is often proposed (e.g. Desimone and Duncan 1995; Woldorff et al. 2004) that sends bias signals to feature maps in ventral occipital cortex. Supporting evidence has come from studies where a representation of a target location has to be developed. In this approach a symbolic cue is given that indicates the likely location of an upcoming target. Participants use this advanced representation to facilitate target detection when it occurs (Hopfinger et al. 2000, Woldorff et al. 2004) or simply attend to that location (Kastner et al. 1999). For example Kastner et al. (1999) asked participants to attend to one location and count the occurrence of one of four complex colorful images presented there. In such studies, during this 'attended' interval, activation is seen in both occipital cortex, consistent with the biasing of visual spatial maps, and frontal and parietal areas, possibly involved in biasing such maps. While somewhat variable across studies the frontal activation typically includes the frontal eye fields. In visual search studies frontal and parietal activation is also often reported,

however it is never clear whether this reflects the source of the bias signal or the attentional movements that are part of later search and match operations. In contrast to the frontal areas that are active in visual search which vary across studies, the parietal activation is highly consistent. An area near the posterior portion of the intraparietal sulcus is active (Donner et al. 2000; Leonards et al. 2000; Nobre et al. 2003). In addition, Shulman et al. (1999) reported increased activation here when participants were maintaining information during an interval, regarding movement direction. Similarly, Giesbrecht et al. (2003) identified a region that includes a similar parietal area as responsible for the representation of task relevant information concerning colored shapes and location. This area has also been reported to be involved in other visual short term memory tasks (McNab and Klingberg, 2008; Todd and Marois, 2004). It seems likely that this area may be involved in maintaining the advanced specification of the target during 'prepare to search' as part of a frontal-parietal control system (Desimone & Duncan 1995).

No study has explicitly isolated the network of areas that support the development and maintenance of an advanced representation of the target in visual search from the other components of the task. Therefore, no study has been able to explore whether neural activation when preparing to search for a target, differs as a function of the anticipated demand of the task. The present study aims to address these limitations by separating the brain activation during the preparation to search for a target, from the later components of a visual search task. By isolating this time period, the changes in neural activity that might underlie the flexible creation of advanced specifications is investigated. This is done by presenting participants with identical targets but in contexts that indicate that their search will be undemanding or demanding, i.e. a feature search or a classic conjunction search. To minimize the interpretative processes that symbolic indication of the current target and


distracter information would have produced, spatial, shape and color information is given in a very concrete way, see Figure 1.

## **2 Results**

### 2.1 Reaction time results

Reaction time results are shown in Figure 2. There was a main effect of the 'type of search' factor with feature search being faster than conjunction search,  $F(1, 15) = 110.29, p < .001$ .

There was a main effect of the 'presence of the target' factor with 'target present' being faster than 'target absent',  $F(1, 15) = 22.47, p < .001$ . There was a significant interaction between the two factors,  $F(1, 15) = 22.80, p < .001$ . As shown in Figure 2, there was little lengthening of reaction time when the target was absent in feature search but a substantial increase when it was absent in conjunction search. Error rates were low in feature search (1.37%) and higher in conjunction search (13.82%). Any trial with an incorrect response was excluded from the subsequent fMRI analysis.



**Fig 2**

Reaction time of correct responses when the target is present and when it is absent in Feature search and Conjunction search conditions. Vertical lines show the standard error of means

## 2.2 Results - fMRI (preparation)

*Effects of preparation for a feature search - “Attend Prepare” (feature) versus “Watch” (feature).*

Full Talairach coordinates are given in Table 1 and Table 2. Preparing to perform a ‘Feature’ search resulted in four clusters of significant BOLD activity in extrastriate visual cortex, ventrally in right Inferior Occipital Gyrus and Fusiform Gyrus BA19, more dorsally in right



Middle Occipital Gyrus BA18, and more posteriorly and medially in the left Lingual Gyrus and right Cuneus (BA 17/18) in the Occipital pole [see Figure 3, top row].

TABLE 1

Furthermore, there were significant clusters of BOLD activity in left ventral Striatum and adjacent Anterior Insula, left posterior Thalamus, right anterior Lateral Cerebellum, and in the Pons.

TABLE 2


*Effects of preparation for a conjunction search - "Attend Prepare" (conjunction) versus "Watch" (conjunction).*

Results are shown in Table 1 and Table 2, Figure 3 middle row. Preparing to perform a conjunction search yielded significant BOLD activity in extrastriate visual cortex, including bilateral Inferior Occipital Gyri and Fusiform Gyri BA19 (ventrally), and more dorsally, bilateral Middle Occipital Gyri BA18/19, in the vicinity of the Transverse Occipital Sulcus (TOS). Additionally, there was significant BOLD activity in right Superior Parietal lobe (BA7), in the proximity of the Intraparietal Sulcus, [see Figure 3, middle row]. Finally, there were clusters in right posterior Thalamus, bilateral ventral Striatum, bilateral anterior Lateral Cerebellum, Midline cerebellum and Pons (Table 2).

*Selective effects of preparing for a feature search compared to preparing for a conjunction search - "Attend Prepare" (conjunction) versus "Attend Prepare" (feature)*

Regions where BOLD activity was greater when preparing for a conjunction relative to a feature search were bilateral ventral occipital cortex (Inferior Occipital/Fusiform Gyri BA19), bilateral dorsal occipital cortex (Middle Occipital Gyri BA18/19, in the proximity of TOS), and right Superior Parietal lobule BA7, in the vicinity of IPS [see Figure 3, bottom row]. Please

note that BOLD effects around the right Intraparietal Sulcus and in dorsal occipital cortex coincided across the two contrasts of “Attend Prepare” (conjunction) versus “Watch” (conjunction) and “Attend Prepare” (conjunction) versus “Attend Prepare” (feature) (Figure 3, bottom and middle rows and Table 1).

**Fig 3**

Group activation map (15 subjects), superimposed on one subject's rendered brain during 'preparing to search'. Superior view, left lateral, right lateral and posterior views are shown.

Top row, activations produced by preparing to search for a feature (Att\_Prep\_Feat) relative

to watching the control display (Watch\_Feat). Middle row, activations produced by preparing to search for a conjunction (Att\_Prep\_Conj) relative to watching the control display (Watch\_Conj). Bottom row, preparing to search for a conjunction compared to preparing to search for a feature. Abbreviations; R, Right; L, Left; GL, Lingual Gyrus; Fus, Fusiform; IPS, intraparietal Sulcus; TOS, Transverse occipital sulcus. Note: Top and Middle row used FWE corrected contrasts. Bottom row uses FDR corrected contrasts and a different scale.

### 3 Discussion

Fast event related fMRI was used to isolate the brain networks that are active during preparing to search for a visual target from the later components of a visual search task. The results show a distinctive network activated during the preparation phase of the task. Of particular interest was the way in which the representation of a given target would vary in anticipation of a 'feature' relative to a 'conjunction' search task. In preparing for either type of search, ventral occipital areas were activated, notably to a greater extent in advance of the conjunction search. In addition, when preparing for a conjunction search, unique activity was seen in bilateral dorsal occipital cortex and in the vicinity of the right intraparietal sulcus (Figure 3). Finally, preparing for either type of search activated ventral striatum, cerebellum, thalamus and pons.

#### 3.1 Varying representations with anticipated task demand

Activations seen in the ventral occipital region are consistent with the idea that an advance specification of the target is implemented by biasing feature maps in extra-striate regions of occipital cortex (Chawla, et al. 1999; Desimone and Duncan 1995; Giesbrecht et al. 2003, Stokes et al. 2009). This portion of extrastriate visual cortex is similar to that which, in previous studies, has been argued to encode both shape and color information (e.g. Beauchamp 1999; Shulman et al. 2003, 1999). Significantly, preparing to search for the identical target, but in the context of expecting it to be amongst very similar distracters rather than very different distracters produced different results. When expecting a conjunction search task, the same ventral occipital area was activated as when expecting a feature search task, but more robustly. This would be consistent with an interpretation that activity in this area reflects the formation of an advanced specification of the target and that

a more extensive representation is formed when participants anticipate a more demanding visual search. For example, when an easy feature search is expected, colour or shape maps may be activated, when a demanding conjunction search is expected colour and shape maps or more complex representations may be activated.

Target and distracter relationships may be established outside of the dorsal occipital region followed by signals sent back to bias early processing. However, at least part of the development of the advanced representation may arise directly out of local comparisons of target and distracter items during the 'preparation' display. During this phase of the task the 'target' needs to be compared to the distracters so as to prepare for the upcoming search task. In the second part of the task similar comparisons are made in the 'search' display. This repeated local comparison of items is similar to the local comparison of display items that is intrinsic in the inter-trial priming procedure (Müller et al. 1995, Found & Müller 1996, Pollmann et al. 2000). Processing on one trial alters the state of the perceptual system for the next search trial. For example, if a search is performed in the colour dimension, that dimension is altered so that it is processed faster on the next trial. This type of activity is thought to produce biasing or weighting of perceptual dimensions e.g. colour (Found & Müller 1996). Similar speeding effects are found when conjunction searches are repeated (Weidner et al. 2002), driven mainly by repeating the distractors in successive trials (Kristjánsson et al. 2002, Geyer et al. 2006,). We might therefore suspect that similar bottom-up biasing develops and stays active during the 'preparation' stage of the current task thus forming part of the preparatory set.

While, something akin to searching the display must have occurred i.e. comparison of target and distracters, this appears to have been done without overt eye movements. The fMRI data show no evidence of frontal eye field activation during this part of the task. As

increased activity in this region is reported to be a consistent finding in human neuroimaging studies (McDowell et al. 2008), the lack of it is indicative of the task being done without eye-movement. While sub-threshold activations cannot be excluded, given the clear instructions, that participants were practice on doing the task without eye movement outside the scanner and reported being able to do so, it seems probable that participants were largely successfully in following the instructions to fixate the central box

The idea of biased feature maps is consistent with a broad range of previous work it does not however fit well with the results of McMains et al. 2007. They demonstrated that there was a general increase in neural activity when preparing for a target event. However, this was non-specific e.g. in brain areas considered specialized for color processing, preparing for a color stimulus produced equal activation as preparing for a movement stimulus. This discrepancy could be understood in a number of ways. A key feature of the current study is that the target changed on every trial. This contrasts with the block design used in McMains et al. (2007) in which the same target identity is used over 18 second blocks of stimuli. It may be that the constant need to establish a new target representation creates a level of activation not seen when a single representation is formed and held. Alternatively, the results of Giesbrecht et al. 2003 indicate that somewhat different results are to be expected when stimuli are presented centrally (as in the current study) or more peripherally (as in McMains et al. 2007). They found that target specific preparatory activity (color or location) was seen with central presentations but this is less clearly the case with more peripheral presentations.

In addition to the ventral occipital areas, activity was observed in the vicinity of the transverse occipital sulci (TOS) when the upcoming visual search task was expected to be demanding. TOS may hence have a similar role as the ventral occipital regions in target

representation. Alternatively, the transverse occipital sulci may contribute to enhanced target representation by the suppression of distracters (Wokciulik and Kanwisher, 1999). Such an explanation seems particularly likely given the current design. Here for reasons of experimental control of visual saliency, during the preparation phase the target is always shown surrounded by distracters and when the upcoming task will be a conjunction search those distracters are always very similar to the target. Furthermore, the suppression of distracters has been shown to be a highly efficient way of biasing the perceptual system so that targets 'pop-out'. This is true for both feature and conjunction searches. One way in which a colour-form conjunction search could be done would be to by inhibiting one colour and then searching within the target colour for a unique shape (e.g. Treisman & Sato 1990). For example when the target is a red o among red xs and blue os, the colour (blue) could be suppressed and the search done on the shape dimension. In this case the unique rounded shape will pop-out from the straight oriented lines of the x distractors. Importantly, when this is used in one trial, it carries over to the next trial (Geyer et al. 2010), i.e. the inhibition remains active. In response to the current 'prepare screen' it seems likely that inhibition is developed in separating the target from distractors and maintained beyond the end of the prepare phase where it actively biases search during the latter part of the task. If so part of the activation we see, perhaps in the transverse occipital sulci may be due to active inhibition of one of the colours or shape maps. Similarly, relevant dimension weighting would be established during the preparation phase of 'feature' search. The features used were highly efficient 'redundantly defined targets' dissimilar from the surrounding distractors in both colour and shape. Search for such targets could be speeded by the active suppression of the non-target features, (Krummenacher et al. 2001, 2002). Neural activity



reflecting the development and maintenance of such suppression would be active during the 'preparation' stage.

It is worth noting that while distracter displays are identical in the Attend Prepare" versus "Watch" analyses they are not in the comparison of the two preparation conditions. This may contribute to the more robust activations seen in the latter analysis.

### 3.2 The posterior brain system

In addition to the occipital activations, a single right lateralized intraparietal activation was seen when preparing to perform a 'conjunction' search task, consistent with a large body of literature reporting activations in foci along the length of the intraparietal sulcus in similar tasks (e.g. Donner et al. 2000; Leonard et al. 2000). The current results restrict the IPS activation during preparation to a single focus. This focus (Figure 3, Table 1), corresponds closely to that described by Donner et al. (2000) as AIP (anterior intraparietal) and by Leonards et al. (2000) as MIPS (medial Intraparietal). It is also close to the location identified by Nobre et al. (2003) as involved in the overall demand of a search task.

Its role in preparing to search needs to be considered in the context of the lack of evidence of frontal activation during this part of the task. Until now it was never clear whether the frontal activations that were seen in earlier visual search studies reflected the representation of the target for which people were searching or some other aspect of the task. The current results provide an answer to this - at least for the concrete stimuli used here (i.e. 'target to be searched for' was indicated by a visually identical stimulus to the 'target presented'). Only posterior cortex and sub-cortical brain areas are found to be active when representing the target and preparing to respond to it in an up-coming visual search display. No evidence of frontal activation was seen. While a negative result, and therefore difficult to interpret, this would be consistent with the view that the biasing of perceptual

maps arises through local comparison of items in the prepare display and not through top-down control. It is possible that frontal cortex may have been engaged while learning the requirements of the task (during the instructions or the practice phase). However, for performance in the scanner the current results indicate that all aspects of the preparation to search are accomplished without frontal cortex involvement. This is consistent with a growing body of work that shows no evidence of frontal involvement in the building and maintenance of specific short-term representations of visual targets. For example Shulman et al. (1999) found no frontal activation while participants prepared to detect a specific direction of motion in a visual display. The results also dovetail with recent Event Related Potential (ERP) studies showing sustained activation over posterior scalp when people are maintaining a template of an item for which they are about to search, Carlisle et al. (2011). More broadly, it has been argued from recent neuroimaging work on visual working memory (see Postle, 2006) that frontal areas only become involved when transformation rather than memory per se, is required. While true for visual features such as motion, shape and color, an exception seems to be the advanced representation of visual stimuli at specific spatial locations. This has been robustly shown to activate frontal areas (e.g., Kastner et al. 1999, Woldorff et al. 2004) and may relate to the close connection between visual spatial attention and motor planning (Deubel & Schneider, 1996). The current results would bolster the position that apart from preparing to detect targets at a specific location, the advanced representation of target features is achieved outside of frontal cortex.

Theoretically this result is important because it is often assumed that in visual search a signal is being sent from cortical regions outside of the visual areas to bias feature maps in an appropriate and flexible way (Desimone and Duncan 1995). However, for the concrete visual stimuli used here, the traditional assumption that frontal-parietal networks are

necessarily involved in forming and maintaining a representation of a target is not supported. Given this, an alternative explanation for the role of the intra-parietal sulcus activation in preparing to search is required. While it is possible that the intra-parietal area by itself is involved in maintaining the advanced specification of the target during 'prepare to search' and sending bias signals to early visual cortex, this seems increasingly unlikely. Recent variants of the Todd and Marois (2004) visual short-term memory paradigm suggest that IPS activity is more related to the various attentional demands of tasks rather than any specific coding (Magen et al. 2009, Mitchell and Cusack 2010,). Magen et al. (2009) argue that attentional demands increase once the delay interval between the target memory display and the probe is lengthened so leading to an increase in activation in the Intra-parietal sulcus. This increased activation is non-specific, being found both in memory for visual information (colors) and spatial information. In the current study attentional demands were increased by changing the task from preparing to search for a feature to preparing to search for a conjunction of two features. It is possible that rather than sending content specific bias signals, the intra-parietal area may support the ongoing activation of occipital neurons that are already encoding target and distractor information. This attentional allocation might increase when a more complex representation needs to be maintained as in the current study or sustained for a longer period as in Magen et al. (2009). An alternative is that the intra-parietal area may be primarily receiving the output from the spatially precise and colour and form specialized occipital neurons, perhaps as part of a process of transforming the visual input into motor space as suggested by Ellison, et al. (2003). When participants anticipate a demanding search task a more detailed representation of the target and distracters may be implemented in these occipital regions. Their output may be what is reflected in increased activation in the intra-parietal sulcus.

### 3.3 Stimulus Response reassignment as target representations change

Concurrent activity in posterior cortex and striatum strongly suggest that the advanced representation of any target in visual search may be best considered as a visual-motor rather than a solely visual representation. In the present study, the identity of the target to which people should prepare to respond varied from trial to trial, thus the stimulus-response representation also changed on every trial. In addition, mixed amongst the 'prepare to search' trials were 'watch' trials which indicated that no response would be needed, in which case a switch from the previous stimulus-response representation would also have to be generated. This may be the processing that is being reflected as activity in the striatum. Such an interpretation would be consistent with earlier work e.g. Cooles et al. (2004). They explicitly examined the substrate of visual stimulus-response rule switching in the striatum and other areas. Participants were cued as to whether to respond to the same object as in the previous trial or to another object. Significant activation in the ventral striatum was found as participants switched between which of the two concrete (i.e. visually identical) objects to respond to. While in Cooles et al. (2004) the analysis was restricted to areas of interest whole brain analysis of a similar task has supported participation of the striatum while also demonstrating cerebellar involvement, Bischoff-Grethe et al. (2002). They explicitly contrasted template switching and response reassignment. They report right anterior-lateral cerebellum (lobule VI) activation during response reassignment, similar to that seen in the current results.

These studies suggests that in the current visual search task with its concrete visual targets, a likely function of ventral striatum and right anterior-lateral cerebellum is response reassignment to a visual stimulus, which is completed during the 'prepare to search' phase

of the task and is independent of task demand. The involvement of the striatum in maintaining a visual –motor template is plausible given that in non-human primates at least, there are substantial input-output connections from higher-order visual areas to the region around the caudate nucleus/ putamen and it has been linked to both perception and memory (Levy et al. 1997, Saint-cyr et al. 1990, Zink et al. 2003).

An alternative role for the putamen is suggested by the results of McNab & Klingberg (2007). They showed that increased activity in the left putamen was seen when participants had to actively ignore yellow colored discs rather than treating them as potential targets in a short term visual memory task. The current task also required ignoring distracters during the preparations stage and a similar function may be accomplished by the putamen here.

## 4 Experimental Procedures


### 4.1 Participants

Seventeen participants took part in the study (8 female, one left-handed, mean age 28.2,  $\pm$  7.89 years) with normal or corrected-to-normal vision. None admitted to current or past history of neurological or psychiatric conditions, learning disabilities, alcohol/substance abuse or current use of prescription medications (as ascertained through a medical history checklist). One subject was discarded for not achieving sufficient proficiency in the visual search task during the training session (see below), and a second was eliminated due to technical problems during the MRI session, yielding a final sample size of 15 subjects. The study was performed in agreement with the regulations of the University of British Columbia Behavioural Ethical Board. Participants took part in a behavioral session outside the scanner (45 min), where they had a chance to practice the visual search task until they exceeded a desired level of performance (>75% accuracy). This session took place within two days prior the fMRI session.

### 4.2 Task

The task was designed to avoid the order invariant problem and so enable the 'prepare to search' phase of a trial to be isolated from the later elements of the visual search task (Ollinger, Shulman & Corbetta, 2001; Ollinger, Corbetta Shulman & 2001). It involved having to decide whether a pre-defined target (a colored letter) was present or not amongst distracters (other colored letters). Visual stimuli were viewed through a periscopic mirror positioned about 10cm above the eyes of the participants. Throughout all trials a

central outline box was present in the middle of the display and participants were asked to keep their eyes fixed on this during a trial. The full sequence is shown in Figure 1.



**Fig 1**

Sequence of screens in a trial (top to bottom) with percent of different trial types. A Conjunction search trial is illustrated. In Feature search the target shared no features with the distracters.

Trials began with the 'Condition Display' in which the outline of the centrally positioned box turned blue or yellow. The colour instructed participants to either "Attend"

to (outline of the box turning blue) or simply “Watch” (outline of the box turning yellow) the upcoming display. This was used to inform the participants as to whether they could simply watch the display on the upcoming trial or should prepare themselves to perform a search task. Activity during the “Watch” condition was later subtracted from activity in the “Attend” conditions in order to control for brain activation caused by simply viewing rather than actively attending to the displays.

This initial ‘Condition Display’ was followed after 200 ms by the onset of a ‘Prepare Display’, which in the “Attend” conditions (“Attend Prepare-Only” trials and “Attend Prepare+Target” trials, see Figure 1) informed the subject as to the target and type of search to prepare for on that trial. The ‘Prepare Display’ was comprised of the target for which participants would shortly have to search, shown inside the central box, surrounded by the 31 or 32 distracters that could be present in the subsequent ‘Search Display’ (see Figure 1, columns 2 and 3). The equivalent display in the “Watch” trials was constructed in the same way except that the central square was filled with a ‘#’ symbol. The distracter sets were matched across conditions.

The ‘Prepare Display’ was presented for 800 ms, and was followed by the white central fixation box remaining on the screen for a further 1000 ms. After this in the “Watch” trials and “Attend Prepare-Only” trials (Figure 1, left and central columns), no further stimuli were presented. The central white outline box remained on the screen and trials ended following a variable interval (mean of 1850 ms, pseudo randomly jittered with a range of 800 - 2,900 ms). In “Attend Prepare+Target” trials however, a ‘Search Display’ followed the 1000 ms fixation and participants had to decide as quickly and accurately as possible if the target was present or absent in the display of 32 letters by pressing one of two keys on a fiber optic keypad with the index fingers of either hand. In the ‘Search Display’ the central



box was empty and the designated target could be either present, replacing one of the distracters that was in the 'Prepare Display' (target present 50%) or it could be omitted (target absent, 50%). The 'Search Display' was shown for 1000 ms and then replaced by a screen with just the white central box.

The target and distracter stimuli used to make the 'Prepare Displays' and the 'Search Displays', varied from trial to trial. The relationship between the target and the distracters determined whether a given visual search trial would be a feature or a conjunction search. In feature search trials, target and distracters had no feature in common (e.g. a yellow M amongst blue Ss). In conjunction search trials, as illustrated in Figure 1, the target and distracters always shared one feature [e.g. a yellow M, amongst yellow Ss (same color) and blue Ms (same shape)]. Equal numbers of feature and conjunction visual search trials were included. The same 'Prepare Displays' used in "Attend Prepare+Target" trials were used in the "Attend Prepare-Only" trials and in "Watch" control trials (but with the central target replaced by a '#'). To enable the isolation of the BOLD signal produced during the prepare phase from that produced by the target search phase, one third of trials were "Watch" trials, one third were "Attend Prepare-Only" and one third were "Attend Prepare+Target", (see, Ollinger, Corbetta & Shulman, 2001).

There were 3 runs of 196 trials. After every run, feedback was given in the form of mean reaction time and the number of their errors shown in the center of the screen for 30 seconds.

4.3 Image Acquisition - Echo-planar images were collected on a Philips Gyroscan Intera 3.0-T scanner, equipped with a 6-channel SENSE coil. Conventional spin-echo T1-weighted sagittal localizers were used to view head position and to graphically prescribe the functional image volumes. Functional image volumes sensitive to the blood oxygen-level dependent (BOLD)

contrast signal were collected with a gradient echo sequence (TR/TE 2000/30 ms, 90° flip angle, field of view 210×143×240 mm (anteroposterior, feet–head, right–left), 3 mm slice thickness, slice gap 1 mm, 36 axial slices).

4.4 Image processing - PAR/REC format data from the 3T Philips system were converted to Analyze format using MRICro (Rorden C: MRICro. <http://www.mricro.com>). The converted images were then analyzed using SPM5 (Wellcome Institute of Cognitive Neurology, <http://www.fil.ion.ucl.ac.uk/spm/>) for image reorientation, realignment, normalization into Montreal Neurological Institute space, and smoothing with a Gaussian kernel (8 mm full width at half maximum) to compensate for inter-subject anatomical differences and optimize the signal to noise ratio.

4.5 fMRI: within subjects Event-Related BOLD responses were modeled for the following trial types: “Watch” (feature), “Watch” (conjunction), “Attend Prepare” (feature) “Attend Prepare” (conjunction) “Attend Target” (feature) and “Attend Target” (conjunction) by the convolution of stimulus-onset vectors for each trial type with the synthetic hemodynamic response function provided in SPM2. The stimulus onset vectors coincided with ‘Condition Display’ onset for “Watch” and “Attend Prepare” trials and with ‘Search Display’ onset in “Attend Target” trials. Eight nuisance regressors (six sets of realignment parameters, and the mean signal from white matter and cerebro-spinal fluid voxels respectively) were included in the model. The magnitude of the BOLD responses for each trial type were calculated using the GLM implemented in SPM2.

To evaluate the selective effects of preparing to search for a target relative to passively looking at a display, the following contrast images were specified: Prepare for a

feature search – “Attend Prepare” (feature) versus “Watch” (feature), and prepare for a conjunction search - “Attend Prepare” (conjunction) versus “Watch” (conjunction). To evaluate the selective effects of preparing for a feature search compared to preparing for a conjunction search, the contrast “Attend Prepare” (feature) versus “Attend Prepare” (conjunction) was specified.

*4.6 fMRI: between subjects.* Contrast images for each subject were entered into two random effects analyses. Pair sample t-tests were set up to test the null hypotheses of no difference between trial types in the mean amplitude of the fitted hemodynamic response for any of these event types. We first applied the more conservative FWE method for correction of multiple comparisons,  $t(14) > 8.71$ ,  $p < .05$ , cluster-size  $> 10$ . This approach yielded several significant clusters for contrasts involving the lower control state (“Watch” trials). However, for the higher level contrast of “Attend Prepare” (feature) versus “Attend Prepare” (conjunction) no clusters reached significance at the .05 level. We then opted for selecting the more liberal FDR method for multiple comparison correction, with the statistical threshold set at  $t(14) > 4.6$ ,  $p < .05$ , cluster size  $> 20$ . This approach was indeed successful in yielding significant activation clusters for this contrast. Figure 3 illustrates the main results of these contrasts, highlighting the common regions activated in the contrasts involving “Attend Prepare” (conjunction) (middle and bottom rows). All reported coordinates are in Talairach space, following conversion from Montreal Neurological Institute (MNI) space, using the program *mni2tal* (Brett et al, 2001).

## 5 Conclusion

The current study identified a network of brain regions activated when preparing to search for a visual target embedded in a display of distracters. This was done by isolating it from the BOLD signal changes produced by the later components of the task. Target identity varied from trial to trial, requiring participants to form a new representation of the target on each trial. In addition, participants knew in advance how demanding the search was likely to be on a given trial. It was hypothesized that for an identical visual target, a simpler representation would be formed when the expected demand of the upcoming search task was low. It was expected that this would lead to a corresponding change in neural activity. The results show a network of neural areas activated in the posterior brain and in sub-cortical areas when 'preparing to search'. Importantly for the hypothesis when preparing to perform a demanding visual search task, identical targets produce new and additional neural activation in occipital and parietal areas. Future work will need to identify which attentional processes are involved in producing this pattern of result e.g. inhibition of distractors or activation of target representations, the relative involvement of the identified areas in different attentional processes and the extent of their involvement when the 'prepare display' is present relative to activity in the interval before the target display. Furthermore, to achieve a full understanding it will also be necessary to establish the directionality of effects and the timing of their activation during visual search. For the latter, fMRI effective connectivity analysis and methodologies with high spatiotemporal resolution (such as MEG) will be needed. These limitations notwithstanding, this is the first fast event-related fMRI study to identify neural correlates of the preparatory phase of visual search and their modulation by the anticipated demand of the visual search.

**Acknowledgments**

We thank Alan Woodford and Brian Luus for their assistance with programming and analysis, anonymous reviewers and Tim Hodgson for valuable feedback on earlier versions of the manuscript and Vince di Lollo for early discussions. This work was supported by the Royal Society London, International Short-term Overseas Visit grant (2004/R6) and The University of Lincoln to (PB) and by the Canada Foundation for Innovation and the National Science and Engineering Research Council of Canada to (ML).

## References

Anderson, E.J., Mannan, S.K., Husain, M., Rees, G., Sumner, P., Mort, D.J., McRobbie, D., Kennard, C., 2007. Involvement of prefrontal cortex in visual search. *Exp Brain Res.* 180, 289-302.

Anderson, E.J., Mannan, S.K., Rees, G., Sumner, P., Kennard C., 2010. Overlapping functional anatomy for working memory and visual search. *Exp Brain Res.* 200, 91-107.

Beauchamp, M.S., Haxby, J.V., Jennings, J.E., DeYoe, E.A., 1999. An fMRI version of the Frnsworth-Munsell 100-Hue test reveals multiple color-selective areas in human ventral occipitotemporal cortex. *Cereb Cortex.* 9, 257-263.

Bischoff-Grethe, A., Ivry, R., Grafton, S., 2002. Cerebellar Involvement in response reassignment rather than attention. *J Neurosci.* 22, 546–553.

Carlisle, N., Arita, J.T., Pardo, D., Woodman, G., 2011. Attentional templates in visual working memory, *J. Neurosci.* 31, 9315-9322.

Chawla, D, Rees, G., Friston, J.K., 1999. The physiological basis of attentional modulation in extrastriate visual areas. *Nat Neurosci,* 671–676.

Coolles, R., Clark, L., Robbins, T.W., 2004. Differential responses in human striatum and prefrontal cortex to changes in object and rule relevance. *J Neurosci.* 24, 1129-1135.

Corbetta, M., Miezin, F.M., Dobmeyer, S., Shulman, G.L., Petersen, S.E., 1990. Attentional modulation of neural processing of shape, color, and velocity in humans. *Science*. 248, 1556-1559.

Corbetta, M., Miezin, F.M., Dobmeyer, S., Shulman, G.L., Petersen S.E., 1991. Shape, color and speed: Functional anatomy by positron emission tomography. *J Neurosci*. 11, 2383-2402.

Desimone, R., Duncan, J., 1995. Neural mechanisms of selective visual attention. *Annu Rev Neurosci*. 18, 193-222.

Donner, T.H., Kettermann, A., Diesch, E., Ostendorf, F., Villringer, A., Brandt, S.A., 2000. Involvement of the frontal eye field and multiple parietal areas in covert visual selection during conjunction search. *Eur J of Neurosci*. 12, 3407-3414.

Deubel, H., Schneider, W., 1996. Saccade target selection and object recognition: evidence for a common attentional mechanism. *Vision Res*. 36, 1827-1839.

Duncan, J., Humphreys, G.W., 1989. Visual search and stimulus similarity, *Psychological Review*. 96, 433-458.

Ellison, A., Rushworth, M., Walsh, V., 2003. The parietal cortex in visual search: a visuomotor hypothesis. *Suppl Clin Neurophysiol*. 56, 321-33.

Found, A., Müller, H., 1996., Searching for unknown feature targets on more than one dimension: Investigating a “dimension-weighting” account. *Percept & Psychophys.* 58, 88-101.

Geyer, T., Müller, H., Krummenacher, J., 2006. Cross-trial priming in visual search for singleton conjunction targets: Role of repeated target and distractor features. *Percept & Psychophys.* 68, 736-749.

Geyer, T., Shi, Z., Müller, H., 2010. Contextual cueing in multiconjunction visual search is dependent on color- and configuration-based intertrial contingencies. *JEP:HPP.* 36, 515-532.

Giesbrecht, B., Woldorff, M.G., Song, A.W., Mangun, G.R., 2003. Neural mechanisms of top-down control during spatial and feature attention. *Neuroimage.* 19, 496-512.

Kim, K.K., Eliassen, J.C., Lee, S.K., Kang, E., (2012). Functional Neuroanatomy of visual search with differential attentional demands: An fMRI study. *Brain Research.* 1475, 49-61.

Kristjánsson, A., Wang, D., Nakayama, K., 2002. The role of priming in conjunctive visual search. *Cognition.* 85, 37-52.

Krummenacher, J., Müller, H., Heller, D., 2001. Visual search for dimensionally redundant pop-out targets: Evidence for parallel-coactive processing of dimensions. *Percept Psychophys.* 63, 901-917.



Krummenacher, J., Müller, H., Heller, D., 2002. Visual search for dimensionally redundant pop-out targets: Parallel-coactive processing of dimensions is location specific. *JEP:HPP*. 28, 1303-1332.

Leonard, U., Sunaert, S., van Hecke, P., Orban, G., 2000. Attention mechanisms in visual search – an fMRI study. *J Cogn Neurosci*. 12 supplement 2:61- 75.

Levy, R., Friedman, H.R., Davachi, L., Goldman-Rakic, P.S., 1997. Differential activation of the caudate nucleus in primates performing spatial and nonspatial working memory tasks. *J Neurosci*. 17, 3870-3882.

Magen, H., Emmanouil, T.A., McMains, S.A., Kastner, S., Treisman, A., 2009. Attentional demands predict short-term memory load response in posterior parietal cortex. *Neuropsychologia*. 47, 1790-1798

McDowell, J.E, Dyckman, K.A, Austin, B.P., Clementz, B.A., 2008. Neurophysiology and neuroanatomy of reflexive and volitional saccades: evidence from studies of humans. *Brain Cogn* 68: 255–271, 2008.

McMains S.A., Fehd H.M., Emmanouil, T, Kastner S., 2007. Mechanisms of feature and space-based attention: response modulation and baseline increases, *J Neurophysiol*. 98:2110-2121.

McNab, F., Klingberg, T., 2008. Prefrontal cortex and basal ganglia control access to working memory, *Nat Neurosci.* 11, 103-107.

Mitchell, D., Cusack, R., 2008. Flexible, capacity-limited activity of posterior parietal cortex in perceptual as well as visual short-term memory tasks. *Cereb Cortex.* 18, 1788-1798.

Müller, H., Heller, D., Ziegler, J., 1995. Visual search for singleton feature targets within and across feature dimensions. *Percept & Psychophys.* 57, 1-17.

Nobre, A., Coull, J., Walsh, V., Frith, C., 2003. Brain activations during visual search: Contributions of search efficiency versus feature binding. *Neuroimage.* 18, 91-103.

Ollinger, J.M., Shulman, G.L., Corbetta, M., 2001. Separating processes within a trial in event-related functional MRI. I. The method. *NeuroImage.* 13, 210–217.

Ollinger J.M., Corbetta, M., Shulman, G.L., 2001. Separating processes within a trial in event-related functional MRI. II. Analysis. *NeuroImage.* 13, 218–229.

Pollmann, S., Weidner, R., Müller, H., von Cramon, D., 2000. A fronto-posterior network involved in visual dimension changes. *J Cogn Neurosci,* 12, 480-494 .

Postle, B.R., 2006. Working memory as an emergent property of the mind and brain. *Neurosci.* 139, 23-38.

Saint-Cyr, J.A., Ungerleider, L.G., Desimone, R., 1990. Organization of visual cortical inputs to the striatum and subsequent outputs to the pallido-nigral complex in the monkey.

*J Compar Neurol.* 298, 129–156.

Shulman, G.L., Ollinger, J.M., Akbudak, E, Conturo, T.E., Snyder, A.Z., Petersen, S.F., Corbetta M., 1999. Areas involved in encoding and applying directional expectations to moving objects. *J Neurosci.* 19, 9480-9496.

Shulman, G.L., McAvoy, M.P., Cowan, M.C., Astafiev, S.V., Tansy, A.P., Avossa, G, Corbetta, M., 2003. Quantitative analysis of attention and detection signals during visual search. *J Neurophysiol* 90, 3384-3397.

Stokes, M., Thompson, R., Nobre, A.C., Duncan, J., 2009. Shape-specific preparatory activity mediates attention to targets in human visual cortex, *Proc R Soc.* 106, 19569–19574.

Treisman, A., Gelade, G., 1980 A feature-integration theory of attention. *Cognitive Psychol.* 12, 97-136.

Treisman, A., Sato, S., 1990. Conjunction search revisited. *JEP:HPP.* 16, 459-478.

Todd, J., Marois, R., 2004. Capacity limit of visual short-term memory in human posterior parietal cortex. *Nature.* 428, 751–754.

Weidner, R., Pollmann, S., Müller, H., von Cramon, D., 2002. Top-down controlled visual dimension weighting: an event-related fMRI study. *Cereb Cortex*.12, 318-328.

Woldorff, M., Hazlett, C., Fichtenholtz, H., Weissman, D., Dale, A., Song, A., 2004. Functional parcellation of attentional control regions of the human brain. *J. Cog. Neurosci*. 16, 149-165.

Wojciulik, E., Kanwisher, N., 1999. The generality of parietal involvement in visual Attention. *Neuron*. 23, 747-764.

Zink, C., Pagnoni, G., Martin, M., Dhamala, M., Berns, G., 2003. Human striatal response to salient non-rewarding stimuli. *J Neurosci*.23, 8092–8097.

