

1 **Title: Parental care influences social immunity in burying beetle larvae**

2

3 **Authors:** C.E. Reavey, L. Beare & S.C. Cotter

4 **Address:** School of Biological Sciences, Queen's University Belfast, MBC, 97 Lisburn Rd,

5 Belfast, BT9 7BL, UK

6 **Corresponding author:** Miss Catherine Reavey, School of Biological Sciences, Queen's

7 University Belfast, MBC, 97 Lisburn Rd, Belfast, BT9 7BL, UK. Tel: +44(0)2890972030,

8 Fax: +44(0)28 9097 5877. Email: [creavey01@qub.ac.uk](mailto:creavey01@qub.ac.uk)

9 **Keywords:** antibacterial, ecological immunology, insect, lysozyme, *Nicrophorus*, parental

10 care, social immunity

11 **Running title:** Social immunity in burying beetle larvae

12

## 13 **Abstract**

14 1. We provide evidence for social immunity in offspring of a sub-social species, the burying  
15 beetle, *Nicrophorus vespilloides*.

16 2. *N. vespilloides* is a carrion breeder, and in a similar fashion to the adult beetles, the  
17 offspring produce exudates that exhibit lytic activity, which are used to coat the breeding  
18 resource. This strategy defends against the microbial community.

19 3. The lytic activity in larval exudates declines as the brood develops, perhaps being most  
20 beneficial at the start of the breeding bout.

21 4. Changing levels of parental care through widowing/orphaning affects lytic activity in the  
22 larval exudates, with levels decreasing in the absence of both parents.

## 23 **Introduction**

24 The burying beetle, *N. vespilloides* (Figure S1, Supplementary Material), breeds on  
25 the carcass of small mammals and exhibits bi-parental care, one component of which is a  
26 social immune response (Cotter & Kilner 2010b). Cotter & Kilner suggest that “any type of  
27 immune response that has been selected to increase the fitness of the challenged individual  
28 and one or more recipients should be classified as social immunity” (2010b). The beetles  
29 coat the carcass with antimicrobial anal exudates (Cotter & Kilner 2010a) in order to  
30 minimise competition from the microbial community, delaying decomposition of the carcass  
31 (Rozen et al. 2008). This social immune response is costly; antibacterial activity is only  
32 upregulated in the presence of a carcass (Cotter & Kilner 2010a) and forced upregulation  
33 reduces lifetime reproductive success (Cotter et al. 2010). In *N. vespilloides*, both sexes  
34 produce exudates, however, the females’ exudates show higher levels of antibacterial activity

35 than the males (Cotter & Kilner 2010a). The beetles can also flexibly adjust the level of  
36 antibacterial activity in response to mate loss.

37 Observations have shown that the larvae also produce exudates throughout their  
38 development, though their potential antibacterial activity is unknown. With production of  
39 these exudates being so costly to the parents, it would benefit them if their offspring were  
40 also able to partake in carcass preservation. Furthermore, we know that larvae of this species  
41 can survive without parental care, albeit at a reduced rate and with a resulting poorer quality  
42 (Eggert et al. 1998). Their ability to survive the loss of both parents may be due in part to the  
43 production of antibacterial substances. Here we ask whether 1) antibacterial activity is  
44 present, 2) how it changes during larval development and 3) whether larvae, like adults, can  
45 flexibly alter levels of antibacterial activity in response to changing conditions, for example,  
46 presence/absence of parents.

## 47 **Materials and Methods**

### 48 ***N. vespilloides* colony**

49 The colony was established from a pedigreed *N. vespilloides* colony at the  
50 Department of Zoology, University of Cambridge. Beetles were maintained as described  
51 previously (Cotter & Kilner 2010a).

### 52 **Experiment 1 – characterising antibacterial activity across the larval stage**

53 Exudates were collected from all larvae in a brood using a capillary tube and pooled  
54 in a single eppendorf tube. This enabled us to use a known volume of exudate in the later  
55 analysis (1 µl). Larvae were sampled from day 1 (hatching) to day 5 (dispersal). Day 1 and 2  
56 larvae were so small that exudate collection with a capillary tube was not possible. Instead,  
57 the tip of the abdomen was gently pressed against a single punched circle of filter paper for

58 every larva in the brood. Whilst volume was impossible to control for in these samples, we  
59 collected <1 µl of exudate sample on each filter paper circle, and so the level of lytic activity  
60 would be a slight under-estimate compared to day 3, 4 and 5 larvae. All samples were stored  
61 at -20°C until testing. Larvae were sampled from fifty-five families, though some were  
62 sampled on more than one day, giving 75 samples in total. Forty-two families were sampled  
63 from once, nine families were sampled from twice and five families were sampled from three  
64 times. The numbers of larvae from which samples were collected and pooled on each day  
65 were as follows: day 1 = 10.33 larvae +/-2.01, day 2 = 14.63 larvae +/- 1.49, day 3 = 9.65  
66 larvae +/-1.24, day 4 = 14.27 larvae +/-1.88 and day 5 = 7.17 larvae +/- 2.29.

## 67 **Experiment 2 – the effect of reduced parental care on larval antibacterial activity**

68 Sixty pairs were established and were assigned to one of four treatment groups, male  
69 removed, female removed, both removed, or neither removed (control). On the day of  
70 hatching, parents were removed according to the treatment and the larval exudate sampled  
71 daily as described above. Exudate collection was more successful for this experiment, sixty  
72 families were sampled daily for five days giving 300 samples in total. Breeding success was  
73 recorded for all families.

## 74 **Antibacterial activity**

75 Lysozyme-like antibacterial activity was measured as described previously (Cotter &  
76 Kilner 2010a). In brief, 1 µl of each sample was pipetted into a hole in an agar plate  
77 inoculated with freeze-dried *Micrococcus lysodeikticus* cells and incubated for 24 hours at  
78 33°C. We selected *Micrococcus lysodeikticus* as it is a soil bacterium, which is the breeding  
79 environment of the burying beetle. It is one of the microorganisms analysed in Hall et al.  
80 (2011) and in that study was inhibited by secretions from *Nicrophorus*. It is also the main  
81 bacteria used for lytic plates in studies on burying beetles (Cotter et al. 2010, Steiger et al.  
82 2011, Arce et al. 2012) and many other insect species e.g. locusts (Wilson et al. 2002). Filter

83 paper circles were placed directly onto the surface of the agar. Clear zones in the agar were  
84 measured with digital calipers and calibrated against a lysozyme standard (Figure S2,  
85 Supplementary Material). In order to test for the potential mechanical differences between  
86 using filter paper and punched holes, we compared samples tested using both methods and  
87 found no difference between the size of the clear zone produced by 1  $\mu$ l of sample pipetted  
88 into a hole and that produced using filter paper placed on the surface of the agar ( $F_{1,25} = 0.27$ ,  
89  $P = 0.61$ ). We were therefore able to reliably compare samples measured using the different  
90 techniques. Whilst other methods are now available to measure antimicrobial activity, the  
91 majority of analyses to date in this and many other insect species have been carried out using  
92 zones of inhibition on agar plates. Using this method allows us to compare across studies to  
93 some extent.

#### 94 **Statistical analyses**

95       Antibacterial activity was log-transformed to approximate normality. Lytic data from  
96 both experiments was analysed with Restricted Estimate Maximum Likelihood (REML)  
97 models in Genstat 15 (VSN International, Hemel Hempstead, UK) with family included as a  
98 random effect to account for multiple testing from each family. Breeding data were analysed  
99 using a generalised linear model (GLM). Carcass weight and interaction terms were included  
100 in all models but were removed due to non-significance. Figures were produced in R 2.15.1  
101 (Development Core Team, 2013).

## 102 **Results and Discussion**

103       We show for the first time in this study that the larval exudates contain antibacterial  
104 substances. It has been shown previously that insect larvae have the ability to produce  
105 antibacterial secretions e.g. blowfly larvae, *L. sericata* (Kerridge et al. 2005), however this  
106 strategy has not been widely documented. Levels of lytic activity were highest in the newly-

107 hatched larvae and declined throughout the brood (REML:  $F_{1,49} = 51.27$ ,  $P < 0.001$ ; Figure 1).  
108 Indeed, as the exudate volume collected from day 1 and day 2 larvae was  $< 1 \mu\text{l}$ , the lytic  
109 activity at this stage is actually slightly underestimated, and so in reality should be even  
110 higher than observed. If the trend had been in the other direction i.e. larval exudate activity  
111 increasing with age, we could not have drawn reliable conclusions as in this case a lower  
112 activity in day 1 and day 2 larvae could have been due to a smaller exudate volume. As we  
113 did not collect a fixed amount of exudate from day 1 and 2 larvae (although volume  $< 1 \mu\text{l}$ ),  
114 we tested whether activity was dependent on the number of larvae from which it was  
115 collected, however there was no effect (REML:  $F_{1,27} = 0.51$ ,  $P = 0.48$ ). The pattern of decline  
116 in larval lytic activity mirrors that of the parents; highest when larvae arrive on the carcass  
117 (Cotter et al. 2013). The need for lytic exudates may fall after a high amount of initial  
118 preservation, resulting in increasing sterility. For the parents, assistance from the larvae may  
119 mean that they do not need to invest as heavily in lytic activity, which could promote further  
120 reproductive success and longevity.

121         With regards parental removal experiments, preliminary data exploration showed that  
122 the effects of removing males and females were very similar but different from the other two  
123 treatment groups. Therefore we grouped these two treatments into a single treatment  
124 representing “one parent removed”. Lytic activity was much lower in larvae where both  
125 parents were removed (REML:  $F_{2,46} = 3.66$ ,  $P = 0.033$ ; Figure 2a) and there was a significant  
126 effect of brood age, with the pattern being very similar to experiment 1 (REML:  $F_{1,210} =$   
127  $119.95$ ,  $P < 0.001$ ). These results illustrate that, like the parents (Cotter & Kilner 2010a), the  
128 larvae exhibit plasticity with regards lytic activity levels. Whilst we hypothesised that this  
129 antibacterial activity may contribute to their survival in orphaned conditions, activity actually  
130 decreased. This drop may be due to the fact that the larvae must now invest in other  
131 activities, for example self-feeding, at the detriment to lytic investment. Their condition is

132 also likely to be compromised even at this stage, which may cause a decline in lytic activity.  
133 A different experimental set-up may be required to unveil an upregulation of lytic activity in  
134 the larvae, for example using a rotten carcass but maintaining parental assistance i.e. a greater  
135 requirement for combatting microorganisms but without the strain of self-feeding. The  
136 number of larvae produced was lower when both parents were removed post-hatching (GLM:  
137  $F_{2,51} = 4.30$ ,  $P = 0.019$ ; Figure 2b), but the mean weight of larvae was not affected (GLM:  
138  $F_{2,49} = 2.07$ ,  $P = 0.137$ ). This is consistent with findings from previous studies (Rozen et al.  
139 2008; Arce et al. 2012).

140 Future experiments should consider the cost of lytic activity from the larval  
141 perspective. The larvae may be constrained developmentally if the mechanism for  
142 production in the parents also drives production in the larval stage. In light of plasticity from  
143 parents and their offspring, future studies should consider the scope for conflict. For  
144 example, if the larvae didn't produce antibacterial exudates, would the parents be forced to  
145 produce more? To conclude, we show that larvae produce antibacterial exudates and that  
146 parental care influences the extent of social immunity.

## 147 **Acknowledgements**

148 CER was supported by a DEL studentship, SCC was supported by a NERC  
149 fellowship (NE/H014225/2). Author contributions: SCC co-designed the experiments,  
150 analysed the data, and co-wrote the paper; LB co-designed the experiments and collected the  
151 data; CER co-designed the experiments, collected the data and co-wrote the paper.

152

153 **References**

- 154 Arce, A. N., Johnston, P. R., Smiseth, P. T. & Rozen, D. E. 2012. Mechanisms and fitness  
155 effects of antibacterial defences in a carrion beetle. *Journal of Evolutionary Biology*, **25**, 930-  
156 937.
- 157 Cotter, S. C. & Kilner, R. M. 2010a. Sexual division of antibacterial resource defence in  
158 breeding burying beetles, *Nicrophorus vespilloides*. *Journal of Animal Ecology*, **79**, 35-43.
- 159 Cotter, S. & Kilner, R. 2010b. Personal immunity versus social immunity. *Behavioral*  
160 *Ecology*, **21**, 663-668.
- 161 Cotter, S., Topham, E., Price, A. & Kilner, R. 2010. Fitness costs associated with mounting a  
162 social immune response. *Ecology Letters*, **13**, 1114-1123.
- 163 Cotter, S. C., Littlefair, J. E., Grantham, P. J. & Kilner, R. M. 2013. A direct physiological  
164 trade-off between personal and social immunity. *Journal of Animal Ecology*, **84**, 846-853.
- 165 Eggert, A. K., Reinking, M. & Müller, J. K. 1998. Parental care improves offspring survival  
166 and growth in burying beetles. *Animal Behaviour*, **55**, 97-107.
- 167 Hall, C.L. et al. (2011) Inhibition of Microorganisms on a Carrion Breeding Resource: The  
168 Antimicrobial Peptide Activity of Burying Beetle (Coleoptera: Silphidae) Oral and Anal  
169 Secretions. *Environmental Entomology*, **40**, 669-678.
- 170 Kerridge, A., Lappin-Scott, H. & Stevens, J. 2005. Antibacterial properties of larval  
171 secretions of the blowfly, *Lucilia sericata*. *Medical and veterinary entomology*, **19**, 333-337.

172 Rozen, D. E., Engelmoer, D. J. P. & Smiseth, P. T. 2008. Antimicrobial strategies in burying  
173 beetles breeding on carrion. *Proceedings of the National Academy of Sciences*, **105**, 17890-  
174 17895.

175 Steiger, S. et al. (2011) Sex differences in immunity and rapid upregulation of immune  
176 defence during parental care in the burying beetle, *Nicrophorus orbicollis*. *Functional*  
177 *Ecology*, **25**, 1368-1378.

178 Wilson, K. et al. (2002) Coping with crowds: Density-dependent disease resistance in desert  
179 locusts. *PNAS*, **99**, 5471-5475.

180

181

182

## 183 **Figure legends**

### 184 **Figure 1**

185           The decline in the antibacterial activity of larval exudate over time. Larvae hatch on  
186 day 1 and disperse from the carcass on day 5.

187


### 188 **Figure 2**

189           The effect of removing one or both parents on a) the antibacterial activity of larval  
190 exudate and b) the number of larvae dispersing from the carcass. Means and SEs in a) are  
191 predicted values from a REML model controlling for family.

192 **Figures**

193 **Figure 1**

194




195

196

197 **Figure 2**


198

199 a)



200

201 b)



202

203