

Accepted Manuscript

Preparation of Azide Biosynthetic Surrogates of *myo*-Inositol

Sandip Pasari, Shareef M. Ismail, Markus R. Wenk, Martin J. Lear

PII: S0040-4039(15)00623-1
DOI: <http://dx.doi.org/10.1016/j.tetlet.2015.04.009>
Reference: TETL 46146

To appear in: *Tetrahedron Letters*

Received Date: 23 January 2015
Revised Date: 29 March 2015
Accepted Date: 2 April 2015



Please cite this article as: Pasari, S., Ismail, S.M., Wenk, M.R., Lear, M.J., Preparation of Azide Biosynthetic Surrogates of *myo*-Inositol, *Tetrahedron Letters* (2015), doi: <http://dx.doi.org/10.1016/j.tetlet.2015.04.009>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.


Graphical Abstract

To create your abstract, type over the instructions in the template box below.
Fonts or abstract dimensions should not be changed or altered.

**Preparation of Azide Biosynthetic
Surrogates of *myo*-inositol**

Leave this area blank for abstract info.

Sandip Pasari, Shareef M. Ismail, Markus R. Wenk, Martin J. Lear*





Tetrahedron Letters
journal homepage: www.elsevier.com

Preparation of Azide Biosynthetic Surrogates of *myo*-Inositol

Sandip Pasari^a, Shareef M. Ismail,^b Markus R. Wenk^{b,c}, Martin J. Lear*^{a,d}

^a Department of Chemistry and Medicinal Chemistry Program, Faculty of Science, Science Drive 3, National University of Singapore, Singapore 117543

^b National University of Singapore (NUS) Graduate School for Integrative Sciences and Engineering (NGS), Centre for Life Sciences, 28 Medical Drive, Singapore 117456

^c Department of Biochemistry and Department of Biological Science, Yong Loo School of Medicine, National University of Singapore, Singapore 117543

^d School of Chemistry, University of Lincoln, Brayford Pool, Lincoln, Lincolnshire, LN6 7TS, UK

ARTICLE INFO

Article history:

Received

Received in revised form

Accepted

Available online

Keywords:

Inositols

Deoxy sugars

Biosynthesis

Glycolipids

ABSTRACT

As a prelude to biomolecular incorporation studies, practical routes to a series of four regioisomeric azido-deoxy derivatives of inositol that mimic the natural *myo*-stereochemistry are described. Starting from commercially available *myo*-inositol, the regioselective and stereoselective introduction of azide functionality was achieved at the C-2, C-3, C-4 and C-5 positions via azide displacement of the corresponding *O*-sulfonates of suitably protected *scyllo*-, *chiro*-, *epi*- and *neo*-inositols, respectively. Notably, a final one-pot acetolysis method conveniently allowed for rapid access to pentaacetate azido-deoxy inositols. Investigations on the metabolic incorporation of these *myo*-inositol azide surrogates in both acetate and free alcohol forms are in progress.

2009 Elsevier Ltd. All rights reserved.

Due to its putative prebiotic history and innate chemical versatility, inositol is a natural carbohydrate central to a plethora of cellular signaling processes and recognition events, both inside and outside cells, and even between different cell types.¹ Typically through initial phosphorylation events, inositol derivatives (e.g., inositol triphosphates, IP₃, and lipidated phosphatidylinositols, PIs) play key roles in modulating calcium signaling, membrane functioning, cell death, cell division and cell-to-cell communication.^{2,3} Additional glycosylation and lipidation events eventually yield higher order glycolipids, such as the glycosylphosphatidylinositols (GPIs) and poly-mannosylated lipids, including the phosphatidyl-*myo*-inositol mannosides (PIMs) and lipoarabinomannans (LAMs). These glycolipids are essential components in anchoring functional proteins on extracellular membrane surfaces and intracellular trafficking processes, and are key to host-cell recognition, immunological processes, and pathogenic infections.⁴

In this study, we have a view to prepare biosynthetic mimics of *myo*-inositol (azide-surrogates) that have the potential to be metabolically incorporated into a chosen cell type (e.g., yeast). For this purpose, the modification or introduction of bioorthogonal functional groups ("click" functionalities) was necessary. Herein, we chose to introduce azide groups to replace the different hydroxyl groups of *myo*-inositol (**1**) (Figure 1). For symmetry reasons, this would result in four regioisomeric series of *myo*-cyclohexanols **2-5**, of which two (the 2-series and 5-series) would be *meso* (achiral).





Figure 1. Azido-deoxy surrogates (**2-5**) of *myo*-inositol (**1**).

A few routes to the synthesis of optically active azido and amino inositols have been reported. Examples of routes start from *p*-benzoquinone⁵ or conduritol-E⁶ via chemo-enzymatic resolution, and also from chiral sources such as L-quebrachitol.⁷ Among various synthetic protocols developed for the synthesis of *myo*-inositol intermediates and its analogues, commercially available *myo*-inositol **1** is the most commonly preferred starting material due to its low cost and pre-defined relative stereochemistry.⁸ For *myo*-inositol, positions C₁, C₃ and C₄, C₆ are equivalent and unsymmetrical protection leads to racemates. The chemical synthesis of optically active inositol analogues would thus necessitate the resolution of racemic inositol intermediates (chemically, enzymatically or *via* desymmetrization techniques)⁹ or by starting with an alternative chiral material (e.g., *via* the Ferrier carbo-cyclization of sugars).¹⁰ For our biological studies, the racemic azido-inositol series **3** and **4**, and the *meso* series **2** and **5**, were considered sufficient to test our hypothesis of metabolic selection and incorporation into live cells. Our synthetic routes to make **2-5** are described herein.

* Corresponding author. Tel.: +44-1522-837411; fax: +44-1522-837411; e-mail: mlear@lincoln.ac.uk

Synthesis of (\pm)-4-Deoxy-4-Azido *myo*-Inositols 4:

We first decided to pursue the dicyclohexylidene diketal protection approach reported by Angyal and co-workers¹¹ as a starting material over diacetals or diisopropylidene derivatives.^{12,13} This allows for the practical, differential hydroxyl protection of *myo*-inositol (**1**) as relatively stable *trans* and *cis* cyclohexylidenes that not only tolerate multi-step synthesis, but also result in reactivity differences between the two remaining hydroxyl groups. The 1,2,4,5-dicyclohexylidene *myo*-inositol **6** was isolated as a white solid in 22% yield after recrystallisation from 1:9 acetone/petroleum ether.¹⁴ The remaining two isomeric ketals (2,3,4,5- and 3,4,5,6-dicyclohexylidenes) can be converted to **6** in a sequence of partial deprotection-reprotection steps.^{11b} Although yields are low, this method is scalable to multi-grams and is convenient in practice.



Scheme 1. Synthesis of (\pm)-4-azido *myo*-inositol analogues **4a,b**. Reagents and conditions: i. Cyclohexanone, toluene/DMF (1:2), pTSA, 110 °C, 22%; ii. BaO, Ba(OH)₂·8H₂O, BnBr, DMF, 60%; iii. DMP, DCM, rt, 93%; iv. NaBH₄, EtOH, 0 °C, 76%; v. Tf₂O, Py, DCM 0 °C to rt, 73%; vi. **11**, AcOH, Ac₂O, 10% H₂SO₄ in Ac₂O, rt; vii. NaOMe, MeOH, rt, 99%.

Barium oxide and barium hydroxide chelation-mediated benzylation of **6** produced the monobenzylated *myo*-inositol derivative **7** as the major product in 60% yield¹⁴ (Scheme 1). The free alcohol at the C₆ position of the *myo*-inositol derivative **7** was subsequently oxidised under Dess-Martin periodinane (DMP) conditions to give the desired ketone **8** in excellent yield (93%). Stereoselective reduction of **8** by sodium borohydride produced the *epi*-alcohol **9**¹⁵ exclusively. The free C₆ alcohol of *epi*-inositol **9** was subsequently activated as its triflate **10** using trifluoromethane sulfonic acid anhydride in pyridine.¹⁶ In accordance with the report of Schlewer *et al.*,¹⁵ azide displacement of triflate **10** gave the desired *myo*-product **11** together with trace amounts of the enol ether side-product **12** via 1,2-*anti* elimination under the basic conditions.

After several methods were explored, global deprotection of **11** in the presence of the azide group was eventually achieved in one step by acetolysis¹⁷ using a mixture of sulphuric acid, acetic anhydride and acetic acid to afford the 4-deoxy-4-azido-*myo*-inositol pentaacetate **4a**^{5,6} in 65% yield. Minor amounts of the partially acetolysed product 3-benzyl-6-azido-*myo*-inositol tetraacetate **13** was also obtained, which could be transformed under the same conditions to yield **4a**. Further methanolysis in the presence of a catalytic amount of sodium methoxide afforded the 4-deoxy-4-azido-*myo*-inositol analogue **4b**. Single crystals of pentaacetate **4a**, from 1:1 hexane/ethyl acetate, confirmed the stereochemistry unambiguously by X-ray analysis (Figure 2).





Figure 2. X-ray structure of racemic pentaacetate 4-azide analogue **4a**.

Synthesis of *meso*-5-Deoxy-5-Azido *myo*-Inositols 5:

Starting with the dicyclohexylidene diol **6**, benzylation of the two remaining hydroxyl groups afforded the fully protected inositol derivative **14** in excellent yield¹⁸ (Scheme 2). The kinetically labile and slightly distorted *trans* ketal of the compound **14** was cleaved selectively by controlled acid hydrolysis using acetyl chloride in DCM/MeOH (3:1) to afford the diol **15** in 68% yield.¹⁹ The diol **15** was regioselectively benzylated under phase transfer catalyst (tetrabutyl ammonium hydrogen sulphate) conditions, resulting in a separable mixture of benzylated products **16** and **17**.²⁰ Due to difficulties to distinguish **16** from **17**, both alcohols were converted to their corresponding methylated derivatives **18a,b** for identification purposes. Subsequent removal of the *cis*-ketal unit and global benzylation of **18a,b** provided the fully protected inositol derivatives **20a** and **20b**, respectively. The ¹H and ¹³C NMR of the derived *meso*-compound (**20b**) clearly indicated the structure to be 5-*O*-methyl-1,2,3,4,6-penta-*O*-benzyl *myo*-inositol. Thus, the corresponding starting alcohol **17** was confirmed to possess the C₅-free alcohol for the synthesis of the desired 5-series azido-analogues **5**.



Scheme 2. Synthesis of *meso*-5-azido *myo*-inositol analogues **5a,b**. Reagents and conditions: i. NaH, DMF, BnBr 0 °C, 97%; ii. CH₃COCl, DCM, MeOH (3:1), 68%; iii. BnBr, Bu₄NHSO₄, DCM, 5% NaOH, reflux; iv. NaH, DMF, MeI, 0 °C to rt, 90%; v. DCM/MeOH (1:1), CH₃COCl, rt, 89%; vi. (i); vii. DMP, NaHCO₃, DCM, rt, 85%; viii. NaBH₄, EtOH, rt, 60%; ix. Tf₂O, Py, 0 °C, 71%; x. NaN₃, DMF, rt, 83%; xi. Ac₂O, AcOH, 10% H₂SO₄ in Ac₂O, 60 °C, 64%; xii. NaOMe, MeOH, rt, 98%.

Having identified the C₅-OH derivative, the C₅-stereochemistry of **17** was inverted by DMP oxidation followed by reduction of the ketone **21** to give the alcohol **22** in the *neo*-configuration (Scheme 2). In the reduction step, the original alcohol **17** was also formed in 36% yield, presumably due to similar sterics being encountered by the hydride reagent when approaching from either the α or β face in **21**. The alcohol **22** was then converted into its trifluoromethane sulfonate **23** by treatment with Tf₂O in pyridine. Next, the azide group was installed by a clean substitution in an S_N2 fashion with sodium azide to

regenerate the *myo*-configured product **24** in good yield. Interestingly, no elimination (enol) product was detected, presumably due to 1,3-diaxial sterics preventing E₂-elimination. Compound **24** was then subjected to exhaustive acetylation to form the 5-deoxy-5-azido-*myo*-inositol pentaacetate **5a**.¹⁷ In this case, an elevated temperature 60 °C was required for complete conversion to the pentaacetate **5a**.⁶ The isomeric, partially acetylated, benzyl ether products were also converted to **5a** under the same acetylation conditions. Deacetylation of **5a** was accomplished by methanolysis in the presence of catalytic amounts of sodium methoxide to yield 5-deoxy-5-azido-*myo*-inositol **5b**. Single crystals of compound **5a** from 1:1 hexane/ether confirmed the structure unambiguously by X-ray analysis (Figure 3).





Figure 3. X-ray structure of meso-pentaacetate 5-azide analogue **5a**.

Synthesis of (±)-3-Deoxy-3-Azido *myo*-Inositols 3:

Inspired by the report of Watanabe *et al.*^{16b} for an efficient S_N2 substitution of C₃-inositol triflates, the synthesis of the 3-series **3a,b** was studied by consecutive double substitution at the C₃ position of an inositol intermediate **26**, which was obtained from the previously synthesized inositol derivative **7** by subsequent methoxy methyl (MOM) ether protection to form the fully protected **25** and Pd-C hydrogenolysis (Scheme 3).²¹ The C₃-free alcohol **26** was treated with trifluoromethane sulfonic acid anhydride in pyridine to yield the corresponding triflate **27**, which was immediately reacted with potassium acetate in DMA to furnish the C₃-inverted *chiro*-acetate **28** in excellent yield (96%)^{16b} without elimination, presumably due to the less basic nature of the acyl anion as compared to azide species.

Deacetylation of *chiro*-**28** in the presence of catalytic amounts of sodium methoxide gave the axial alcohol **29** (Scheme 3). After completion, acidic resin was added to the reaction mixture to remove sodium ions by ion exchange. Unexpectedly, the *trans* ketal unit of **29** rearranged into a more stable *cis* C3/C4-ketal (**30**) by ketal migration under the slightly acidic conditions. Hence, a basic aqueous work up procedure using ethylacetate-water was employed while scaling up. The free axial alcohol in **29** was subsequently treated with triflic anhydride and pyridine to form the triflate **31**. Excess pyridine led to formation of an eliminated product (**35**) in minor amounts. Two equivalents of pyridine in dichloromethane, however, generated the trifluoromethane sulfonylated product **31** in 90% yield without elimination. On the other hand, the free axial alcohol of **29** could be smoothly mesylated in pyridine as the solvent to form **32** without elimination, presumably due to the lower leaving group aptitude of OMs as compared to OTf.²² The isolated triflate **31** was examined first. Treatment with sodium azide in DMF generated the *myo*-configured substitution product **33** in low yield (15%) together with a minor S_N1 substitution product **34** (10%) and the elimination product **35** in 56% yield.²³ In comparison, the mesylate derivative **32** failed to undergo azide displacement with sodium azide in DMF, even upon heating at 70 °C. Also, Mitsunobu reaction of the alcohol **29** in order to achieve direct azide displacement was unsuccessful.




Scheme 3. Synthesis of (±)-3-azido *myo*-inositol analogues **3a,b**. Reagents and conditions: i. MOMCl, DIPEA, 0 °C to rt, 87%; ii. EtOAc, THF, 20% Pd-C, H₂, 60%; iii. Tf₂O, Py, DCM, 0 °C, 78%; iv. KOAc, DMA, 70 °C, 96%; v. NaOMe, MeOH, rt, 97%; vi. (iii), 90%; vii. MeSO₂Cl, py, 94%; viii. NaN₃, DMF, rt; ix. **31**, TMSN₃, TBAF, THF, 43%.

We thus opted to optimise the azide displacement of triflate **31** by screening differing azide reagents and solvents: NaN₃/DMA, NaN₃/Me₂CO/H₂O, ⁿBu₄NN₃/DCM, TMSN₃/TBAF/THF. Eventually, the combination of excess TMSN₃ (8 equiv.) in the presence of (0.5 equiv.) of TBAF in THF medium provided *myo*-azide **33** in a 43% optimal yield.²⁴ To complete the 3-series, global acetylation of compound **33** formed the desired 3-deoxy-3-azido *myo*-inositol pentaacetate **3a**⁵ in moderate yield (Scheme 3) under mildly acidic conditions (2% H₂SO₄ in acetic anhydride). Treatment of **33** in dry, ethereal HCl further allowed all acid labile protecting groups to be cleaved in one step and 3-deoxy-3-azido-*myo*-inositol **3b** could be isolated cleanly as a white solid after an ether wash.

Synthesis of meso-2-Deoxy-2-Azido *myo*-Inositols 2:

Following the orthoformate protection approach of Kishi *et al.*²⁵, which allows for large differences in reactivity between equatorial and axial hydroxyl groups of inositol in terms of sterics and electronics,²⁶ a route to the 2-series of azide analogues was developed (Scheme 4). The symmetric dibenzyl *myo*-inositol orthoformate **37**²⁷ was prepared from the dibenzoate derivative **36** through a prolonged silver(I) oxide chelation mediated bis-benzylation²⁸ and aminolysis sequence. The equatorial free hydroxyl group of **37** was oxidised under DMP condition to give the ketone **38** in excellent yield. Next, stereoselective reduction of ketone **38** produced the inverted axial alcohol **39** in the *scyllo*-configuration exclusively.²⁵ The rigidity of the orthoformate unit presumably accounts for this highly stereoselective reduction step. The axially positioned free alcohol of **39** was then sulfonylated²² by treatment with methane sulfonyl chloride in pyridine to give **40** in excellent yield. At this stage, an S_N2 attack at C₂ of compound **40** was presumed to be sterically challenging. Hence, orthoformate cleavage of compound **40** was performed first, and the triol **41** was formed by mild methanolysis with pTSA.²⁹ Next, the mesylate derivative **41** was heated with sodium azide in DMF, which formed the *myo*-configured azide substituted product **42** in moderate yield, along with minor amounts of the epoxide **43** and some unreacted starting material **41**. No elimination product was identified in this case; however, the reaction required heating at 80 °C, which led to decomposition of some material. Final exhaustive acetylation of **42** completed the synthesis of the desired 2-deoxy-2-azido-*myo*-inositol pentaacetate **2a** in good yield. Clean methanolysis under

basic conditions regenerated the free hydroxyl groups to produce *meso* 2-deoxy-2-azido-*myo*-inositol **2b**.



Scheme 4. Synthesis of 2-azido *myo*-inositol analogues **2a,b**. Reagents and conditions: i. (a) Ag_2O , BnBr , DMF, rt, (b) $(\text{CH}_3)_2\text{CHCH}_2\text{NH}_2$, MeOH, reflux, 70%; ii. DMP, DCM, rt, 95%; iii. MeOH, THF, NaBH_4 , 87%; iv. $\text{CH}_3\text{SO}_2\text{Cl}$, py, 92%; v. *p*TSA, MeOH, 97%; vi. NaN_3 , DMF, 80 °C; vii. Ac_2O , AcOH, 15% H_2SO_4 in Ac_2O , 50 °C, 64%; viii. NaOMe, MeOH, 99%.

Summary

In this letter, we have described straightforward routes to various azido-deoxy inositol analogues through azide installation. As a common strategy, we adopted a consecutive $\text{S}_{\text{N}}2$ double-inversion approach on suitably protected inositol derivatives, which were derived via convenient oxidation-reduction sequences from *myo*-inositol. A final, global acetolysis step in the presence of the azide group (2-15% sulphuric acid in acetic anhydride/acid) enabled the direct and convenient synthesis of azido-inositol pentaacetate analogues **2a-5a**. Methanolysis by treatment with catalytic sodium methoxide subsequently provided the fully unprotected azido *myo*-inositol surrogates **2b-5b** cleanly. The routes are convenient and provide sufficient material for biological study. Having genuine azide-surrogates of *myo*-inositol in hand, a more concise and diversified strategy is under investigation. In the meantime, the metabolic incorporation of these modified azido inositol analogues into various inositol lipids of yeast cells are in progress, and lipid profiling and biosynthetic cell compatibilities will be reported in due course.

Acknowledgments

We thank the National University of Singapore for graduate Scholarships (to S.P. and S.M.I.). This work was funded (to M.R.W. and M.J.L.) by the Competitive Research Program (CRP) of the National Research Foundation (NRF) of Singapore (NRF-G-CRP 2007-04).

Supplementary data

Supplementary data associated with this article can be found, in the online version, at ...

References and notes

- Agranoff, B. W.; Bradley, R. M.; Brady, R. O. *J. Biol. Chem.* **1958**, *233*, 1077.
- Shears, S. B. *Mol. Cell. Biol.* **1998**, *14*, 36, 49.
- Hammond, G.; Thomas, C. L.; Schiavo, G. *Curr. Top. Microbiol.* **2004**, *282*, 177.

- (a) Paulick, M. G.; Bertozzi, C. R. *Biochemistry* **2008**, *47*, 6991; (b) Baker, G. R.; Billington, D. C.; Gani, D. *Bioorg. Med. Chem. Lett.* **1991**, *1*, 17.
- Podeschwa, M. A. L.; Plettenburg, O.; Altenbach, H. *J. Org. Biomol. Chem.* **2003**, *1*, 1919.
- (a) Sanfilippo, C.; Patti, A.; Piattelli, M.; Nicolosi, G. *Tetrahedron Asymm.* **1998**, *9*, 2809; (b) Serrano, P.; Llebaria, A.; Delgado, A. *J. Org. Chem.* **2005**, *70*, 7829.
- (a) Johnson, S. C.; Dahl, J.; Shih, T. L.; Schedler, D. J.; Anderson, L.; Benjamin, T. L.; Baker, D. C. *J. Med. Chem.* **1993**, *36*, 3628; (c) De Almeida, M. V.; Figueiredo, R. M.; Dos Santos, H. F.; Da Silva, A. D.; De Almeida, W. B. *Tetrahedron Lett.* **2001**, *42*, 2767.
- (a) Sureshan, K. M.; Ikeda, K.; Asano, N.; Watanabe, Y. *Tetrahedron Lett.* **2004**, *45*, 8367; (b) Riley, A. M.; Jenkins, D. J.; Potter, B. V. L. *Carbohydr. Res.* **1998**, *314*, 277; (c) Gurale, B. P.; Shashidhar, M. S.; Gonnade, R. G. *J. Org. Chem.* **2012**, *77*, 5801; (d) Murali, C.; Gurale, B. P.; Shashidhar, M. S. *Eur. J. Org. Chem.* **2010**, 755.
- (a) Bruzik, K. S.; Myers, J.; Tsai, M.-D. *Tetrahedron Lett.* **1992**, *33*, 1009; (b) Martin, S. F.; Josey, J. A.; Wong, Y.-L.; Dean, D. W. *J. Org. Chem.* **1994**, *59*, 4805; (c) Rudolf, M. T.; Schultz, C. *Liebigs Ann.* **1996**, 533 and references therein; (d) Mayer, T. G.; Schmidt, R. R. *Liebigs Ann.* **1997**, 859; (e) Suzuki, T.; Tanaka, S.; Yamada, I.; Koashi, Y.; Yamada, K.; Chida, N. *Org. Lett.* **2000**, *2*, 1137; (f) Riley, A. M.; Correa, V.; Mahon, M. F.; Taylor, C. W.; Potter, B. V. L. *J. Med. Chem.* **2001**, *44*, 2108; (g) Sureshan, K. M.; Yamasaki, T.; Hayashi, M.; Watanabe, Y. *Tetrahedron Asymm.* **2003**, *14*, 1771.
- (a) Estevez, V. A.; Prestwich, G. D. *J. Am. Chem. Soc.* **1991**, *113*, 9885; (b) Chen, J.; Feng, L.; Prestwich, G. D. *J. Org. Chem.* **1998**, *63*, 6511; (c) Collins, P.; Ferrier, R. In *Monosaccharides; Their chemistry and their roles in natural products*: John Wiley & Sons Ltd.: West Sussex, England, 1995; pp. 449.
- (a) Angyal, S. J.; Tate, M. E.; Gero, S. D. *J. Chem. Soc.* **1961**, 4116; (b) Angyal, S. J.; Irving, G. C.; Rutherford, D.; Tate, M. E. *J. Chem. Soc.* **1965**, 6662.
- (a) Angyal, S. J.; Macdonald, C. G. *J. Chem. Soc.* **1952**, 686; (b) Leiserow, L.; Rabinovi, D. *J. Chem. Soc. A* **1969**, 2367; (c) Gigg, J.; Gigg, R.; Payne, S.; Conant, R. *Carbohydr. Res.* **1985**, *142*, 132; (d) Desai, T.; Gigg, J.; Gigg, R.; Martinzamor, E.; Schnetz, N. *Carbohydr. Res.* **1994**, *258*, 135.
- Baker, G. R.; Billington, D. C.; Gani, D. *Tetrahedron* **1991**, *47*, 3895.
- (a) Vacca, J. P.; Desolms, S. J.; Huff, J. R.; Billington, D. C.; Baker, R.; Kulagowski, J. J.; Mawer, I. M. *Tetrahedron* **1989**, *45*, 5679; (b) Dreef, C. E.; Tuinman, R. J.; Lefeber, A. W. M.; Elle, C. J. J.; Vandermaer, G. A.; Vanboom, J. H. *Tetrahedron* **1991**, *47*, 4709.
- Ballereau, S.; Guedat, P.; Poirier, S. N.; Guillemette, G.; Spiess, B.; Schlewer, G. *J. Med. Chem.* **1999**, *42*, 4824.
- (a) Tagliaferri, F.; Wang, S. N.; Berlin, W. K.; Outten, R. A.; Shen, T. Y. *Tetrahedron Lett.* **1990**, *31*, 1105-1108; (b) Sureshan, K. M.; Ikeda, K.; Asano, N.; Watanabe, Y. *Tetrahedron* **2008**, *64*, 4072.
- (a) van Dijkum, E.; Danac, R.; Hughes, D. J.; Wood, R.; Rees, A.; Wilkinson, B. L.; Fairbanks, A. *J. Org. Biomol. Chem.* **2009**, *7*, 1097; (b) Cao, Y.; Okada, Y.; Yamada, H. *Carbohydr. Res.* **2006**, *341*, 2219.
- Ozaki, S.; Watanabe, Y.; Ogasawara, T.; Kondo, Y.; Shiotani, N.; Nishii, H.; Matsuki, T. *Tetrahedron Lett.* **1986**, *27*, 3157.
- (a) Lu, P. J.; Gou, D. M.; Shieh, W. R.; Chen, C. S. *Biochemistry* **1994**, *33*, 11586; (b) Wang, D. S.; Hsu, A. L.; Song, X. Q.; Chiou, C. M.; Chen, C. S. *J. Org. Chem.* **1998**, *63*, 5430.
- Jiang, C.; Moyer, J. D.; Baker, D. C. *J. Carbohydr. Chem.* **1987**, *6*, 319.
- Lin, G.; Tsai, M. D. *J. Am. Chem. Soc.* **1989**, *111*, 3099.
- Fabris, F.; Rosso, E.; Paulon, A.; De Lucchi, O. *Tetrahedron Lett.* **2006**, *47*, 4835.
- Johnson, S. C.; Dahl, J.; Shih, T. L.; Schedler, D. J. A.; Anderson, L.; Benjamin, T. L.; Baker, D. C. *J. Med. Chem.* **1993**, *36*, 3628.
- Anilkumar, G.; Nair, L. G.; Olsson, L.; Daniels, J. K.; Fraser-Reid, B. *Tetrahedron Lett.* **2000**, *41*, 7605.
- Lee, H. W.; Kishi, Y. *J. Org. Chem.* **1985**, *50*, 4402.
- Sureshan, K. M.; Shashidhar, M. S.; Praveen, T.; Das, T. *Chem. Rev.* **2003**, *103*, 4477.
- Praveen, T.; Shashidhar, M. S. *Carbohydr. Res.* **2001**, *330*, 409.
- Banerjee, T.; Srikantiah, S. M. *Tetrahedron Lett.* **1994**, *35*, 8053.
- Sureshan, K. M.; Das, T.; Shashidhar, M. S.; Gonnade, R. G.; Bhabhade, M. M. *Eur. J. Org. Chem.* **2003**, 1035.

- Practical routes to all azido-deoxy derivatives of *myo*-inositol are described.
- Regio- and stereo-controlled introduction of azide functionality were achieved.
- Key *scyllo*-, *chiro*-, *epi*- and *neo*-inositols were prepared as intermediates.
- A final, one-pot acetolysis gave convenient access to the pentaacetate forms.
- Metabolic, biosynthetic incorporation of these azide surrogates are in progress.

ACCEPTED MANUSCRIPT