

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-(Hydroxymethyl)pyrene

 Tobias Gruber,[‡] Wilhelm Seichter and Edwin Weber*

Institut für Organische Chemie, TU Bergakademie Freiberg, Leipziger Strasse 29, D-09596 Freiberg/Sachsen, Germany

Correspondence e-mail: edwin.weber@chemie.tu-freiberg.de


Received 19 December 2009; accepted 19 January 2010

 Key indicators: single-crystal X-ray study; $T = 153$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.038; wR factor = 0.111; data-to-parameter ratio = 15.4.

The asymmetric unit of the title compound, $\text{C}_{17}\text{H}_{12}\text{O}$, contains two molecules, in which the fused aromatic ring systems are almost planar [maximum deviations = 0.0529 (9) and 0.0256 (9) Å]. In the crystal, aromatic π - π stacking interactions (perpendicular distance of centroids of about 3.4 Å) and strong $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds result in a helical arrangement of pyrenyl dimers.

Related literature

For the solid-state structures of pyrenes, see: Robertson & White (1947); Camerman & Trotter (1965); Allmann (1970); Hazell *et al.* (1972); Kai *et al.* (1978); Frampton *et al.* (2000). For the synthesis and structures of pyrene derivatives, see: Steward (1960); Gruber *et al.* (2006, 2008, 2009). For the use of pyrenes in fluorescence sensors, see: Bren (2001).



Experimental

Crystal data

 $\text{C}_{17}\text{H}_{12}\text{O}$
 $M_r = 232.27$
 Monoclinic, $P2_1/c$
 $a = 19.9182$ (6) Å

 $b = 8.8880$ (3) Å
 $c = 13.0882$ (4) Å
 $\beta = 91.719$ (2)°
 $V = 2316.00$ (13) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.08$ mm⁻¹
 $T = 153$ K
 $0.59 \times 0.29 \times 0.12$ mm

Data collection

 Bruker APEXII CCD area-detector diffractometer
 29632 measured reflections

 5051 independent reflections
 3801 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.111$
 $S = 1.06$
 5051 reflections

 327 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.21$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.18$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1}\cdots\text{O1A}^i$	0.84	1.87	2.6972 (12)	167
$\text{O1A}-\text{H1A}\cdots\text{O1}^{ii}$	0.84	1.89	2.7163 (12)	167

 Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINTE* (Bruker, 2004); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

The authors are indebted to F. Eissmann for his swift assistance. Financial support from the German Federal Ministry of Economics and Technology (BMW) under grant No. 16IN0218 'ChemoChips' is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2186).

References

- Allmann, R. (1970). *Z. Kristallogr. Kristallgeom. Kristallphys. Kristallchem.* **132**, 129–151.
 Bren, V. A. (2001). *Russ. Chem. Rev.* **70**, 1017–1036.
 Bruker (2004). *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Camerman, A. & Trotter, J. (1965). *Acta Cryst.* **18**, 636–643.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Frampton, C. S., Knight, K. S., Shankland, N. & Shankland, K. (2000). *J. Mol. Struct.* **520**, 29–32.
 Gruber, T., Fischer, C., Felsmann, M., Seichter, W. & Weber, E. (2009). *Org. Biomol. Chem.* **7**, 4904–4917.
 Gruber, T., Seichter, W. & Weber, E. (2006). *Acta Cryst.* **E62**, o2569–o2570.
 Gruber, T., Seichter, W. & Weber, E. (2008). *Supramol. Chem.* **20**, 753–760.
 Hazell, A. C., Larsen, F. K. & Lehmann, M. S. (1972). *Acta Cryst.* **B28**, 2977–2984.
 Kai, Y., Hama, F., Yasuoka, N. & Kasai, N. (1978). *Acta Cryst.* **B34**, 1263–1270.
 Robertson, J. M. & White, J. G. (1947). *J. Chem. Soc.* pp. 358–368.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Steward, F. H. L. (1960). *Aust. J. Chem.* **13**, 478–487.

[‡] Current address: University of Oxford, Department of Chemistry, Chemistry Research Laboratory, Mansfield Road, Oxford OX1 3TA, England.

supplementary materials

Acta Cryst. (2010). E66, o443 [doi:10.1107/S1600536810002424]

1-(Hydroxymethyl)pyrene

T. Gruber, W. Seichter and E. Weber

Comment

Owing to their electronic, optical and geometric properties, monofunctionalized pyrenes, attachable to a receptor platform, are of special interest for fluorescent sensor development (Bren, 2001). In this respect, 1-(hydroxymethyl)pyrene was prepared as part of our studies on the solid state structure of fluorogenic calixarenes with possible analytical applications (Gruber *et al.*, 2008; Gruber *et al.*, 2009).

Being composed of a plane aromatic region and a methylene bridged hydroxy group, the hybrid nature of the title compound is striking. The pyrene moiety alone shows no significant deviations of bond lengths and angles compared with those of the unsubstituted analogue (Robertson & White, 1947; Camerman & Trotter, 1965; Allmann, 1970; Hazell *et al.*, 1972; Kai *et al.*, 1978), and is almost planar. The largest deviation from the mean plane through the carbon framework of the pyrene unit is observed for atoms C2 [0.0529 (9)Å] and C1A [0.0256 (9)Å], respectively. Similar to the unsubstituted parent substance, the pyrene moieties of two molecules of 1-(hydroxymethyl)pyrene are forming a slightly displaced face-to-face dimer with an average distance of the aromatic units of about 3.4Å, though the latter are not arranged entirely coplanar [2.43 (3)°]. Additionally, within the dimer a strong hydrogen bond involving the two hydroxy groups can be observed [$d(\text{O}\cdots\text{O}) = 2.6972$ (12)Å]. Worth mentioning is the varying conformation of the hydroxymethyl residue in both molecules of the asymmetric unit. In molecule 1, a nearly coplanar arrangement with regard to the aromatic plane can be observed [C2–C1–C17–O1 = 3.46 (15)°], whereas in molecule 2 the same torsion angle of 116.54 (12)° is adopted (Fig. 1). These findings are explained by the steric demands of a strong hydrogen bond between two hydroxy groups [$d(\text{O}\cdots\text{O}) = 2.7163$ (12)Å], which links the pyrene dimers mentioned above in a helical manner in the direction of the crystallographic *b* axis. Considering the packing, two of these helices, each in the opposite direction, are connected by edge-to-face interactions of the pyrenyl groups as shown in Fig. 2.

Experimental

The title compound was synthesized from commercially available pyrene-1-carbaldehyde, which was reduced with sodium borohydride in boiling methanol, following an analogous procedure described for the reduction of anthracene-9-carbaldehyde (Steward, 1960; Gruber *et al.*, 2006). Colourless plates (m.p. 393–394 K) of the solvent-free 1-(hydroxymethyl)pyrene suitable for X-ray diffraction were obtained by recrystallization from *n*-hexane/dichloromethane (1:2).

Refinement

The H atoms were positioned geometrically and allowed to ride on their parent atoms, with O–H = 0.84Å, C–H = 0.95–0.99Å and $U_{\text{iso}} = 1.2\text{--}1.5 U_{\text{eq}}(\text{parent atom})$.

Figures




Fig. 1. Molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at 30% probability level. H atoms are presented as a small circles of arbitrary radius.



Fig. 2. Packing diagram of the title compound, viewed down the *a* axis. Hydrogen atoms not involved in hydrogen bonding have been omitted.

1-(Hydroxymethyl)pyrene

Crystal data

$C_{17}H_{12}O$

$M_r = 232.27$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 19.9182\ (6)\ \text{\AA}$

$b = 8.8880\ (3)\ \text{\AA}$

$c = 13.0882\ (4)\ \text{\AA}$

$\beta = 91.719\ (2)^\circ$

$V = 2316.00\ (13)\ \text{\AA}^3$

$Z = 8$

$F(000) = 976$

$D_x = 1.332\ \text{Mg m}^{-3}$

Melting point: 393 K

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 8740 reflections

$\theta = 2.5\text{--}32.3^\circ$

$\mu = 0.08\ \text{mm}^{-1}$

$T = 153\ \text{K}$

Plate, colourless

$0.59 \times 0.29 \times 0.12\ \text{mm}$

Data collection

Bruker APEXII CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube
graphite

φ and ω scans

29632 measured reflections

5051 independent reflections

3801 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\text{max}} = 27.0^\circ$, $\theta_{\text{min}} = 3.0^\circ$

$h = -25 \rightarrow 25$

$k = -11 \rightarrow 10$

$l = -16 \rightarrow 11$

Refinement

Refinement on F^2

Least-squares matrix: full

Primary atom site location: structure-invariant direct
methods

Secondary atom site location: difference Fourier map

$$R[F^2 > 2\sigma(F^2)] = 0.038$$

$$wR(F^2) = 0.111$$

$$S = 1.06$$

5051 reflections

327 parameters

0 restraints

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0591P)^2 + 0.2704P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.21 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.18 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.53989 (4)	-0.02633 (10)	0.21742 (7)	0.0412 (2)
H1	0.5295	-0.0899	0.2617	0.062*
C1	0.65534 (6)	0.05248 (12)	0.26246 (8)	0.0262 (2)
C2	0.63001 (6)	0.15072 (13)	0.33402 (8)	0.0316 (3)
H2	0.5828	0.1582	0.3409	0.038*
C3	0.67218 (6)	0.23825 (13)	0.39567 (8)	0.0336 (3)
H3	0.6534	0.3036	0.4447	0.040*
C4	0.74142 (6)	0.23177 (12)	0.38682 (8)	0.0296 (3)
C5	0.78651 (7)	0.32180 (13)	0.44853 (9)	0.0392 (3)
H5	0.7688	0.3864	0.4989	0.047*
C6	0.85288 (7)	0.31708 (14)	0.43697 (10)	0.0444 (3)
H6	0.8812	0.3781	0.4794	0.053*
C7	0.88234 (7)	0.22197 (14)	0.36198 (10)	0.0388 (3)
C8	0.95132 (7)	0.21558 (18)	0.34778 (12)	0.0540 (4)
H8	0.9807	0.2747	0.3899	0.065*
C9	0.97759 (7)	0.1249 (2)	0.27363 (13)	0.0606 (5)
H9	1.0247	0.1236	0.2645	0.073*
C10	0.93612 (7)	0.03560 (18)	0.21215 (11)	0.0494 (4)
H10	0.9550	-0.0256	0.1609	0.059*
C11	0.86674 (6)	0.03471 (14)	0.22482 (9)	0.0334 (3)
C12	0.82208 (6)	-0.05924 (13)	0.16581 (9)	0.0334 (3)
H12	0.8400	-0.1256	0.1167	0.040*
C13	0.75524 (6)	-0.05603 (12)	0.17800 (8)	0.0282 (3)
H13	0.7272	-0.1204	0.1374	0.034*

supplementary materials

C14	0.72512 (5)	0.04227 (11)	0.25087 (7)	0.0236 (2)
C15	0.76866 (6)	0.13372 (11)	0.31289 (8)	0.0250 (2)
C16	0.83916 (6)	0.12977 (12)	0.29987 (8)	0.0296 (3)
C17	0.60896 (6)	-0.04252 (14)	0.19512 (9)	0.0335 (3)
H17A	0.6153	-0.0146	0.1228	0.040*
H17B	0.6218	-0.1496	0.2032	0.040*
O1A	0.48228 (4)	0.26209 (9)	0.65830 (6)	0.0368 (2)
H1A	0.4998	0.3385	0.6859	0.055*
C1A	0.41705 (5)	0.19195 (13)	0.50620 (8)	0.0286 (3)
C2A	0.43258 (6)	0.08575 (14)	0.43237 (9)	0.0343 (3)
H2A	0.4781	0.0724	0.4148	0.041*
C3A	0.38356 (6)	-0.00083 (14)	0.38391 (8)	0.0337 (3)
H3A	0.3959	-0.0728	0.3342	0.040*
C4A	0.31636 (6)	0.01636 (12)	0.40716 (8)	0.0277 (2)
C5A	0.26394 (7)	-0.06959 (13)	0.35757 (8)	0.0341 (3)
H5A	0.2754	-0.1425	0.3080	0.041*
C6A	0.19914 (7)	-0.04986 (13)	0.37933 (9)	0.0360 (3)
H6A	0.1658	-0.1081	0.3442	0.043*
C7A	0.17913 (6)	0.05748 (13)	0.45456 (8)	0.0308 (3)
C8A	0.11218 (6)	0.08017 (15)	0.47954 (10)	0.0397 (3)
H8A	0.0779	0.0234	0.4455	0.048*
C9A	0.09507 (6)	0.18397 (16)	0.55308 (10)	0.0426 (3)
H9A	0.0492	0.1978	0.5688	0.051*
C10A	0.14416 (6)	0.26767 (15)	0.60382 (9)	0.0371 (3)
H10A	0.1317	0.3383	0.6543	0.045*
C11A	0.21194 (6)	0.24959 (12)	0.58179 (8)	0.0281 (3)
C12A	0.26408 (6)	0.33496 (13)	0.63165 (8)	0.0305 (3)
H12A	0.2526	0.4059	0.6826	0.037*
C13A	0.32911 (6)	0.31781 (12)	0.60855 (8)	0.0286 (3)
H13A	0.3622	0.3777	0.6429	0.034*
C14A	0.34959 (5)	0.21105 (12)	0.53313 (8)	0.0245 (2)
C15A	0.29880 (5)	0.12329 (11)	0.48278 (7)	0.0232 (2)
C16A	0.23004 (5)	0.14309 (12)	0.50639 (8)	0.0250 (2)
C17A	0.47224 (6)	0.28837 (15)	0.55113 (9)	0.0370 (3)
H17C	0.5145	0.2669	0.5159	0.044*
H17D	0.4609	0.3956	0.5397	0.044*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0302 (5)	0.0379 (5)	0.0551 (6)	-0.0008 (4)	-0.0045 (4)	0.0141 (4)
C1	0.0342 (6)	0.0209 (5)	0.0234 (5)	0.0014 (5)	-0.0018 (4)	0.0054 (4)
C2	0.0356 (6)	0.0297 (6)	0.0297 (6)	0.0062 (5)	0.0051 (5)	0.0060 (5)
C3	0.0529 (8)	0.0252 (6)	0.0231 (6)	0.0098 (5)	0.0058 (5)	-0.0004 (5)
C4	0.0489 (7)	0.0184 (6)	0.0211 (5)	0.0018 (5)	-0.0035 (5)	0.0024 (4)
C5	0.0675 (9)	0.0225 (6)	0.0269 (6)	-0.0012 (6)	-0.0102 (6)	-0.0014 (5)
C6	0.0669 (10)	0.0280 (7)	0.0369 (7)	-0.0142 (6)	-0.0201 (6)	0.0039 (5)
C7	0.0451 (7)	0.0328 (7)	0.0377 (7)	-0.0109 (6)	-0.0120 (5)	0.0143 (5)

C8	0.0460 (8)	0.0608 (10)	0.0545 (9)	-0.0216 (7)	-0.0128 (7)	0.0200 (8)
C9	0.0329 (7)	0.0859 (12)	0.0627 (10)	-0.0102 (8)	-0.0017 (7)	0.0288 (9)
C10	0.0401 (8)	0.0603 (9)	0.0481 (8)	0.0083 (7)	0.0065 (6)	0.0171 (7)
C11	0.0344 (6)	0.0344 (7)	0.0316 (6)	0.0045 (5)	0.0012 (5)	0.0120 (5)
C12	0.0445 (7)	0.0297 (6)	0.0262 (6)	0.0108 (5)	0.0044 (5)	0.0020 (5)
C13	0.0392 (7)	0.0224 (6)	0.0229 (5)	0.0024 (5)	-0.0027 (4)	-0.0003 (4)
C14	0.0342 (6)	0.0176 (5)	0.0188 (5)	0.0018 (4)	-0.0016 (4)	0.0033 (4)
C15	0.0371 (6)	0.0177 (5)	0.0202 (5)	0.0015 (4)	-0.0030 (4)	0.0050 (4)
C16	0.0374 (7)	0.0244 (6)	0.0268 (6)	-0.0026 (5)	-0.0049 (5)	0.0102 (4)
C17	0.0322 (6)	0.0327 (7)	0.0351 (6)	-0.0021 (5)	-0.0036 (5)	0.0026 (5)
O1A	0.0400 (5)	0.0354 (5)	0.0342 (5)	-0.0092 (4)	-0.0105 (3)	0.0040 (4)
C1A	0.0319 (6)	0.0281 (6)	0.0257 (6)	-0.0012 (5)	-0.0023 (4)	0.0072 (4)
C2A	0.0324 (6)	0.0411 (7)	0.0297 (6)	0.0056 (5)	0.0043 (5)	0.0065 (5)
C3A	0.0468 (7)	0.0307 (6)	0.0236 (6)	0.0090 (5)	0.0033 (5)	-0.0016 (5)
C4A	0.0420 (7)	0.0206 (5)	0.0202 (5)	0.0009 (5)	-0.0029 (4)	0.0029 (4)
C5A	0.0568 (8)	0.0233 (6)	0.0219 (6)	-0.0028 (5)	-0.0059 (5)	-0.0013 (4)
C6A	0.0508 (8)	0.0282 (6)	0.0281 (6)	-0.0129 (6)	-0.0143 (5)	0.0037 (5)
C7A	0.0358 (7)	0.0281 (6)	0.0279 (6)	-0.0056 (5)	-0.0083 (5)	0.0104 (5)
C8A	0.0346 (7)	0.0413 (7)	0.0425 (7)	-0.0083 (6)	-0.0101 (5)	0.0161 (6)
C9A	0.0292 (6)	0.0506 (8)	0.0481 (8)	0.0017 (6)	0.0014 (5)	0.0206 (7)
C10A	0.0387 (7)	0.0384 (7)	0.0345 (6)	0.0106 (5)	0.0069 (5)	0.0104 (5)
C11A	0.0341 (6)	0.0257 (6)	0.0244 (5)	0.0044 (5)	-0.0002 (4)	0.0070 (4)
C12A	0.0430 (7)	0.0244 (6)	0.0239 (5)	0.0067 (5)	-0.0017 (5)	-0.0022 (4)
C13A	0.0383 (6)	0.0214 (6)	0.0257 (6)	-0.0009 (5)	-0.0080 (4)	-0.0012 (4)
C14A	0.0315 (6)	0.0197 (5)	0.0220 (5)	0.0001 (4)	-0.0032 (4)	0.0042 (4)
C15A	0.0330 (6)	0.0174 (5)	0.0190 (5)	0.0002 (4)	-0.0033 (4)	0.0036 (4)
C16A	0.0321 (6)	0.0205 (5)	0.0221 (5)	-0.0002 (4)	-0.0043 (4)	0.0072 (4)
C17A	0.0349 (6)	0.0411 (7)	0.0349 (7)	-0.0081 (6)	-0.0028 (5)	0.0093 (5)

Geometric parameters (Å, °)

O1—C17	1.4222 (14)	O1A—C17A	1.4301 (14)
O1—H1	0.8400	O1A—H1A	0.8400
C1—C2	1.3865 (15)	C1A—C2A	1.3922 (17)
C1—C14	1.4056 (15)	C1A—C14A	1.4094 (15)
C1—C17	1.5146 (15)	C1A—C17A	1.5000 (16)
C2—C3	1.3860 (17)	C2A—C3A	1.3824 (17)
C2—H2	0.9500	C2A—H2A	0.9500
C3—C4	1.3887 (17)	C3A—C4A	1.3900 (16)
C3—H3	0.9500	C3A—H3A	0.9500
C4—C15	1.4218 (15)	C4A—C15A	1.4232 (15)
C4—C5	1.4338 (16)	C4A—C5A	1.4334 (16)
C5—C6	1.3356 (19)	C5A—C6A	1.3415 (18)
C5—H5	0.9500	C5A—H5A	0.9500
C6—C7	1.434 (2)	C6A—C7A	1.4360 (17)
C6—H6	0.9500	C6A—H6A	0.9500
C7—C8	1.393 (2)	C7A—C8A	1.3971 (17)
C7—C16	1.4247 (16)	C7A—C16A	1.4233 (15)
C8—C9	1.377 (2)	C8A—C9A	1.383 (2)

supplementary materials

C8—H8	0.9500	C8A—H8A	0.9500
C9—C10	1.386 (2)	C9A—C10A	1.3825 (19)
C9—H9	0.9500	C9A—H9A	0.9500
C10—C11	1.3969 (18)	C10A—C11A	1.3983 (16)
C10—H10	0.9500	C10A—H10A	0.9500
C11—C16	1.4187 (17)	C11A—C16A	1.4216 (15)
C11—C12	1.4295 (17)	C11A—C12A	1.4281 (16)
C12—C13	1.3459 (16)	C12A—C13A	1.3476 (16)
C12—H12	0.9500	C12A—H12A	0.9500
C13—C14	1.4377 (15)	C13A—C14A	1.4371 (15)
C13—H13	0.9500	C13A—H13A	0.9500
C14—C15	1.4246 (14)	C14A—C15A	1.4235 (14)
C15—C16	1.4201 (16)	C15A—C16A	1.4241 (15)
C17—H17A	0.9900	C17A—H17C	0.9900
C17—H17B	0.9900	C17A—H17D	0.9900
C17—O1—H1	109.5	C17A—O1A—H1A	109.5
C2—C1—C14	119.64 (10)	C2A—C1A—C14A	119.25 (10)
C2—C1—C17	121.08 (10)	C2A—C1A—C17A	118.94 (11)
C14—C1—C17	119.28 (10)	C14A—C1A—C17A	121.74 (11)
C3—C2—C1	121.36 (11)	C3A—C2A—C1A	121.82 (11)
C3—C2—H2	119.3	C3A—C2A—H2A	119.1
C1—C2—H2	119.3	C1A—C2A—H2A	119.1
C2—C3—C4	120.94 (10)	C2A—C3A—C4A	120.66 (11)
C2—C3—H3	119.5	C2A—C3A—H3A	119.7
C4—C3—H3	119.5	C4A—C3A—H3A	119.7
C3—C4—C15	118.84 (10)	C3A—C4A—C15A	118.95 (10)
C3—C4—C5	122.47 (11)	C3A—C4A—C5A	122.37 (10)
C15—C4—C5	118.69 (11)	C15A—C4A—C5A	118.68 (10)
C6—C5—C4	121.60 (12)	C6A—C5A—C4A	121.78 (11)
C6—C5—H5	119.2	C6A—C5A—H5A	119.1
C4—C5—H5	119.2	C4A—C5A—H5A	119.1
C5—C6—C7	121.51 (11)	C5A—C6A—C7A	121.40 (10)
C5—C6—H6	119.2	C5A—C6A—H6A	119.3
C7—C6—H6	119.2	C7A—C6A—H6A	119.3
C8—C7—C16	118.77 (13)	C8A—C7A—C16A	118.86 (11)
C8—C7—C6	122.73 (12)	C8A—C7A—C6A	122.87 (11)
C16—C7—C6	118.50 (12)	C16A—C7A—C6A	118.27 (11)
C9—C8—C7	121.00 (13)	C9A—C8A—C7A	121.06 (12)
C9—C8—H8	119.5	C9A—C8A—H8A	119.5
C7—C8—H8	119.5	C7A—C8A—H8A	119.5
C8—C9—C10	120.81 (13)	C10A—C9A—C8A	120.51 (12)
C8—C9—H9	119.6	C10A—C9A—H9A	119.7
C10—C9—H9	119.6	C8A—C9A—H9A	119.7
C9—C10—C11	120.59 (14)	C9A—C10A—C11A	120.82 (12)
C9—C10—H10	119.7	C9A—C10A—H10A	119.6
C11—C10—H10	119.7	C11A—C10A—H10A	119.6
C10—C11—C16	118.93 (12)	C10A—C11A—C16A	119.09 (11)
C10—C11—C12	122.74 (12)	C10A—C11A—C12A	122.55 (11)
C16—C11—C12	118.32 (10)	C16A—C11A—C12A	118.36 (10)

C13—C12—C11	121.68 (11)	C13A—C12A—C11A	121.88 (10)
C13—C12—H12	119.2	C13A—C12A—H12A	119.1
C11—C12—H12	119.2	C11A—C12A—H12A	119.1
C12—C13—C14	121.70 (10)	C12A—C13A—C14A	121.56 (10)
C12—C13—H13	119.2	C12A—C13A—H13A	119.2
C14—C13—H13	119.2	C14A—C13A—H13A	119.2
C1—C14—C15	119.25 (9)	C1A—C14A—C15A	119.19 (10)
C1—C14—C13	122.99 (10)	C1A—C14A—C13A	122.93 (10)
C15—C14—C13	117.76 (10)	C15A—C14A—C13A	117.87 (10)
C16—C15—C4	119.72 (10)	C4A—C15A—C14A	120.12 (10)
C16—C15—C14	120.31 (10)	C4A—C15A—C16A	119.43 (10)
C4—C15—C14	119.95 (10)	C14A—C15A—C16A	120.44 (9)
C11—C16—C15	120.17 (10)	C11A—C16A—C7A	119.67 (10)
C11—C16—C7	119.86 (11)	C11A—C16A—C15A	119.90 (10)
C15—C16—C7	119.96 (11)	C7A—C16A—C15A	120.43 (10)
O1—C17—C1	113.64 (10)	O1A—C17A—C1A	111.75 (9)
O1—C17—H17A	108.8	O1A—C17A—H17C	109.3
C1—C17—H17A	108.8	C1A—C17A—H17C	109.3
O1—C17—H17B	108.8	O1A—C17A—H17D	109.3
C1—C17—H17B	108.8	C1A—C17A—H17D	109.3
H17A—C17—H17B	107.7	H17C—C17A—H17D	107.9
C14—C1—C2—C3	-0.83 (16)	C14A—C1A—C2A—C3A	0.59 (17)
C17—C1—C2—C3	-179.82 (10)	C17A—C1A—C2A—C3A	-176.38 (10)
C1—C2—C3—C4	0.83 (17)	C1A—C2A—C3A—C4A	0.44 (18)
C2—C3—C4—C15	0.19 (16)	C2A—C3A—C4A—C15A	-0.83 (16)
C2—C3—C4—C5	179.32 (10)	C2A—C3A—C4A—C5A	179.05 (10)
C3—C4—C5—C6	-178.12 (11)	C3A—C4A—C5A—C6A	-178.71 (11)
C15—C4—C5—C6	1.01 (17)	C15A—C4A—C5A—C6A	1.17 (16)
C4—C5—C6—C7	0.19 (18)	C4A—C5A—C6A—C7A	-0.76 (17)
C5—C6—C7—C8	179.54 (12)	C5A—C6A—C7A—C8A	-179.58 (11)
C5—C6—C7—C16	-0.77 (18)	C5A—C6A—C7A—C16A	-0.22 (16)
C16—C7—C8—C9	1.55 (19)	C16A—C7A—C8A—C9A	0.14 (17)
C6—C7—C8—C9	-178.76 (13)	C6A—C7A—C8A—C9A	179.50 (11)
C7—C8—C9—C10	-1.0 (2)	C7A—C8A—C9A—C10A	-0.13 (18)
C8—C9—C10—C11	-0.6 (2)	C8A—C9A—C10A—C11A	0.17 (18)
C9—C10—C11—C16	1.65 (19)	C9A—C10A—C11A—C16A	-0.23 (16)
C9—C10—C11—C12	-177.77 (12)	C9A—C10A—C11A—C12A	179.31 (11)
C10—C11—C12—C13	-178.69 (11)	C10A—C11A—C12A—C13A	-179.03 (10)
C16—C11—C12—C13	1.88 (16)	C16A—C11A—C12A—C13A	0.50 (16)
C11—C12—C13—C14	0.14 (17)	C11A—C12A—C13A—C14A	-0.75 (17)
C2—C1—C14—C15	-0.18 (15)	C2A—C1A—C14A—C15A	-1.18 (15)
C17—C1—C14—C15	178.83 (9)	C17A—C1A—C14A—C15A	175.69 (9)
C2—C1—C14—C13	-179.59 (10)	C2A—C1A—C14A—C13A	179.92 (10)
C17—C1—C14—C13	-0.58 (15)	C17A—C1A—C14A—C13A	-3.20 (16)
C12—C13—C14—C1	177.38 (10)	C12A—C13A—C14A—C1A	179.07 (10)
C12—C13—C14—C15	-2.04 (15)	C12A—C13A—C14A—C15A	0.15 (15)
C3—C4—C15—C16	177.55 (9)	C3A—C4A—C15A—C14A	0.21 (15)
C5—C4—C15—C16	-1.62 (15)	C5A—C4A—C15A—C14A	-179.68 (9)
C3—C4—C15—C14	-1.19 (15)	C3A—C4A—C15A—C16A	179.28 (9)

supplementary materials

C5—C4—C15—C14	179.65 (9)	C5A—C4A—C15A—C16A	-0.60 (14)
C1—C14—C15—C16	-177.54 (9)	C1A—C14A—C15A—C4A	0.80 (15)
C13—C14—C15—C16	1.90 (14)	C13A—C14A—C15A—C4A	179.75 (9)
C1—C14—C15—C4	1.18 (14)	C1A—C14A—C15A—C16A	-178.27 (9)
C13—C14—C15—C4	-179.38 (9)	C13A—C14A—C15A—C16A	0.68 (14)
C10—C11—C16—C15	178.59 (10)	C10A—C11A—C16A—C7A	0.24 (15)
C12—C11—C16—C15	-1.96 (15)	C12A—C11A—C16A—C7A	-179.32 (9)
C10—C11—C16—C7	-1.12 (16)	C10A—C11A—C16A—C15A	179.89 (9)
C12—C11—C16—C7	178.33 (10)	C12A—C11A—C16A—C15A	0.34 (15)
C4—C15—C16—C11	-178.65 (9)	C8A—C7A—C16A—C11A	-0.19 (15)
C14—C15—C16—C11	0.08 (15)	C6A—C7A—C16A—C11A	-179.58 (9)
C4—C15—C16—C7	1.05 (15)	C8A—C7A—C16A—C15A	-179.84 (9)
C14—C15—C16—C7	179.78 (9)	C6A—C7A—C16A—C15A	0.77 (15)
C8—C7—C16—C11	-0.46 (16)	C4A—C15A—C16A—C11A	180.00 (9)
C6—C7—C16—C11	179.84 (10)	C14A—C15A—C16A—C11A	-0.92 (15)
C8—C7—C16—C15	179.83 (10)	C4A—C15A—C16A—C7A	-0.35 (15)
C6—C7—C16—C15	0.13 (16)	C14A—C15A—C16A—C7A	178.73 (9)
C2—C1—C17—O1	-3.46 (15)	C2A—C1A—C17A—O1A	-116.54 (12)
C14—C1—C17—O1	177.53 (9)	C14A—C1A—C17A—O1A	66.57 (14)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1 \cdots O1A ⁱ	0.84	1.87	2.6972 (12)	167.
O1A—H1A \cdots O1 ⁱⁱ	0.84	1.89	2.7163 (12)	167.

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $x, -y+1/2, z+1/2$.

Fig. 1




Fig. 2

