

1 **Aposematism in the burying beetle? Dual function of anal fluid in parental**  
2 **care and chemical defence**

3 Carita Lindstedt<sup>1,2</sup>, Giuseppe Boncoraglio<sup>2</sup>, Sheena Cotter<sup>3,2</sup>, James Gilbert<sup>4,2</sup> and Rebecca M.  
4 Kilner<sup>2</sup>

5 1. Centre of Excellence in Biological Interactions, Department of Biological and  
6 Environmental Sciences, University of Jyväskylä, Finland

7 2. Department of Zoology, University of Cambridge, UK

8 3. School of Life Sciences, University of Lincoln, UK

9 4. School of Biological, Biomedical and Environmental Sciences, The University of Hull, UK.

10

11 *A short title: Aposematic display of burying beetle *N. vespilloides*?*

12

13

14 *Corresponding author:*

15 Carita Lindstedt

16 Centre of Excellence in Biological interactions,

17 Department of Biological and Environmental Sciences,

18 University of Jyväskylä, P.O. Box 35,

19 FI-40014 University of Jyväskylä, Finland

20 carita.a.lindstedt@jyu.fi

21

22 ABSTRACT

23 Burying beetles (*Nicrophorus vespilloides*) bear distinctive and variable orange-black  
24 patterning on their elytra and produce an anal exudate from their abdomen when threatened.  
25 During breeding, the anal exudates contribute to the antimicrobial defence of the breeding  
26 resource. We investigated whether the anal exudates also provide a responsive chemical  
27 defence, which is advertised to potential avian predators by the beetle's orange and black elytral  
28 markings. We found that that the orange-black elytral markings of the burying beetle are highly  
29 conspicuous for avian predators against range of backgrounds, by using computer simulations.  
30 Using bioassays with wood ants, we also showed that the burying beetle's anal exudates are  
31 aversive to potential predators. From these results, and other evidence in the literature, we  
32 conclude that the evidence for aposematism in the burying beetle is as strong as the evidence  
33 for many other classically aposematic species, such as defended Hymenopterans, ladybirds or  
34 poisonous frogs. Nevertheless, we also report unexpectedly high levels of individual variation  
35 in coloration and chemical defences, as well as sex differences. We suggest that this variation  
36 might be due partly to conflicting selection pressures, particularly on the dual function of the  
37 exudates, and partly to nutritional differences in the developmental environment. The ecology  
38 of the burying beetles (*Nicrophorus* spp.) differs markedly from better-studied aposematic  
39 insects. This genus thus offers new potential for understanding the evolution of aposematism  
40 in general.

41

42

43

44

45

46

47 INTRODUCTION

48 Prey individuals with toxic defences educate predators to avoid prey of similar appearance in  
49 future encounters (Speed et al. 2012). The avoidance learning rate of predators will be further  
50 enhanced if a defended prey bears a distinctive and memorable signal, such as bright coloration  
51 or a conspicuous display that predators can associate with the toxicity (i.e. aposematism) and  
52 so avoid attacking prey animals that carry that signal in future (Poulton 1890, Guilford 1990,  
53 Alatalo and Mappes 1996, Ruxton, Sherratt and Speed 2004). Predators have been shown to  
54 select for pronounced warning signals (Forsman and Merilaita 1999, Lindström et al. 1999,  
55 Lindstedt et al. 2008, Mappes et al. 2014) and signal uniformity (e.g., Mallet and Barton 1989,  
56 Joron and Mallet 1998, Kapan 2001, Beatty et al. 2004, Rowland et al. 2007) as well as high  
57 levels of chemical defence (Leimar et al. 1986, Skelhorn and Rowe 2006, Ihalainen et al. 2007,  
58 Rowland et al. 2007) because all these characteristics enhance the efficiency of avoidance  
59 learning in the predator. Therefore, directional selection by predators is expected to decrease  
60 variation in the expression of these traits.

61                 Nevertheless, it is widely acknowledged that both aposematic coloration (Ojala  
62 et al. 2007, Stevens and Ruxton 2011) and levels of chemical defence (Speed et al. 2012) can  
63 vary considerably among individuals. One explanation is that intrinsic constraints limit the  
64 response to directional selection from predators. For example, physiological costs of producing  
65 pigmentation (Grill and Moore 1998, Bezzerides et al. 2007, Ojala et al. 2007, Sandre et al.  
66 2007, Lindstedt et al. 2010) or defensive chemicals (Higginson et al. 2011) can maintain  
67 variation in each of these traits. These costs can be further shaped by ecological (Grill and  
68 Moore 1998, Bezzerides et al. 2007, Ojala et al. 2007, Sandre et al. 2007, Lindstedt et al. 2010)  
69 and social (Daly et al. 2012) environments. In addition, the heritability of an aposematic trait  
70 and how it is genetically correlated with other traits can also influence the way in which it  
71 responds to directional selection from predators, and is a measure of the extent of variation in  
72 that trait (Lindstedt et al. 2016).

73                 A different explanation for the persistence of variation is that aposematic  
74 coloration serves multiple functions, for example in thermoregulation (Brakefield 1985,  
75 Lindstedt et al. 2009, Hegna et al. 2013) or in mate choice (Summers et al. 1999, Maan and  
76 Cummings 2009). Thus, one of the key steps in understanding how this variation is maintained,  
77 has been to move the focus from the two-way interaction of the predator and prey towards  
78 considering the interactions of the prey species in greater complexity. This approach can  
79 identify additional selection pressures that may oppose directional selection imposed by  
80 predators, and thereby maintain variation in aposematic coloration (Friman et al. 2009,  
81 Nokelainen et al. 2011, Gordon et al. 2015, Rojas et al. 2015, Crothers and Cummings 2013).  
82 Likewise, defensive compounds can also serve multiple functions and consequently be  
83 subjected to selection in different directions. For example, defensive toxins sequestered from  
84 the diet can sometimes be used to enhance immunological defence against parasites (Laurentz  
85 et al. 2012, Kollberg et al. 2014) or to produce pheromones at reproductive stage (Conner et  
86 al. 1981). Therefore to understand how variation in aposematic displays persists, despite  
87 directional selection from predators, it is important to establish new independent model species  
88 that differ ecologically and are therefore exposed to diverse selection pressures.


89 Here we consider whether the burying beetle (*Nicrophorus vespilloides*) exhibits  
90 aposematism and describe the extent of individual variation in its chemical defences and  
91 putative aposematic coloration. Burying beetles (*Nicrophorus* spp) are carnivorous Silphid  
92 beetles that are best known for their elaborate biparental care (Scott 1998, Eggert et al. 1998).  
93 They prepare carrion during reproduction, which they defend, maintain and feed to their  
94 offspring. Larvae of burying beetles feed on the carcass which parents smear with foul smelling  
95 dark brown anal exudate (Degenkolb et al. 2011), inhibiting microbial growth (Cotter et al.  
96 2010, Cotter et al. 2013) and increasing larval survival (Arce et al. 2012).

97 The majority of *Nicrophorus* species also bear the distinctive orange-black  
98 coloration that is typical of other aposematic insects (Sillen-Tullberg 1985, Mappes and Alatalo  
99 1997, Gamberale-Stille and Tullberg 1999, Exnerová et al. 2006; Sikes et al. 2002, Figure 1a.).  
100 Several reports in the literature suggest that the orange-black elytral markings of the burying  
101 beetle could function as part of a warning display (Morton Jones 1932, Lane and Rothschild  
102 1965, Anderson and Beck 1985, Young 2014). Many Silphid beetles commonly feature in the  
103 diet of diverse vertebrates (Young 2014) and burying beetles specifically are potential prey for  
104 crows that scavenge upon carrion (Morton Jones 1932). Yet black Silphidae are more  
105 commonly described as prey than the orange and black *Nicrophorus* spp (Young 2014).  
106 Furthermore, Morton Jones (1932) reports that none of three different North American  
107 *Nicrophorus* spp were eaten by birds when presented alongside other Coleopteran species. The  
108 burying beetle species were unique among those species in being orange and black, whereas  
109 the species that were consumed were not. Further circumstantial evidence that the orange and  
110 black coloration of the burying beetle is aposematic comes from observations by Lane and  
111 Rothschild (1965), who describe a marked increase in agitation shown by captive blue tits  
112 (*Cyanistes caeruleus*) when orange-black *N. investigator* beetles were placed in their cages.  
113 These agitated behaviours are a characteristic avian response to several different species of  
114 aposematic insects (Rothschild and Lane 1960).

115 The orange-black colouration is just one component of a burying beetle's putative  
116 warning display. Upon handling, they also make a conspicuous 'buzzing' sound (Lane and  
117 Rothschild 1965, Hall et al 2013, C. Lindstedt pers obs). *N. investigator* even moves its  
118 abdomen in a style purported to resemble the stinging movements of bumble-bees (Lane and  
119 Rothschild 1965). These visual and auditory elements of the display accompany the responsive  
120 production of chemical defences. Upon handling, burying beetles produce the same anal  
121 exudate from their abdomen that is used by beetles to defend the carcass from rival microbes  
122 (Lane and Rothschild 1965, Cotter and Kilner 2010, Cotter et al. 2010, Degenkolb et al. 2011,  
123 Duarte et al. 2017). The odour of the exudate reportedly lingers for more than a year on  
124 unwashed 'inanimate objects' (Lane and Rothschild 1960), is very pungently putractive and  
125 has a very high pH (Degenkolb et al. 2011). In addition to compounds with antimicrobial  
126 properties, the anal exudate of *N. vespilloides* includes over 10 chemical compounds known to  
127 be repellent against invertebrates and vertebrates and some of these compounds can serve both  
128 antimicrobial and repellent functions (Degenkolb et al. 2011). Many of these repellent  
129 compounds have been found also in the defensive glands of other Coleopteran and  
130 Hymenopteran species (Degenkolb et al. 2011) suggesting that they could function in chemical

131 defence of the adult beetles as well as assist in defending the carcass from the rival microbes  
132 (Duarte et al. 2017). During the breeding chemical profile of the anal exudate changes as the  
133 number of antimicrobial compounds produced by *N. vespilloides* beetles increases. However,  
134 the repellent compounds are still present in the anal exudate during the breeding (Degenkolb  
135 et al. 2011, Haberer et al. 2014).

136



**Figure 1.** Individual variation in the aposematic signal for a) the size of the striking orange elytral pattern, and b) the quantity of anal exudate *N. vespilloides* produces when disturbed.

137

138 We have three aims in this paper: 1) to determine the salience of the burying  
139 beetle's orange and black coloration to avian predators, against a range of natural backgrounds  
140 (Stevens 2007); 2) to test whether the chemical defences in the burying beetle's anal exudates  
141 are aversive, using a standard bioassay with ants; 3) to quantify phenotypic variation and broad-  
142 sense heritability in each of these traits. Aims 1) and 2) are linked to understanding the nature  
143 of selection acting on the burying beetle's elytral markings and chemical defences, whereas  
144 aim 3) helps to understand how these traits might respond to selection.

145

## 146 METHODS

### 147 *N. vespilloides* colony

148 We used burying beetles from an outbred laboratory population established in 2005 at  
149 Cambridge University, and supplemented annually with wild-caught individuals from sites  
150 close to Cambridge, UK. Adults were housed alone in plastic boxes (12x8x2 cm) filled with  
151 moist soil, food (minced beef) was available *ad libitum* and boxes were kept at a constant  
152 temperature of 21 °C and 16h:8h light:dark cycle. Boxes were cleaned twice a week and at the  
153 same time old food was replaced. For breeding, unrelated pairs were placed in plastic boxes  
154 (17x12x6 cm) half filled with moist soil, provided with a freshly thawed mouse carcass (21.94  
155  $\pm$  0.33 SE g, range 15-35g) and kept in the dark. Larvae disperse from the carcass ca. 8 days  
156 after hatching and sexual maturity is reached ca. 5 weeks after dispersal.

157

### 158 **Aim 1: Quantifying the salience of the orange-black coloration to avian predators**

159 To test how insectivorous birds perceive the colour, luminance and contrast of colour patterns  
160 of beetles against various natural backgrounds, we used an avian vision model that assumes  
161 that receptor noise limits visual discrimination (Vorobyev and Osorio 1998, Vorobyev et al.  
162 1998). This model is included in the Image Calibration and Analysis Toolbox (Troscianko and  
163 Stevens 2015). First, the regions of interest (ROIs) from the normalized and linearized images  
164 of beetles and different backgrounds (twigs from Scotch pine; stones; skin of museum samples  
165 of bank vole (*Myodes glareolus*); and birch leaf (*Betula pubescens*) were converted to predicted  
166 photoreceptor responses of single and double cone types of a blue tit (Hart, Partridge and  
167 Cuthill 2000, Hart 2001, Troscianko and Stevens 2015) by using a mapping function of the  
168 Image Calibration and Analysis Toolbox. This mapping is highly accurate compared to  
169 reflectance-based calculations of predicted cone responses (Stevens and Cuthill 2006, Pike  
170 2011, Troscianko and Stevens 2015). Colour vision in birds stems from the four single cone  
171 types (Cuthill 2006), while the double cones are likely responsible for luminance-based tasks  
172 (Vorobyev et al. 1998, Osorio and Vorobyev 2005), such as detecting achromatic contrast  
173 differences. The vision model converts the ROIs to cone-catch data, i.e. to the relative photon  
174 catches of a blue tit's four single cones: longwave (LW), mediumwave (MW), shortwave (SW)  
175 and ultraviolet (UV) cones, as well as to luminance values based on the double cone sensitivity.  
176 To analyse the phenotypic and genetic variation in colour of the beetles, we calculated

177 saturation values (colour richness) similar to (Arenas et al. 2015) and brightness (double cone  
178 sensitivity) for the ROIs of the first and second orange stripes and black pattern.

179 To analyse the conspicuousness of burying beetles to avian predators, colour and  
180 luminance discrimination models (Vorobyev and Osorio 1998) were conducted on cone-catch  
181 data of backgrounds and colour patterns of beetles with ImageJ Toolbox (MICA) (Troscianko  
182 and Stevens 2015). We first tested how well blue tits can discriminate between the orange and  
183 black pattern elements of beetles against various natural backgrounds. Family mean values of  
184 cone catch data for the first and second orange stripe and black pattern of colour and luminance  
185 were compared against different backgrounds. To test the intrapattern contrast of orange and  
186 black pattern elements, we compared mean values of cone catch data of the first and second  
187 orange stripes and the black pattern within an individual. Finally, to test whether birds can  
188 detect the variation in conspicuousness of the colouration among *N. vespilloides* families, we  
189 compared the family mean values of cone catch data of different pattern elements among  
190 families. The discrimination model uses units called just noticeable differences (hereafter,  
191 JNDs) where values <1-3 indicate that the two colours are likely indistinguishable under  
192 optimal light conditions and values >3 indicate that two objects are likely discriminable and by  
193 increasing degrees: the greater the value the more distinguishable the colours should be even  
194 under less optimal light conditions (Siddiqi et al. 2004). Four single cones were used for the  
195 colour discrimination model, whereas the luminance discrimination model was based on the  
196 double cones (Siddiqi et al. 2004). In the colour discrimination model, a Weber fraction of 0.05  
197 was used for the most abundant cone type, and the relative proportions of cone types in the  
198 blue tit retina (longwave = 0.96, mediumwave = 1, shortwave = 0.85, and ultraviolet sensitive  
199 = 0.46). A Weber fraction 0.05 was also used for modelling luminance discrimination using  
200 the double cones (Siddiqi et al. 2004, Sandre et al. 2010).

201

## 202 **Aim 2: Measuring noxiousness of the anal exudates using bioassays with ants**

203 Ants are important predators of insects (Molleman et al. 2010, Pavis et al. 1992, Way and Khoo  
204 1992) and one of the most important competitors with burying beetles for carcasses (Scott  
205 1998). Ants can also reliably recognize the presence of repellent compounds, and thus are ideal  
206 for conducting bioassays of potentially noxious substances (Deroe and Pasteels 1977, Hare and  
207 Eisner 1993, Dyer and Floyd 1993). Often deterrence against ants correlates with the deterrence  
208 against avian predators (Lindstedt et al. 2006 and 2011, Lindstedt et al. 2008, Reudler et al.  
209 2015).

210 We collected anal exudates from approximately 100 sexually matured beetles  
211 from the lab stock reared in standardized conditions. Anal exudates were collected by poking  
212 the abdomen of each beetle gently 1-2 times from the ventral side with a capillary tube, which  
213 caused the beetles to spray the fluid. Fluid was collected into the capillary tubes and pooled  
214 into 3 separate Eppendorf tubes and placed in a freezer (-20 C). Before presentation to the ants,  
215 samples were thawed and then diluted with a 20 % sugar solution (20% sugar, 80% water) to  
216 motivate the ants to feed on the solution. We conducted two separate bioassays with two  
217 concentrations to test how the variation in the concentration affected ants' willingness to feed  
218 on it. In the first assay, we tested the deterrence of anal exudate by offering 10% exudate  
219 solution (10% anal exudate / 90% sugar water) and palatable control solution (10% of plain

220 water / 90% sugar water) to ants simultaneously. In the second assay, we used 1 % exudate  
221 solution (1% anal exudate / 99% sugar water) and 1% control solution (1% plain water / 99%  
222 sugar water).

223 Bioassays were conducted similar to Reudler et al. 2015. We performed tests with  
224 the 10% exudate and control solutions on 10 different ant (*Formica sp.*) nests in the field in  
225 central Finland (62 °N, 26° E) in sunny and warm weather (15-20°C). To standardize the  
226 potential variation in activity and ant traffic among ant nests, we presented ants simultaneously  
227 with droplets of exudate and control solutions. In the vicinity of each nest we chose a spot on  
228 the trail where ant traffic was about 10 to 20 individuals/minute. We put 10 µl of both the  
229 control and exudate solutions close to each other (<2 cm) on a transparent, sterilized plastic  
230 circle (4 cm in diameter) and offered it to the ants. We repeated the assay three times per nest,  
231 each on a different ant trail, and order of control and exudate droplets was changed between  
232 repetitions. During the experiment we calculated the number of ants drinking from the different  
233 solutions in 1 minute intervals during the 10 minutes and counted the mean number of ants that  
234 drank each type of fluid to measure its aversiveness (Reudler et al. 2015). Recording was  
235 started after the first ant worker arrived at either of the droplets. We repeated exactly the same  
236 procedure one week later with the 1 % control and exudate solutions, using five of the same  
237 nests as those used in 10% solution assays. All of the experiments were run within a 2 week  
238 period in August 2010.

239

### 240 **Aim 3: Variation in chemical defence, orange elytra pattern and colour**

241 We set up 25 pairs for breeding with a carcass (mean ± S.E. carcass mass given above, in  
242 description of breeding conditions). Both parents remained with the offspring until larvae  
243 dispersed, at which point they were discarded and the larvae were transferred to separate  
244 individual boxes to pupate. After eclosion, when individuals had developed the typical black  
245 and orange coloration, they were sexed and the quantity of the defence fluid was measured by  
246 poking the abdomen of each beetle gently 1-2 times from the ventral side with a capillary tube,  
247 which caused the beetles to spray the fluid. Fluid was collected into the capillary tubes and the  
248 quantity produced was measured. Beetles were then weighed and killed by storing them in a  
249 freezer for -20°C. Frozen individuals were photographed after the experiment using a calibrated  
250 Fuji IS digital camera, which records both ultraviolet and human visible signals. From the  
251 photographs, the size of the elytra and orange patterns were measured with the ImageJ -  
252 program and hue and brightness of the pattern components analysed with the Image Calibration  
253 and Analysis Toolbox (Troscianko and Stevens 2015) with the method described above. In  
254 total, we aimed to measure the anal exudate volume from 5 females and 5 males from each of  
255 25 families. One individual was left out from the analyses as we failed to measure the defensive  
256 response and for some families the number of offspring was less than 10 individuals. We  
257 sampled 3-10 individuals per family (mean 8.96 ± 0.35 S.E.) yielding 224 samples in total.  
258 Signal size and colour measurements were taken from 98 individuals across 14 families.

259

### 260 **Statistics**

261 To take into account possible variation in ant behaviour and activity among the nests and trails,  
262 we used pairwise t-tests to compare the mean number of ants feeding on exudate solution and

263 control solution for the bioassays with 10% concentration and 1% exudate and control  
264 solutions. Data from the ant experiments were analysed using IBM SPSS Statistics 20 (IBM  
265 Corporation, NY, USA).

266 We used general linear mixed models to analyse the relationship between sex and  
267 elytra size on the volume of anal exudate produced, and on each of the other measures of the  
268 aposematic signal: the size, brightness and saturation of the two orange stripes and the  
269 brightness and saturation of the black portions of the elytra. The fit of each model was checked  
270 by examination of the residuals. The two measures of the black colour were log transformed  
271 as inspection of residuals suggested deviations from a normal distribution. We applied model  
272 selection by comparing nested models with ANOVA. In all models, family was included as a  
273 random effect to account for variation due to genetic or maternal effects. Variance components  
274 from the random model associated with family ( $V_G$ ) and residual variance ( $V_R$ ) were used to  
275 calculate broad sense heritability ( $H^2$ ) for each of the traits, where  $H^2 = V_G/(V_G + V_R)$ . For  
276 mixed models, we used the “lme4” package in R (Bates et al. 2013);  $t$ -statistics, degrees of  
277 freedom and  $p$ -values were calculated using Satterthwaite’s approximation, with the  
278 “lmerTest” package in R (Kuznetsova et al. 2013). The significance of the random effects was  
279 tested against a Chi-squared distribution. The coefficients of genetic ( $CV_G$ ) and residual ( $CV_R$ )  
280 variation were calculated using untransformed data, as values for transformed data are  
281 meaningless (Houle 1992).

282

## 283 RESULTS

284

### 285 **Aim 1: Quantifying the salience of the orange-black coloration to avian predators**

286 The avian vision model for blue tits shows that avian predators should be able to discriminate  
287 orange and black patterns of burying beetles against various backgrounds (green leaves, grey  
288 stones, twigs, vole fur) both in terms of colour and luminance (Table 1). Within-pattern contrast  
289 of black and orange patterns was high and clearly visible for birds both in terms of colour and  
290 luminance (Table 1). Also, interestingly, the differences in the mean contrast values of the hue  
291 of pattern elements among families should be clearly visible for avian predators (Table 1).  
292 However, variation in the luminance contrasts of orange pattern elements among families are  
293 probably more difficult for birds to discriminate (Table 1).

294

### 295 **Aim 2: Measuring noxiousness of the anal exudates using bioassays with ants**

296 We found that significantly more ants took the sugar water than sugar water mixed with anal  
297 exudate of beetles (10% exudate: 90% sugar water) ( $t = -6.678$ ,  $n = 30$ ,  $p < 0.001$ ). However,  
298 when the concentration was decreased (1% exudate: 99% sugarwater), we could not detect any  
299 difference between the treatments ( $t = -0.400$ ,  $n = 15$ ,  $p = 0.695$ ) (Fig. 2). Thus, a higher  
300 concentration of anal exudates resulted in better defence against ants.

301

302 **Table 1. Discrimination values (JND) for colour (hue) and brightness of different elytra**  
 303 **pattern elements of *N. vespilloides* (values are average of 11 families) against various**  
 304 **natural backgrounds according to model by Vorobyev et al. 1998. Variation in**  
 305 **conspicuousness among families is based on two-way comparisons of average**  
 306 **discrimination values of 11 families. Brackets show the range between minimum and**  
 307 **maximum values). Values > 3 are easy to tell apart in most conditions.**  
 308

Comparisons of elytra pattern elements	Colour (hue)	Brightness
	Mean discrimination value (min-max)	Mean discrimination value (min-max)
<i>Intrapattern contrasts</i>		
Black versus orange in the first stripe	90.09 (56.69-110.83)	34.20 (26.93-39.05)
Black versus orange in the second stripe	94.88 (77.39-111.17)	31.03 (25.59-35.86)
Orange in the first stripe versus orange in the second stripe	13.85 (53.31-3.00)	3.17 (0.20-7.06)
<i>Elytra pattern contrasts against natural backgrounds</i>		
Black against the pine twig	16.36 (3.66-23.26)	26.27 (19.53-30.61)
Black against the birch leaf	44.95 (34.07-53.40)	36.27 (26.09-43.20)
Black against the stone	12.17 (6.81-20.94)	39.63 (31.66-46.22)
Black against the bank vole fur	14.71 (3.12-24.07)	28.38 (13.94-38.77)
Orange in the 1st stripe against the pine twig	76.77 (49.94-94.63)	7.93 (3.47-12.70)
Orange in the 1st stripe against the birch leaf	58.55 (40.93-71.28)	3.22 (0.04-9.13)
Orange in the 1st stripe against the stone	93.93 (63.11-114.46)	5.47 (0.37-12.14)
Orange in the 1st stripe against the bank vole fur	79.61 (49.49-100.83)	6.18 (0.05-18.30)
Orange in the 2nd stripe against the pine twig	81.41 (72.12-101.33)	4.82 (0.44-8.27)
Orange in the 2nd stripe against the birch leaf	58.31 (46.57-82.53)	5.34 (0 - 13.04)
Orange in the 2nd stripe against the stone	98.97 (87.29-121.26)	8.60 (3.85-16.05)
Orange in the 2nd stripe against the bank vole fur	84.39 (72.03-107.53)	4.42 (0.08-13.87)
<i>Variation in conspicuousness among families</i>		
Black	8.60 (1.84 - 24.89)	3.78 (0.07-11.07)
Orange in the first stripe	53.72 (2.86-18.82)	2.33 (0.01-9.23)
Orange in the second stripe	11.67 (1.79-34.42)	2.66 (0.02-8.71)

309

310 **Aim 3a): Variation in chemical defence**

311 Body size was not associated with the amount of anal exudate beetles produced (Table 2).  
 312 However, females produced significantly higher quantities of fluid than males (REML:  
 313 Estimate=  $2.89 \pm 0.855$ ; Table 2; Fig. 3). The amount of anal exudate produced upon  
 314 disturbance showed a moderate broad-sense heritability of 0.38 (Table 3).

315

316 **Table 2. ANOVA results for the fixed effects of sex, elytra size and their interaction on**  
 317 **the amount of eclosion fluid produced and elements of the aposematic signal.**


318

Trait	Sex	Elytra size	Sex:Elytra size
<b>Eclosion fluid</b>	<b>F<sub>1,198</sub> = 11.4</b> <b>P &lt; 0.001</b>	F <sub>1,99</sub> = 2.06 P = 0.15	F <sub>1,94</sub> = 2.88 P = 0.09
<b>Orange total (mm)</b>	F <sub>1,98</sub> = 0.17 P = 0.68	<b>F<sub>1,109</sub> = 92.68</b> <b>P &lt; 0.001</b>	F <sub>1,96</sub> = 0.61 P = 0.44
<b>First stripe (mm)</b>	F <sub>1,98</sub> = 0.01 P = 0.93	<b>F<sub>1,110</sub> = 78.03</b> <b>P &lt; 0.001</b>	F <sub>1,96</sub> = 0.01 P = 0.92
<b>Second stripe (mm)</b>	F <sub>1,99</sub> = 1.12 P = 0.29	<b>F<sub>1,110</sub> = 65.95</b> <b>P &lt; 0.001</b>	F <sub>1,97</sub> = 2.56 P = 0.11
<b>Brightness stripe 1</b>	F <sub>1,87</sub> = 1.05 P = 0.31	<b>F<sub>1,35</sub> = 11.13</b> <b>P = 0.002</b>	F <sub>1,87</sub> = 0.00 P = 0.96
<b>Saturation stripe 1</b>	F <sub>1,85</sub> = 0.49 P = 0.48	<b>F<sub>1,41</sub> = 4.70</b> <b>P = 0.036</b>	F <sub>1,86</sub> = 1.26 P = 0.26
<b>Brightness stripe 2</b>	F <sub>1,88</sub> = 1.36 P = 0.25	F <sub>1,24</sub> = 0.95 P = 0.34	F <sub>1,86</sub> = 0.00 P = 0.98
<b>Saturation stripe 2</b>	F <sub>1,87</sub> = 0.04 P = 0.84	<b>F<sub>1,39</sub> = 24.83</b> <b>P &lt; 0.001</b>	F <sub>1,87</sub> = 0.23 P = 0.63
<b>Brightness black</b>	<b>F<sub>1,87</sub> = 4.27</b> <b>P = 0.04</b>	F <sub>1,32</sub> = 0.59 P = 0.45	F <sub>1,85</sub> = 1.01 P = 0.32
<b>Saturation black</b>	F <sub>1,87</sub> = 1.09 P = 0.30	F <sub>1,35</sub> = 0.10 P = 0.76	F <sub>1,85</sub> = 1.68 P = 0.20


319

320

321



**Figure 2.** Mean number of ants (+- 1 SE) drinking the control solution (20% sugarwater) indicated by open circles and 10% (10% anal exudate: 90 % sugarwater) and 1 % experimental solution (1 % anal exudate: 99% sugarwater) indicated by closed circles.



**Figure 3.** Mean volume of anal exudate produced under disturbance by *N. vespilloides* females and males.

323 **Aim 3b): Variation in orange elytra pattern and colour**

324 The total size of the orange elytra pattern did not differ between males and females but did  
 325 increase with the size of the elytra (REML: Estimate =  $0.42 \pm 0.043$ ; Table 2). The same pattern  
 326 was found if the two orange stripes were considered independently (Table 2).

327 The brightness of the first orange stripe was significantly higher than the second  
 328 (Paired t-test,  $t_{184}=6.9$ ,  $P<0.001$ ), although the saturation of the stripes did not differ (Paired t-  
 329 test,  $t_{187}=0.69$ ,  $P=0.49$ ). However, whilst the saturation of both stripes increased with elytra  
 330 size (REML: Stripe 1 estimate =  $0.0006 \pm 0.00027$ , stripe 2 estimate =  $0.0010 \pm 0.00019$ ; Table  
 331 2), the brightness of the first stripe decreased as beetles got bigger (REML: Estimate =  $68.13$   
 332  $\pm 20.42$ ; Table 2) and elytra size had no effect on the brightness of the second stripe (Table 2).  
 333 The brightness of the black sections of the elytra were lower in males (REML: Estimate = -  
 334  $195.12 \pm 89.21$ ; Table 2) but were not affected by the size of the beetles (Table 2).

335 The size of the orange pattern, both in total and in the first and second stripe  
 336 separately, showed high broad sense heritabilities (range = 0.57-0.65, Table 3). None of the  
 337 measures of saturation and brightness was significantly heritable, though the saturation of the  
 338 first stripe and the brightness of the black were marginally non-significant (range - 0.03-0.12,  
 339 Table 3).

340  
 341

342 **Table 3. Genetic and residual variance in the amount of eclosion fluid produced and**  
 343 **elements of the aposematic signal as estimated by REML using the lmer package in R.  $V_G$**   
 344 **represents additive, dominance and epistatic variation.  $H^2$  is the broad sense heritability**  
 345 **estimate  $V_G/ V_R$ ,  $CV_G$  and  $CV_R$  are the coefficients of genetic and residual variance**  
 346 **respectively. Significance was tested with chi squared.  $P>0.10$  <sup>n.s.</sup>,  $P<0.10$  <sup>+</sup>,  $P<0.001$  <sup>\*\*\*</sup>**  
 347

Trait	No. families	$V_G$ (SD)	$V_R$ (SD)	$H^2$	chi	$CV_G$	$CV_R$
<b>EF</b>	25	24.53 (4.95)	40.02 (6.33)	0.38	54.6***	20.1	15.8
<b>Orange total (mm)</b>	14	12.09 (3.48)	6.51 (2.55)	0.65	81.8***	28.8	39.2
<b>First stripe (mm)</b>	14	5.57 (2.36)	3.15 (1.78)	0.64	75.3***	42.4	56.3
<b>Second stripe (mm)</b>	14	1.68 (1.30)	1.26 (1.12)	0.57	65.3***	77.2	89.1
<b>Brightness stripe 1</b>	11	158064 (398)	2244302 (1498)	0.06	1.25	0.25	0.07
<b>Saturation stripe 1</b>	11	4.84e-05 (0.007)	0.00019 (0.014)	0.12	2.93 <sup>+</sup>	14374	5323
<b>Brightness stripe 2</b>	11	99341 (315)	1660766 (1289)	0.06	2.07 <sup>n.s.</sup>	0.32	0.08
<b>Saturation stripe 2</b>	11	1.81e-05 (0.004)	1.92e-04 (0.014)	0.09	0.39 <sup>n.s.</sup>	23525	7206
<b>Brightness black</b>	11	18432 (136)	191849 (438)	0.09	2.43 <sup>+</sup>	0.74	0.23
<b>Saturation black</b>	11	1.73e-04 (0.013)	0.002 (0.045)	0.08	1.64 <sup>n.s.</sup>	7604	2224

348

349

350 DISCUSSION

351 Our first aim was to determine the salience of the burying beetle's orange and black coloration  
352 to avian predators, against a range of natural backgrounds. We found that these elytral markings  
353 of the burying beetle are highly conspicuous for avian predators. Objectively, the burying  
354 beetle's orange-black elytral patterning does not differ much from the orange-black patterning  
355 of other insect species which are widely recognised to be aposematic, such as *Arctia plantaginis*  
356 larvae (Lindstedt et al. 2008) and adult females (Lindstedt et al. 2011), ladybirds (Linas et al.  
357 2015) or *Heliconius* butterflies (Langham 2004). Furthermore, some *Nicrophorus* species have  
358 also been suggested to be Müllerian mimics of wasps and bumble-bees (Morton Jones 1932,  
359 Milne and Milne 1944, Lane and Rothschild 1965, Anderson and Beck 1985), each of which  
360 is known to deter avian predators. These observations, in conjunction with earlier reports that  
361 birds find burying beetles highly aversive (summarised in the Introduction), strongly suggest  
362 that many species of burying beetle use their orange and black elytral patterns as part of a  
363 warning display, and that these markings are under selection from avian predators. Collectively  
364 the evidence for aposematism (visual analyses about the conspicuousness of coloration  
365 combined with the bioassay for toxicity and presence of responsive defence) in the burying  
366 beetle is as strong as the evidence for a many other classical examples of an aposematism and  
367 Müllerian mimics such as defended Hymenopterans (e.g. Penney et al. 2012, Wilson et al.  
368 2012), poison frogs (e.g. Maan & Cummings 2012), ladybirds (e.g. Linas et al. 2015) or marine  
369 opisthobranchs (e.g. Cortesi and Cheney 2010).

370 We fulfilled our second aim by demonstrating that the chemical defences in the  
371 burying beetle's anal exudates are aversive, using a standard bioassay with wood ants (Reudler  
372 et al. 2015). In our experiments, a greater concentration of anal exudate resulted in better  
373 defence against ants, suggesting that the production of more potent exudates should enhance  
374 the efficacy of the beetle's chemical defence. The most conservative interpretation of these  
375 results is that burying beetles can defend themselves, and their carrion breeding resource,  
376 specifically against ants (e.g. Scott et al. 1987). However, deterrence against ants often correlates  
377 with the deterrence against avian predators in chemically defended species (Deroe and Pasteels  
378 1977, Hare and Eisner 1993, Dyer and Floyd 1993, Lindstedt et al. 2006 and 2011, Lindstedt  
379 et al. 2008, Reudler et al. 2015). Therefore a wider possible interpretation is that burying beetles  
380 possess a general chemical defence against their potential predators. If this is true, it means that  
381 the burying beetle's anal exudates serve a dual function by contributing to two public resources:  
382 the defence of the carrion breeding resource against microbes (Duarte et al. 2016, Duarte et al.  
383 2017) as well as the collective education of potential predators via warning displays (Speed et  
384 al. 2012). The constituents within the exudates are therefore likely to be subjected to differing  
385 selection pressures from each of these two functions.

386 These contrasting selection pressures might explain why we found high levels of  
387 individual variation in the volume of anal exudate produced. We also found a sex difference in  
388 the volume of anal exudates produced by burying beetles, though this is harder to explain. One  
389 possibility is connected with a sex difference in the function of the anal exudates, namely the  
390 antimicrobial defence of the carcass during reproduction. When preparing carrion for  
391 reproduction, burying beetles strip the body of fur or feathers, mould the the flesh into a ball

392 and smear it with antimicrobial anal exudates (Scott 1998, Rozen et al 2008, Cotter and Kilner  
393 2010). Females contribute exudates with greater lytic activity than males to this defence (Cotter  
394 and Kilner 2010), and likewise secrete a greater volume of fluid than males when handled (this  
395 study). In future work it would be interesting to test whether, and in what direction, the  
396 antimicrobial activity is correlated with the repellence of the anal exudate.

397           A second possibility is that females secrete a greater volume of exudates when  
398 threatened because they are more vulnerable to attacks by potential predators. The carcass is  
399 an attractive resource to scavengers and yet attended by parents during reproduction. Females  
400 spend much longer than males associated with the carcass, since males leave the brood before  
401 larval development is complete (Scott 1998, Boncoraglio and Kilner 2012, de Gasperin et al.  
402 2015). Females might therefore be more likely than males to encounter a potential predator,  
403 and this could explain why they produce more exudate when threatened. However, it is  
404 important to remember that we only measured the quantity of the fluid here. Thus, it is possible  
405 that males can compensate the lower amount of exudate by making it more noxious. In addition,  
406 we measured the quantity of fluid only once per individual and therefore we do not know if  
407 males are not able to produce more fluid or if they were just not willing to do so.

408           Whatever the reason for this sex difference, it suggests that higher volumes  
409 produced by females are potentially contributing more to the education of naïve predators than  
410 are males. Understanding the evolutionary significance of this difference will again come down  
411 to understanding the cost of the chemical defence. If females can produce more anal exudates  
412 than males for the same cost, then they are simply contributing to a public good in relation to  
413 their ability to pay, as predicted by theory (Frank 2010, Duarte et al. 2016). But if females are  
414 paying a higher cost for educating predators with their greater noxiousness then they are  
415 vulnerable to exploitation by males, who can potentially gain the same protection from  
416 predation but for a lower price. If this is indeed the case then the puzzle for future work is to  
417 explain why such exploitation persists.

418           We have assumed throughout that an individual's chemical defences are fixed in  
419 their potency and producing higher volumes is favoured for both parental care and chemical  
420 defence. Yet burying beetles can flexibly adjust the antimicrobial function of their anal  
421 exudates, up-regulating it only when reproducing and varying its potency in relation to their  
422 partner's contributions, and the scale of microbial threat to the carcass (Cotter and Kilner 2010,  
423 Cotter et al. 2010, Haberer et al. 2014). Although a plastic response like this cannot account  
424 for our measurements, because they were taken when beetles were not breeding, it would be  
425 interesting to test whether burying beetles are similarly capable of adjusting the concentration  
426 of fluid they exude when threatened, increasing the potency when the threat of attack is greater  
427 during reproduction on the carcass.

428           We found high levels of individual variation in elytral markings as well as in the  
429 volume of the exudates produced. Each might be attributable to an environmental or genetic  
430 constraint upon the production of each trait (Lindstedt et al. 2009, Lindstedt et al. 2016). To  
431 understand how variation in colour patterning and chemical defences arise we need to know  
432 more about the costs associated with these traits and how they are affected by early  
433 developmental environment of the beetles. In addition, it is important to know the chemical  
434 structure of pigments (e.g. Lindstedt et al. 2010b) and defence chemicals. Burying beetles are

435 carnivorous insects and their diet is scarce in antioxidants in comparison to herbivores (Olson  
436 and Owens 1998, Bortolotti et al. 2000). If the orange pigmentation is protein based, it might  
437 be relatively cheaper for a carnivore to produce than if the orange colour was dependent on  
438 carotenoids or flavonoids, which are much rarer in a carnivorous diet. In the latter case, burying  
439 beetles would need to synthesize pigments and defensive chemicals *de novo* and this may  
440 require energy and resources that are scarce in their diet. It might even involve recruiting  
441 microbial symbionts for this purpose (Moran and Jarvik 2010, Tsuchida et al. 2010). For the  
442 repellent compounds in anal exudate it is already known that they are mainly based on amino-  
443 acids (Degenkolb et al. 2011) and therefore likely to be synthesized *de novo* and constrained  
444 by the quality and availability of proteins in the diet.

445         Since variation in both burying beetle elytral markings and their anal exudates  
446 are potentially connected to diet, it would be interesting in future work to determine the extent  
447 to which individual variation can be explained by variation in the level post-hatching care  
448 received during early life. Our calculations suggest that the broad-sense heritability of each  
449 trait is relatively high, but our measures cannot partition out the separate effects of the  
450 developmental environment from inherited genetic variation. Previous work on other burying  
451 beetle traits has found that once the developmental environment is accounted for, trait  
452 heritability is relatively low (e.g. Lock et al 2004). Nevertheless, this does not necessarily mean  
453 that traits cannot respond to selection by predators or other agents (Kilner et al 2015, Jarrett et  
454 al 2017) and exactly how this happens will need to be determined more explicitly in future  
455 work.

456         In conclusion, our experiments, together with evidence in the literature, strongly  
457 suggest that the orange-black colouring of the burying beetle's elytra serves an aposematic  
458 function and anal exudate of beetles can serve multiple functions in antipredator defence and  
459 parental care. The challenge for future work is to deduce the costs associated with producing  
460 both the colourful display and the chemical defence so as to better explain the intra-specific  
461 variation we have found. We also need more information about the selection pressures that  
462 visual predators, namely birds, impose on the colour and size of the pattern as well as toxicity  
463 of the anal exudate. We note that not all *Nicrophorus* species are orange and black, and that  
464 some entirely black species still produce a malodorous fluid when handled (e.g. *N. humator*,  
465 R. M. Kilner pers. obs.). Therefore, the genus *Nicrophorus* in general provides the opportunity  
466 to test: 1) why some chemically defended species have evolved conspicuous marking while  
467 others have not; 2) how aposematism is linked to different life-history strategies and social  
468 behaviour; and 3) how individuals can balance their contributions to two different sorts of  
469 public goods: chemical defence and antimicrobial defence of a carrion breeding resource.

470  
471

472 FUNDING

473 This study was funded by the Academy of Finland via the projects no 136387, 257581 (CL).  
474 GB was funded by Marie Curie Intra-European Fellowship 252120 LIFHISBURBEE. SC was  
475 supported by a Natural Environment Research Council grant (NE/H014225/2). RMK was  
476 supported in part by the European Research Council (ERC) Consolidators grant 310785  
477 BALDWINIAN\_BEETLES and by a Wolfson Merit Award from the Royal Society.

478

479 ACKNOWLEDGEMENTS

480 We are grateful for all the research assistants helping with the maintenance of lab population  
481 and Martin Stevens and Jolyon Troscianco for helping with the colour analyses.

482

483 DATA ACCESSIBILITY STATEMENT

484 Analyses reported in this article can be reproduced using the data sets provided by Lindstedt  
485 et al. 2017. Data for the individual variation in the chemical defence, elytra colour and size of  
486 the markings in *Nicrophorus vespilloides*, will be released on 1<sup>st</sup> of August 2018. However,  
487 all reasonable requests for materials will be respected before that time on request.

488

489 REFERENCES

- 490 Alatalo R, Mappes J. 1996. Tracking the evolution of warning signals. *Nature*. 382:708-710.
- 491 Anderson RS, Peck SB. 1985. Geographic patterns of colour variation oin North American  
492 *Nicrophorus* burying beetles (Coleoptera; Silphidae). *J. Nat. His.* 20:283-297.
- 493 Arce AN, Johnston PR, Smiseth PT, Rozen DE. 2012. Mechanisms and fitness effects of  
494 antibacterial defences in a carrion beetle. *J. Evol. Biol.* 25:930-937.
- 495 Arenas LM, Walter D, Stevens M. 2015. Signal honesty and predation risk among a closely  
496 related group of aposematic species. *Sci. Rep.* 5:1-12.
- 497 Beatty CD, Beirinckx K, Sherratt TN. 2004. The evolution of Müllerian mimicry in  
498 multispecies communities. *Nature*. 431:63-67.
- 499 Bezzerides A, MacGraw K, Parker R, Hussein J. 2007. Elytra colour as a signal of chemical  
500 defence in the Asian ladybird beetle *Harmonia axyridis*. *Behav. Ecol. Sociobiol.* 61:  
501 1401-1408.
- 502 Boncoraglio G, Kilner RM. 2012. Female burying beetles benefit from male desertion: sexual  
503 conflict and counter-adaptation over parental investment. *PLoS One*. 7:e31713.
- 504 Brakefield PM. 1985. Polymorphic Müllerian mimicry and interactions with thermal melanism  
505 in ladybirds and a soldier beetle: a hypothesis. *Biol. J. Linnean Soc.* 26:243-267.
- 506 Brower L, Edmunds M, Moffitt C. 1975. Cardenolide content and palatability of a population  
507 of *Danaus chrysippus* butterflies from West Africa. *J. Entomol. Series A, General*  
508 *Entomology* 49:183–196.
- 509 Conner WE, Eisner T, Vander Meer RK, Guerrero A, Meinwald J. 1981. Precopulatory  
510 sexual interaction in an arctiid moth (*Utetheisa ornatrix*): Role of a pheromone derived  
511 from dietary alkaloids. *Behav. Ecol. Sociobiol.* 9:227.
- 512 Cotter SC, Kilner RM. 2010. Personal immunity versus social immunity. *Behav. Ecol.* 21:  
513 663-668.

514 Cotter SC, Topham E, Price AJP, Kilner RM. 2010. Fitness costs associated with mounting a  
515 social immune response. *Ecol. Lett.* 13:1114-1123.

516 Cotter SC, Littlefair JE, Grantham PJ, Kilner RM. 2013. A direct physiological trade-off  
517 between personal and social immunity. *J. Anim. Ecol.* 82:846-853.

518 Cortesi F, Cheney KL. 2010. Conspicuousness is correlated with toxicity in marine opisthobranchs. *J.*  
519 *Evol. Biol.* 23:1509–1518.

520 Crothers LR, Cummings ME. 2013. Warning signal brightness variation: Sexual selection may  
521 work under the radar of natural selection in populations of a polytypic poison frog. *Am.*  
522 *Nat.* 181: E116-E124.

523 Cuthill IC. 2006. vol. I: Mechanisms and Measurements. *Bird Colouration* (eds. G. E. Hill and  
524 K. J. McGraw). Cambridge, Massachusetts: Harvard University Press. p. 3-40

525 Daly D, Higginson AD, Chen D, Ruxton GD, Speed MP. 2012. Density-dependent investment  
526 in costly anti-predator defences: an explanation for the weak survival benefit of group  
527 living. *Ecol. Lett.* 15:576-583.

528 de Gasperin O, Kilner RM. 2015. Interspecific interactions change the outcome of sexual  
529 conflict over prehatching parental investment in the burying beetle *Nicrophorus*  
530 *vespilloides*. *Ecol. Evol.* 5:5552-5560.

531 Degenkolb T, Düring RA, Vilcinskas A. 2011. Secondary metabolites released by the burying  
532 beetle *Nicrophorus vespilloides*: Chemical analyses and possible ecological functions.  
533 *J. Chem. Ecol.* 37:724-735.

534 Deroe C, Pasteels JM. 1977. Defensive mechanisms against predation in the Colorado beetle  
535 (*Leptinotarsa decemlineata* Say). *Arch. Biol. Sci.* 88:289–304.

536 Duarte A, Cotter SC, Reavey RE, Ward RJS, de Gasperin O, Kilner RM. 2016. Social  
537 immunity of the family: parental contributions to a public good modulated by brood  
538 size. *Evol. Ecol.* 30:123-135.

539 Duarte S, Welch M, Swannack C, Wagner J, and Kilner RM. 2017. Strategies for managing  
540 rival bacterial communities: lessons from burying beetles. *J. Anim. Ecol.* in press

541 Dyer LA, Floyd T. 1993. Determinants of predation on phytophagous insects: the importance  
542 of diet breadth. *Oecologia* 96:575-582.

543 Dyer LA, Dodson CD, Gentry G. 2003. A bioassay for insect deterrent compounds found in  
544 plant and animal tissues. *Phytochem. Anal.* 14:381–388.

545 Eggert AK, Reinking M, Müller JK. 1998. Parental care improves offspring survival and  
546 growth in burying beetles. *Anim. Beh.* 55:97-107.

547 Exnerová A, Svadová K, Štys P, Barcalová S, Landová E, Prokopová M, Fuchs R, Socha R.  
548 2006. Importance of colour in the reaction of passerine predators to aposematic prey:  
549 experiments with mutants of *Pyrrhocoris apterus* (Heteroptera). *Biol. J. Linnean Soc.*  
550 88:143-153.

551 Forsman A, Merilaita S. 1999. Fearful symmetry: pattern size and asymmetry affects  
552 aposematic signal efficacy. *Evol. Ecol.* 13:131-140.

553 Frank SA. 2010. ‘A general model of the public goods dilemma’ *J. Evol. Biol.* 23:1245–1250.

554 Friman V, Lindstedt C, Hiltunen T, Laakso J, Mappes J. 2009. Predation on multiple trophic  
555 levels shapes the evolution of pathogen virulence. *PLoS ONE* 4:e6761.

556 Gamberale-Stille G, Tullberg B. 1999. Experienced chicks show biased avoidance of stronger  
557 signals: an experiment with natural colour variation in live aposematic prey. *Evol. Ecol.*  
558 13:579-589.

559 Gordon SP, Kokko H, Rojas B, Nokelainen O, Mappes J. 2015. Colour polymorphism torn  
560 apart by opposing positive frequency-dependent selection, yet maintained in space. *J.*  
561 *Anim. Ecol.* 84:1555-1564.

562 Guilford T. 1990. The evolution of aposematism. Insect defence: adaptive mechanisms and  
563 strategies of prey and predators (eds. D. L. Evans and Schmidt, J. O.). New York, State  
564 University of New York Press: p. 23-61.

565 Grill CP, Moore AJ. 1998. Effects of a larval antipredator response and larval diet on adult  
566 phenotype in an aposematic ladybird beetle. *Oecologia* 114:274-282.

567 Haberer W, Steiger S, Müller JK. 2014. Dynamic changes in volatile emissions of breeding  
568 burying beetles. *Phys. Entomol.* 39:153-164.

569 Hall CL, Mason AC, Howard DR, Padhi A, Smith RJ. 2013. Description of acoustic characters  
570 and stridulatory pars stridens of *Nicrophorus* (Coleoptera: Silphidae): a comparison of  
571 eight North American species. *Ann. Entomol. Soc. N. Amer.* 106:661-669.

572 Hare JF, Eisner T. 1993. Pyrrolizidine alkaloid deters ant predators of *Utetheisa ornatix* eggs-  
573 effects of alkaloid concentration, oxidation-state, and prior exposure of ants to alkaloid-  
574 laden prey. *Oecologia* 96:9-18.

575 Hart NS, Partridge JC, Cuthill IC. 2000. Retinal asymmetry in birds. *Curr. Biology* 10:115-  
576 117.

577 Hart NS. 2001. The visual ecology of avian photoreceptors. *Prog. Retin. Eye Res.* 20:675-703.

578 Hegna RH, Nokelainen O, Hegna J, Mappes J. 2013. To quiver or to shiver: increased  
579 melanization benefits thermoregulation, but reduces warning signal efficacy in the wood  
580 tiger moth. *Proc. R. Soc. Lond. B.* 280:2012-2812.

581 Higginson AD, Delf J, Ruxton GD, Speed M. 2011. Growth and reproductive costs of larval  
582 defence in the aposematic lepidopteran *Pieris brassicae*. *J. Anim. Ecol.* 80:384-392.

583 Houle D. 1992. Comparing evolvability and variability of quantitative traits. *Genetics* 130:  
584 195-204.

585 Ihalainen E, Lindström L, Mappes J. 2007. Investigating Müllerian mimicry: predator learning  
586 and variation in prey defences. *J. Evol. Biol.* 20:780-791.

587 Jarrett BJM, Schrader M, Rebar D, Houslay TM, Kilner RM. 2017. Cooperative interactions  
588 within the family enhance the capacity for evolutionary change in body size. *Nat. Ecol.*  
589 *Evol.* 1:0178.

590 Joron M, Mallet JLB. 1998. Diversity in mimicry: paradox or paradigm? *Trends in Ecology*  
591 *and Evolution* 13:461-466.

592 Kapan DD. 2001. Three-butterfly system provides a field test of müllerian mimicry. *Nature*  
593 409:338-340.

594 Kilner RM, Boncoraglio G, Henshaw JM, Jarrett BJM, De Gasperin O, Attisano A, Kokko H.  
595 2015. Parental effects alter the adaptive value of an adult behavioural trait. *eLIFE*  
596 4:e07340.

597 Kollberg I, Bylund H, Jonsson T, Schmidt A, Gershenson J, Björkman C. 2015. Temperature  
598 affects insect outbreak risk through tritrophic interactions mediated by plant secondary  
599 compounds. *Ecosphere* 6:1-17.

600 Kuznetsova A, Brockhoff PB, Christensen RHB. 2013. lmerTest: tests for random and fixed  
601 effects for linear mixed effect models (lmer objects of lme4 package) [WWW document].  
602 URL <http://CRAN.R-project.org/package=lmerTest> [accessed on 12 February 2015].

603 Lane C, Rothschild M. 1965. A case of Müllerian mimicry of sound. *Proc. R. Entomol. Soc.*  
604 *Lond. A* 40:156-158.

605 Langham GM. 2004. Specialized avian predators repeatedly attack novel color morphs of  
606 *Heliconius* butterflies. *Evolution*. 58:2783-2787.

607 Laurentz M, Reudler JH, Mappes J, Friman V, Ikonen S, Lindstedt C. 2012. Diet quality can  
608 play a critical role in defence efficacy against parasitoids and pathogens in the Glanville  
609 Fritillary (*Melitaea cinxia*). *J. Chem. Ecol.* 38:116-125.

610 Leimar O, Enquist M, Sillén-Tullberg B. 1986. Evolutionary stability of aposematic coloration  
611 and prey unprofitability: a theoretical analysis. *Am. Nat.* 128:469-490.

612 Lindstedt C, Lindström L, Mappes J. 2008. Hairiness and warning colours as components of  
613 antipredator defence: additive or interactive benefits? *Anim. Beh.* 75:1703-1713.

614 Lindstedt C, Lindström L, Mappes J. 2009. Thermoregulation constrains effective warning  
615 signal expression. *Evolution*. 63:469–478.

616 Lindstedt C, Reudler Talsma JH, Ihalainen E, Lindström L, Mappes J. 2010a. Diet quality  
617 affects coloration indirectly: excretion costs in a generalist herbivore. *Evolution*. 64: 68–  
618 78.

619 Lindstedt C, Morehouse N, Pakkanen H, Casas J, Christides J, Kemppainen K, Lindström L,  
620 Mappes J. 2010b. Pigment composition of a variable warning signal of *Parasemia*  
621 *plantaginis* larvae (Arctiidae). *Func. Ecol.* 24:759–766.

622 Lindstedt C, Eager H, Ihalainen E, Kahilainen A, Stevens M, Mappes J. 2011. Direction and  
623 strength of selection by predators for the hindwing color of the aposematic wood tiger  
624 moth. *Beh. Ecol.* 22:580-587.

625 Lindstedt C, Schroderus E, Lindström L, Mappes T, Mappes J. 2016. Evolutionary constraints  
626 of warning signals: A genetic trade-off between the efficacy of larval and adult warning  
627 coloration can maintain variation in signal expression. *Evolution*. 70:2562-2572.

628 Lindstedt C, Boncoraglio G, Cotter S, Gilbert J, Kilner RM. 2016. Data from: Aposematism in  
629 the burying beetle? Dual function of anal fluid in parental care and chemical defence.  
630 *Behavioral Ecology*. <http://dx.doi.org/10.5061/dryad.770sn>

631 Lindström L, Alatalo RV, Mappes J, Riipi M, Vertainen L. 1999. Can aposematic signals  
632 evolve by gradual change? *Nature*. 397:249-251.

633 Lock JE, Smiseth PT, Moore AJ. 2004. Selection, inheritance, and the evolution of parent-  
634 offspring interactions. *Am. Nat.* 164(1):13-24.

635 Maan ME, Cummings ME. 2008. Female preferences for aposematic signal components in a  
636 polymorphic poison frog. *Evolution*. 62:2334–2345.

637 Maan ME, Cummings ME. 2009. Sexual dimorphism and directional sexual selection on  
638 aposematic signals in a poison frog. *PNAS* 106:19072-19077.

639 Maan ME, Cummings ME. 2012. Poison frog colors are honest signals of toxicity, particularly  
640 for bird predators. *Am. Nat.* 179:E1-E14.

641 Mallet JLB, Barton NH. 1989. Strong natural selection in a warning-color Müllerian mimicry.  
642 *Evol. Ecol.* 13:777-806.

643 Mappes J, Alatalo RV. 1997a. Effect of novelty and gregariousness in survival of aposematic  
644 prey. *Beh. Ecol.* 8:174-177.

645 Mappes J, Kokko H, Ojala K, Lindström L. 2014. Seasonal changes in predator community  
646 switch the direction of selection for anti-predatory defences. *Nat. Com.* 5:5016.

647 Milne LJ, Milne MJ. 1944. Notes on the behaviour of burying beetles. (*Nicrophorus* spp.). *J.*  
648 *New York Entomol. Soc.* 52:311-327.

649 Molleman F, Kaasik A, Whitaker MR, Carey JR. 2012. Partitioning variation in duration of ant  
650 feeding bouts can offer insights into the palatability of insects: experiments of African  
651 fruit-feeding butterflies. *J. Res. Lepid.* 45:65–75.

652 Moran NA, Jarvik T. 2010. Lateral transfer of genes from fungi underlies carotenoid  
653 production in aphids. *Science.* 328:624-627.

654 Morton Jones F. 1932. Insect coloration and the relative acceptability of insects to birds. *Trans.*  
655 *Entomol. Soc. Lond.* 80:345-371.

656 Nokelainen O, Lindstedt C, Mappes J. 2013. Environment-mediated morph-linked immune and  
657 life-history responses in the aposematic wood tiger moth. *J. Anim. Ecol.* 3:653–662.

658 Nokelainen O, Valkonen J, Lindstedt C, Mappes J. 2014. Changes in predator community  
659 structure shifts the efficacy of two warning signals in Arctiid moths. *J. Anim. Ecol.* 83:  
660 598–605.

661 Ojala K, Lindström L, Mappes J. 2007. Life history constraints and warning signal expression  
662 in arctiid moth. *Func. Ecol.* 21:1162-1167.

663 Olson VA, Owens IPF. 1999. Costly sexual signals: are carotenoids rare, risky or required?  
664 *TREE* 14:510-514.

665 Osorio D, Vorobyev M. 2005. Photoreceptor spectral sensitivities in terrestrial animals:  
666 adaptations for luminance and color vision. *Proc. R. Entomol. Soc. Lond. A.* 272:1745–  
667 1752.

668 Pavis C, Malusse PH, Ducrot F, Jaffe K, Descoins C. 1992. Defensive secretions of first-  
669 instar larvae of rootstalk borer weevil, *Diaprepes abbreviatus* L. (Coleoptera:  
670 Curculionidae), to the fire-ant *Solenopsis geminata* (F.) (Hymenoptera: Formicidae). *J.*  
671 *Chem. Ecol.* 18:2055-2068.

672 Penney HD, Hassall C, Skevington JH, Abbott KR, Sherratt TN. 2012. A comparative analysis  
673 of the evolution of imperfect mimicry. *Nature.* 483:461-464.

674 Poulton EB. 1890. The colours of animals: their meaning and use especially considered in the  
675 case of insects (Edn 2), xiii, 360 pp. Kegan Paul, Trench, Trubner and co. London.

676 Reudler JH, Lindstedt C, Pakkanen H, Lehtinen I, Mappes J. 2015. Costs and benefits of plant  
677 allelochemicals in herbivore diet in a multi enemy world. *Oecologia.* 179:  
678 (DOI: 10.1007/s00442-015-3425-0).

679 Rojas B, Gordon SP, Mappes J. 2015. Frequency-dependent flight activity in the colour  
680 polymorphic wood tiger moth. *Curr. Zool.* 61:765–772.

681 Rothschild M, Lane C. 1960. Warning and alarm signals by birds seizing aposematic insects.  
682 *Ibis.* 102:328-330

683 Rowland HM, Ihalainen E, Lindström L, Mappes J, Speed MP. 2007. Co-mimics have a  
684 mutualistic relationship despite unequal defence levels. *Nature.* 448:64-66.

685 Rozen DE, Engelmoer DJP, Smiseth PT. 2008. Antimicrobial strategies in burying beetles  
686 breeding on carrion. *PNAS* 105:17890-17895.

687 Ruxton GD, Sherratt TN, Speed MP. 2004. Avoiding attack. Evolutionary ecology of crypsis,  
688 warning signals and mimicry. p. 249. Oxford University Press, New York.

689 Sandre S, Tammaru T, Esperk T, Julkunen-Tiitto R, Mappes J. 2007. Carotenoid-based colour  
690 polyphenism in a moth species: search for fitness correlates. *Entomol. Exp. Appl.*  
691 124:269-277.

692 Sandre S-L, Stevens M, Mappes J. 2010. The effect of predator appetite, prey warning  
693 coloration and luminance on predator foraging decisions. *Behaviour* 147:1121–1143.

694 Scott MP, Traniello JFA, Fetherston IA. 1987. Competition for prey between ants and burying  
695 beetles (*Nicrophorus* spp): Differences between northern and southern temperate sites.  
696 *Psyche* 94:325-332.

697 Scott MP. 1998. The ecology and behaviour of burying beetles. *Ann. Rev. Entomol.* 43:595-  
698 618.

699 Siddiqi A, Cronin TW, Loew ER, Vorobyev M, Summers K. 2004. Interspecific and  
700 intraspecific views of color signals in the strawberry poison frog *Dendrobates pumilio*. *J.*  
701 *Exp. Biol.* 207:2471–2485.

702 Sillén-Tullberg B. 1985. Higher survival of an aposematic than of a cryptic form of a distasteful  
703 bug. *Oecologia.* 67:411–415.

704 Sikes DS, Madge RB, Newton AD. A catalog of the Nicrophorinae (Coleoptera: Silphidae) of  
705 the world. 2002. *Zootaxa* 65:1-304.

706 Skelhorn J, Rowe C. 2006. Predator avoidance learning of prey with secreted or stored defences  
707 and the evolution of insect defences. *Anim. Behav.* 72:827-834

708 Speed M, Ruxton GD, Mappes J, Sherratt TN. 2012. Why are defensive toxins so variable? An  
709 evolutionary perspective. *Biol. Rev.* 87:874-884.

710 Stevens M. 2007. Predator perception and the interrelation between different forms of  
711 protective coloration. *Proc. R. Entomol. Soc. Lond. A.* 274:1457–1464.

712 Stevens M, Cuthill I. 2006. Disruptive coloration, crypsis and edge detection in early visual  
713 processing. *Proc. R. Entomol. S. Lond. A.* 7:2141-2147.

714 Stevens M, Ruxton GD. 2011. Linking the evolution and form of warning coloration in nature.  
715 *Proc. R. Soc. Lond. B.* DOI: 10.1098/rspb.2011.1932.

716 Troscianko J, Stevens M. 2015. Image calibration and analysis toolbox - a free software suite  
717 for objectively measuring reflectance, colour and pattern. *Methods in Ecology and*  
718 *Evolution* DOI: 10.1111/2041-210X.12439

719 Tsuchida, Tsutomu, Koga, Ryuichi, Horikawa M, Tsunoda T, Maoka T, Matsumoto S, Simon  
720 J-C, Fukatsu T. 2010. Symbiotic bacterium modifies aphid body color. *Science.* 19:  
721 1102-1104.

722 Valkonen JK, Nokelainen O, Niskanen M, Kilpimaa J, Björklund M, Mappes J. 2012. Variation  
723 in predator species abundance can cause variable selection pressure on warning signaling  
724 prey. *Ecol. and Evol.* 2:1971–1976.

725 Vorobyev M, Osorio D. 1998. Receptor noise as a determinant of color thresholds. *Proc. R.*  
726 *Entomol. Soc. Lond. A.* 265:315–358.

727 Vorobyev M, Osorio D, Bennett ATD, Marshall NJ, Cuthill IC. 1998. Tetrachromacy, oil  
728 droplets and bird plumage colors. *J. Compar. Physiol. A.* 183:621–633.

729 Wilson JS, Williams KA, Forister ML, von Dohlen CD, Pitts JP. 2012. Repeated evolution in  
730 overlapping mimicry rings among North American velvet ants. *Nat. Com.* 13:1272.

731 Young OP. 2014. Vertebrate predation on Silphidae (Coleoptera): a literature review. *The*  
732 *Coleopt. Bull.* 68:221-234  
733 Way MJ, Khoo KC. 1992. Role of ants in pest management. *Ann. Rev. Entomol.* 37:479-503.  
734 Weller S, Jacobson N, Conner W. 1999. The evolution of chemical defences and mating  
735 systems in tiger moths (Lepidoptera: Arctiidae). *Biol. J. Linn. Soc. Lond.* 68:557-578.  
736  
  
737  
738