

Children can optimally integrate multisensory information after a short action-like mini game training

Elena Nava<sup>1</sup> \* <sup>°</sup>, Julia Föcker<sup>2</sup> & Monica Gori<sup>3</sup>

\* Corresponding author: Elena Nava

<sup>1</sup> University of Milan-Bicocca

Department of Psychology

Piazza dell'Ateneo Nuovo 1

20126 Milan, Italy

email: [elena.nava@unimib.it](mailto:elena.nava@unimib.it)

<sup>°</sup> NeuroMi, Milan Center for Neuroscience, Milan, Italy

<sup>2</sup> Ludwig-Maximilians-University

College of Social Science

University of Lincoln

Brayford Pool, Lincoln, UK

<sup>3</sup> Istituto Italiano di Tecnologia

Via Enrico Melen 82

Genoa, Italy

### **Conflict of Interest**

All authors declare no conflict of interest.

### **Acknowledgments**

We would like to thank Martina Solbiati and Federica Turazzi for their amazing help in recruiting and testing the children involved in the study. We would also like to thank all children and their families for their willing participation in this study.

## Highlights

- Adults combine multisensory estimates into a coherent multimodal percept
- At least until 8-10 years of age, children do not integrate in an optimal fashion
- We show that a short training with action-like mini games can push the boundaries of developmental plasticity in 4-5 year-olds, who learn to optimally integrate multimodal information
- A short training with action-like mini games also enhances visuo-spatial attention skills
- These plastic changes persist up to 3 months following the end of the training

## Abstract

Combining information across different sensory modalities is of critical importance for the animal's survival and a core feature of human's everyday life.

In adulthood, sensory information is often integrated in a statistically optimal fashion, so that the combined estimates of two or more senses are more reliable than the best single one. Several studies have shown that young children use one sense to calibrate the others, which results in unisensory dominance and undermines their optimal multisensory integration abilities.

In this study we trained children aged 4-5 years with action-like mini games, to determine whether it could improve their multisensory as well as their visuo-spatial skills. Multisensory integration abilities were assessed using a visuo-haptic size discrimination task, while visuo-spatial attention skills were investigated using a multiple object tracking task (MOT).

We found that 2-weeks training were sufficient to observe both optimal multisensory integration and visuo-spatial enhancements selectively in the group trained with action-like mini games. This plastic change persisted up to 3 months, as assessed in a follow-up.

Our novel findings reveal that abilities that are commonly known to emerge in late childhood can be promoted in younger children through action-like mini games and have long-lasting effects. Our data have clinical implications, in that they suggest that specific trainings could potentially help children with multisensory integration deficits.

## Keywords

plasticity; video games; multisensory integration; visuo-spatial skills; learning; development

## Introduction

Adults typically integrate multisensory information in a statistically optimal fashion, that is, the information sources are weighted according to their reliabilities depending on the context, so that the combined estimates are more reliable than the best single source of information (Parise, Spence & Ernst, 2012). Developmental studies have shown that optimal multisensory integration has a protracted time course of development, becoming mature only around the age of 8-10 years (Gori, Del Viva, Sandini & Burr, 2008; Petrini, Remark, Smith & Nardini, 2014; Nardini, Jones, Bedford & Braddick, 2008). For example, Gori et al. (2008) measured visuo-haptic integration abilities of form perception in children aged 5 to 10 years by asking them to discriminate the height of two blocks presented in sequence and found that only older children were able to optimally integrate visuo-haptic information, while younger children relied more on haptic cues. Similar findings were found in comparable age groups using different tasks (Gori, Sandini & Burr, 2012; Nardini et al., 2008), thus suggesting that lack of multisensory integration may represent a distinctive (cognitive) feature before age 10. This late multisensory development has been interpreted as a trade-off between integration benefits and the child's body size and sensory systems continuously recalibrating and adjusting (Burr & Gori, 2012). This weighting of the sensory systems is indeed a signature of the heightened plasticity of the developing brain, but to date, no study has addressed to what extent it is possible to 'push the boundaries' of such plasticity.

Here we assessed this issue by training children aged 4-5 years with a short action-like mini game training for 2 weeks (45 min/day). Children were assigned to one of the following training groups: action video games (AVG), non-action video games (NAVG). A third group (CTRL) served as control group to assess re-test effects.

The use of video games was motivated by the growing literature documenting striking changes in adult action-video-game players (Bavelier, Green, Pouget & Schrater, 2012; Green & Bavelier, 2003). In particular, these studies show behavioural improvements in several cognitive domains, mostly involving visuo-spatial attention, but also memory, and executive control (Bediou, Adams, Mayer, Tipton, Green & Bavelier, 2018). These skills appear to improve in children too following action video games play, as shown in studies conducted in typically (Dye & Bavelier, 2004), as well as atypically developing populations (Franceschini, Gori, Ruffino, Viola, Molteni & Facoetti, 2013; Gambacorta, Huang, Vedamurthy, Nahum, Bayliss, Bavelier & Levi, 2014). In particular, the study of Franceschini et al. (2013) has shown that exposing children with dyslexia for only 12 hours to action-like mini games can dramatically improve their reading skills, suggesting a possible transfer of learning across very different skills that are mediated by attention.

Interestingly, there appears to be a bidirectional interplay between attention and multisensory processing (Talsma, Senkowsi, Soto-Faraco & Woldorff, 2010). For example, temporally and spatially aligned multisensory inputs have a higher probability to capture one individual's attention, as shown in infant studies revealing that redundant and synchronous multisensory stimuli recruit attention and facilitate

perceptual differentiation more than presenting the same information from a single modality (Bahrick & Lickliter, 2000). However, also top-down attention facilitates the integration of multisensory stimuli, as shown by behavioural and neuroimaging studies that have identified several higher level factors, such as voluntarily oriented spatial attention and semantic congruency, which can influence whether and how integration across senses occurs (Welch, 1999).

In this study, by taking advantage of the benefits of video games on cognitive improvements (Bavelier et al., 2012; Franceschini et al., 2013), we investigated whether multisensory integration and visuo-attentional skills can be enhanced following a short action-like mini game training in children aged 4-5 years using a simplified version of the paradigm adopted by Gori et al. (2008), and a multiple object tracking task (MOT) adapted from a study by Green & Bavelier (2006). Because video games are multisensory in nature (i.e., players use tactile, visual and auditory information to achieve their purpose during playing), we predicted improvements in multisensory integration and attention skills particularly in the group of children trained with action-like mini games. The reason is that in action video games, the feedback provided by the sensory modalities is more relevant to the aim of the game (i.e., winning), in comparison to non-action video games. In the latter, players are not pressured by time, thus paying attention to the feedback provided by the concurrent presentation of the sensory modalities is not strictly necessary to the aim of the game. That is, even though both action and non-action video games possess multisensory features, it is only in the action video games that multisensory feedback is functional to achieve a goal.

## **Method**

### *Participants*

19 males and 22 females (mean age = 5.2, range 4.5-6.2 years) were recruited to take part in this study from kindergartens around the city of Milan (Italy). All children had no cognitive or neurological impairment. All but four children were right-handed, and 5 wore glasses. The parents of all children gave their informed consent before the experiment started. The study was approved by the ethical committee of the University of Milano-Bicocca, in accord with the Declaration of Helsinki.

Each child was assigned to one of the following three groups: action-like video games (AVG), non-action-like video game (NAVIG) and controls (CTRL). In order to obtain homogeneous groups (i.e., groups of children with comparable experience playing video games), we administered a questionnaire to the parents assessing the familiarity of their child with video games. The questionnaire was developed by the experimenters in order to address the familiarity of the children with video games and the amount of time spent playing (see Supplementary Material). The questionnaire simply assessed when the child started playing video games (if ever), intended as actively playing on, e.g., tablet, and for how long the child plays daily. We defined as “active playing” those activities involving playing video games in first-person and not, for example, passively watching TV or videos on YouTube.

## Optimal multisensory integration after video game training

The results of the questionnaire showed that none of the children played video games before the age of 4 years, and none of the children spent over 60 minutes a day playing.

We then labelled the children according to the overall time spent per day playing video game: “Intense Players” (N = 9, 30-60 minutes a day), “Medium Players” (N = 14, maximum 30 minutes a day), and “Non Players” (N = 16, no experience playing video games).

The training groups were then balanced, so that an equal number of intense, medium and no players were assigned to each group: N = 14 children were assigned to the AVG group (4 intense players, 5 medium players and 5 non-players), N = 13 children were assigned to the NAVG group (3 intense players, 5 medium players and 5 non-players), N = 14 children were assigned to the CTRL group (4 intense players, 4 medium players and 6 non-players). Table 1 reports a summary about the number of children assigned to each group. The sample size was determined based on previous studies using a Bayesian approach to the data analysis (see Gori et al., 2008). Note that, in comparison to Gori et al. (2008), each group in our study included over 10 children in order to achieve more power and a total of 41 children between 4 and 6 years of age.

### *Description of the tasks*

#### *Multisensory task*

The multisensory size-discrimination task was adapted from Gori et al. (2008) and consisted in presenting children with pairs of wooden blocks of fixed width (97 mm) but variable height, ranging between 45 and 65 mm (with 2 mm increments, see Fig. 1). The blocks were attached on a vertical wooden surface, which was designed to resemble the original set-up of Gori et al. (2008).

All children performed an overall 60 trials, divided into 3 conditions: unimodal visual, unimodal haptic, and visuo-haptic. In the unimodal visual condition, the experimenter presented for about 3 seconds the first block in front of the child, sit at about 45 cm from the stimulus at eye-level. To prevent the child from seeing the experimenter changing the block, the experimenter covered the child’s eyes with the hand. Then, the child had another ca. 3 seconds to look at the block. The experimenter then asked which of the two blocks was taller, and the response of the child (i.e., “first” or “second” responses) was manually recorded by the experimenter.

In the unimodal haptic condition, the blocks were attached behind the wooden surface facing the child, so that she/he could only touch without seeing them. The procedure was the same as for the unimodal visual condition, with the only difference that children could keep their eyes open throughout the testing.

In the visuo-haptic condition, children were presented with two blocks at the same time, one that they could see, and one that they could touch of same size.

The conditions were performed in separate sessions and were counterbalanced across children.

In all conditions, the standard block was always 55 mm high, and was presented either first or second with respect to the other block. The task took overall 30 minutes to complete.

### *Multiple Object Tracking (MOT) task*

In this task, children had to keep track of the spatial position of dynamic target stimuli, embedded among dynamic distractors. The target stimuli consisted in 0.7 cm diameter blue cartoon faces, with a sad facial expression, which appeared among 12 distractors, represented by yellow happy faces (see Fig. 2). The stimuli appeared within a 14 cm diameter grey circle on a portable 15-inches HP laptop. The target and distractors moved randomly within the circle at a velocity of ca. 5°/sec.

The trial was always started by the experimenter, who pressed a button on the keyboard. The 12 distractors appeared, followed shortly after by the target stimulus that remained visible for 2 sec. During this time (Cueing Phase), children were instructed to keep track of the target stimulus until it turned into a yellow face too, thus leaving all faces indistinguishable (Tracking Phase). In the tracking phase, children were asked to track the previously marked blue target items for a duration of 5 seconds. Once the stimulus had changed into a distractor, a “?” appeared on the screen, and children had to tell whether in that specific spatial location there was a target before. Children had two buttons of the keyboard that could be pressed: one that corresponded to a “yes”, and one to a “no” response in relation to the target stimulus. The next trial would start only after the child response; thus, the experimenter encouraged the participants to enter a response even if they were uncertain. On average, children took a maximum of 5 seconds to respond.

The task followed a staircase method, by which following 3 consecutive correct responses, the number of target stimuli would increase of 1 stimulus, up to a maximum of 7 stimuli. In other words, following 3 consecutive correct responses provided by the child, the number of target stimuli would increase to 2; after another 3 correct responses the number of target stimuli would increase to 3, and so on. This staircase converges to 65% in terms of performance (see Brown, 1996, Table 1).

If the child gave two incorrect responses in a row, the number of target stimuli would decrease of 1 stimulus. The threshold would be reached after 5 reversal (i.e., the change in number of target stimuli presented), or after a maximum number of 40 trials. The task lasted about 15 minutes, and the overall experimental session for each child lasted approximately one hour, including at least one break between the two tasks.

### *General procedure*

The tasks were administered before and after the two-weeks training. Note that the children only played during week-days, so overall 10 days. Both tasks were administered to each child and they were counterbalanced among children, so that half of the children started with the multisensory task, and the other half with the MOT task.

The testing was conducted by two expert experimenters, who administered them inside the kindergartens which provided a quiet room. The first testing lasted one week, after which the training immediately started. The training consisted in the AVG and NAVG participants playing for two consecutive weeks with video games of different types (see below for a list of video games used in both trainings) for 45 minutes/day. The training was supervised by the two expert experimenters, who made sure that the children played with the

## Optimal multisensory integration after video game training

video games for the exact amount of time, on the specific video games assigned to each group (see below the list of video games used).

The control group only served to control for re-test effects and thus did not make any training between the two experimental sessions.

Following the training, the two tasks were administered again, and again it took ca. one week to test all children.

### *List of video games played:*

The video games were classified as action or non-action mini games according to the characteristics provided by Green, Li, and Bavelier (2009), who classified action video as all sharing a set of qualitative features, including “extraordinary speed (both in terms of very transient events and in terms of the velocity of moving objects), a high degree of perceptual, cognitive, and motor load in the service of an accurate motor plan (multiple items that need to be tracked and/or kept in memory, multiple action plans that need to be considered and quickly executed typically through precise and timely aiming at a target), unpredictability (both temporal and spatial), and an emphasis on peripheral processing (with important items most often appearing away from the centre of the screen)”. It is important to note that all the video games we used provided auditory, visual and tactile feedback; that is, children actively touched the tablet, watched the scenes and heard auditory tones.

On the contrary, non-action video games are those that do not require speed (i.e., objects do not move fast) and are not necessarily presented in the periphery. Non-action video games are more accuracy-based and time does not pressure the participants. Most importantly though, we selected the video games so that none contained violent scenes or actions. In other words, in none of the selected video game there was any shooting, blood, death, or anything that could remind the hurting of individuals. All video games were played on an iPad during the training.

For the action-like mini games training we used the following games:

FRUIT NINJA, SMASH MASTER, NINJUMP, RAIL RUSH, BRICKS BREAKER RE, BANANA KONG, PJ. MASKS: MOONLIGHT HEROES, FRUIT SMASHER, LEP’S WORLD 3, LIGHTNING McQUEEN, TRAIN CONDUCTOR WORLD, ROLLING SKY, MINION RUSH, JUMP FEVER.

For the non-action-like mini games training, we used the following games:

PUZZINGO, BUBBLE SHOOTER, DRAWING for KIDS, KIDS MAZES, MEMORY GAME, DOCTOR KIDS, ANGRY BIRDS, ANIMALS PUZZLE, FROSTY SHOT, DIRTY FARM, PLAYING CARDS for KIDS, FIND THE DIFFERENCE, COLOURING BOOKS, MASHA and the BEAR.

## Results

### *Multisensory task*

Several studies have shown that adults integrate visual and haptic information (and information from other modalities) in a statistically optimal fashion, weighting each sense according to its reliability (Deneve & Pouget, 2004; Ernst & Banks, 2002; Parise, Spence & Ernst, 2012). The maximum likelihood estimate (MLE) predicts as main signature for cross-modal integration that the thresholds for dual-modality presentation is lower than either visual or haptic thresholds. This model, described below, is a perfect way to quantify the level of multisensory integration in children. Since Gori et al. (2008) demonstrated that MLE predicts visual-haptic multisensory integration only after 8-10 years of age, we used this model to quantify the improvement of multisensory integration after the training with video games. To make the MLE calculation, we first extracted the threshold for each unimodal condition (visual and haptic alone) by calculating the proportion of trials in which the child responded that the standard stimulus was taller than the other variable stimuli and by fitting for each child the data into a cumulative Gaussian function, in which the obtained standard deviation estimated the discrimination threshold.

To assess optimal integration, we compared the discrimination thresholds with the threshold predicted by the Maximum Likelihood Estimation model before and after the training. The MLE calculation assumes that for size judgments, the optimal bimodal MLE prediction for the visuo-haptic threshold ( $\sigma_{vH}$ ) is given by:

$$\sigma_{vH}^2 = \frac{\sigma_v^2 \sigma_H^2}{\sigma_v^2 + \sigma_H^2} \leq \min(\sigma_v^2, \sigma_H^2)$$

where  $\sigma_v$  and  $\sigma_H$  are the visual and haptic unimodal thresholds extracted from the psychometric function. The improvement is greatest ( $\sqrt{2}$ ) when  $\sigma_v = \sigma_H$ . The prediction of the model is good when the bimodal threshold predicted from the unimodal estimates as per the equation above is not different from the measured bimodal hreshold. The critical test for optimal multi-sensory integration is that the measured bimodal threshold be as good as the MLE estimate of the bi-modal threshold, derived from optimally combining the unimodal performances.

Following this model, we compared the thresholds in all modalities to the MLE prediction by means of paired t-tests, and found that in the visuo-haptic modality, before starting the training, all groups of children presented higher thresholds than the MLE prediction (all  $p < 0.003$ ), suggesting no optimal multisensory integration capabilities. However, after the training, a specific improvement was observed in the group of

## Optimal multisensory integration after video game training

children that performed the action-like mini-game training. Indeed, while the NAVG [ $t(12) = 2.39, p = 0.03$ ] and CTRL group [ $t(13) = 4.45, p < 0.001$ ] still showed higher measured threshold than the MLE prediction, the AVG group showed optimal multisensory integration, as their measured thresholds did not differ from the MLE prediction ( $p = 0.10$ ).

The comparison of the haptic and visual modalities alone with the MLE prediction showed that all groups presented higher thresholds than the MLE prediction before and after the training (all  $p < 0.003$ ).

We also conducted a repeated measures ANOVA on the discrimination thresholds of the three groups of children, with Modality (haptic alone, visual alone and visuo-haptic) and Session (pre- and post-training) as within-subjects factors, and Group (AVG, NAVG and controls) as between-subjects factor.

The analysis revealed a main effect of Session [ $F(1, 38) = 11.94, p = 0.001$ ] and Modality [ $F(2, 76) = 14.13, p < 0.001$ ], a Session X Group interaction [ $F(2, 38) = 4.02, p = 0.03$ ], due to the AVG group performing better after the training ( $p = 0.003$ ), while the other two groups did not show any difference between sessions (both  $p > 0.99$ ). There was also a Modality X Group [ $F(4, 76) = 3.46, p = 0.01$ ], and a Session X Modality interaction [ $F(2, 76) = 3.48, p = 0.04$ ]. To further explore whether the performance on each modality changed across groups, we conducted other ANOVAs on each modality. The analysis conducted on the visuo-haptic thresholds showed a main effect of Group [ $F(2, 38) = 6.96, p = 0.003$ ], a main effect of Session, [ $F(1, 38) = 27.97, p < 0.001$ ], and a Group X Session interaction [ $F(2, 38) = 5.81, p = 0.006$ ], the latter caused by children only in the AVG group improving their visuo-haptic discrimination abilities following training ( $p < 0.001$  on Bonferroni post-hoc test). On the contrary, the performance of the other two groups did not vary following either the non-action video game training ( $p = 0.38$ ) or the re-test ( $p = 0.99$ , see Fig. 3a, upper left panel for a summary of results, and Fig. 4 for psychometric functions of the whole three groups).

Furthermore, analysis on both the haptic and visual modality alone did not reveal any main effect or interaction in any group and in any session (all  $p > 0.14$ ). That is, the training specifically improved multisensory integration abilities, but not the performance in the single modality.

### *MOT task*

Because less than 50% of the children reached the threshold, for each child we calculated the average number of target stimuli she/he could track on each trial. This average was calculated on the correct responses only. Thus, the mean number of tracked objects was entered as dependent variable in a repeated measure ANOVA, with the within subject factor Session and the between subject Group as main factors. The analysis revealed a main effect of Session [ $F(1, 38) = 24.99, p < 0.001$ ], Group [ $F(2, 38) = 3.25, p = 0.04$ ], and a Session X Group interaction [ $F(2, 38) = 9.24, p < 0.001$ ], caused by the AVG group increasing the

## Optimal multisensory integration after video game training

average number of object tracked following the training (pre-training:  $M = 1.86$ ,  $SD = 0.58$ ; post-training:  $M = 3.04$ ,  $SD = 0.59$ ,  $p < 0.001$ ) in comparison to the NAVG (pre-training:  $M = 1.88$ ,  $SD = 0.53$ ; post-training:  $M = 2.34$ ,  $SD = 0.72$ ,  $p = 0.39$ ) and the control group (pre-training:  $M = 2.05$ ,  $SD = 0.50$ ; post-training:  $M = 2.08$ ,  $SD = 0.40$ ,  $p = 0.99$ ). Importantly, no difference across the three groups emerged in the testing conducted before the training (all  $p = 0.99$ , see Fig. 3b, upper panel for a summary of results).

To observe whether changes in visuo-spatial attentional skills could subserved changes observed in multisensory integration, we performed correlations between the two tasks. This was done by subtracting the mean performance in the post-training phase to the pre-training phase performance separately for group. However, in all groups, we did not find any correlation between the two tasks (all  $p > 0.22$  on two-tailed Pearson correlation).

### **3 month-follow-up**

#### *Participants, tasks and procedure*

To assess the stability of the changes observed selectively in the AVG group, we conducted a follow-up in this same group of children after 3 months from the end of the training.

Only  $N = 13$  children of the AVG group took part, because 1 child could not participate. This child was removed from the analyses aimed at comparing the three experimental sessions (pre-training, post-training and follow-up). Note that the questionnaire regarding the familiarity with video game use was also administered again to the families of the children, to make sure that their playing pattern did not vary within these 3 months. Indeed, they confirmed that the pattern did not vary in any child.

The stimuli, tasks and procedure were identical to the other sessions. The children were again tested in the same kindergarten they were tested in the previous sessions.

### **Results of the 3 month-follow up**

The analyses conducted in the follow-up revealed that the measured threshold of the AVG in the follow-up did not differ from the MLE prediction ( $p = 0.34$ ), thus showing optimal multisensory integration.

Furthermore, to observe differences across sessions, we performed a repeated measure ANOVA with Session (pre-training, post-training and 3-month-follow-up) as only factor. This analysis revealed a main effect of Session [ $F(2, 24) = 13.95$ ,  $p < 0.001$ ], caused by differences in visuo-tactile integration between the pre-training and the post-training ( $p < 0.001$ ), the pre-training and the follow-up ( $p = 0.01$ ), but not between the post-training and the follow-up ( $p = 0.13$ ). That is, the improvement in multisensory integration was maintained after 3 months following the end of the training (see Fig. 3a, lower panel for a summary of results).

Stability of changes were also observed in the MOT, though less robust than in the multisensory task. That is, we found a main effect of Session [ $F(2, 24) = 19.47$ ,  $p < 0.001$ ], caused by a difference in

performance between the pre-and post-training ( $p < 0.001$ ), and between the pre-training and follow-up ( $p = 0.01$ ). Note however, that the follow-up differed from the post-training ( $p = 0.02$ ), suggesting that after 3 months, visuo-spatial enhancement was slightly decaying (follow-up:  $M = 2.47$ ,  $SD = 0.54$ ,  $p = 0.01$ , see Fig. 3b, lower panel for a summary of results).

### Discussion

The results of this study show that a short training with action-like mini games can promote optimal integration of multisensory signals during a particular developmental phase in which this ability has been reported to be sub-optimal.

This suggests that selectively training multisensory abilities aids the developing brain to better combine redundant multisensory estimates into a coherent multimodal percept.

In line with previous studies (Green, Pouget & Bavelier, 2010; Green & Bavelier, 2012), our findings suggest that the enhanced visuo-spatial attention and multisensory processes seen in children in the AVG group may reflect a change in learning strategies. It should be discussed which mechanisms may be responsible for the changes observed. That is, did the training promote better multisensory binding, or did the children learn to better estimate the sensory weights? While our data cannot provide a conclusive answer to these questions, two aspects should be noted. First, it might be that children learned (or improved) the correspondence between the sensory signals. That is, the perceptual system has to determine whether two signals belong to the same object or event before deciding whether to integrate them or not. Thus, it could be that children at this age are yet not able to causally infer that there is a connection between the two senses, and the training may have improved such correspondence.

It could also be claimed that instead of being a purely binding problem, children may have learned to better estimate the sensory weights; this, in turn, might have favoured optimal integration. Optimal integration, indeed, means that signals from unimodal cues are integrated in proportion to their reliability as predicted by Bayes's rule. Estimating the sensory weights may however be a very difficult task for children, as the sensory systems involved with spatial perception must recalibrate continuously during development because of the body rapidly changing in size. As suggested by previous studies (Gori et al., 2008; Burr & Gori, 2012; Gori, 2015), it may be that for the developing child, calibration is more important than optimizing perception by integration; and if sensory information is integrated, one sense cannot be used to calibrate the other. The training with the action-like mini games may have facilitated the binding of information between haptic and visual information but the effect does not appear to be related to a change in the precision of unimodal inputs. Indeed, we did not observe any difference in the precision of the haptic and visual systems before and after the training but only a specific change in the bimodal condition after the training in the group trained with the action video games.

Furthermore, even though our findings may suggest that multisensory integration improvements go hand in hand with visual attention enhancement, it should be noted that we did not find any correlation between the two tasks, which in turn suggests that the improvements observed in the two tasks may reflect different mechanisms underlying attention and multisensory learning skills. In particular, it could be that learning to optimally integrate the senses could occur in a very automatic fashion, without any attentional requirements.

Most importantly, we also found that these plastic changes have long-term effects, at least lasting up to 3 months, and are observed both at a multisensory integration and visuo-spatial level. Importantly, the fact that performance in the MOT after 3 months was less robust than in the multisensory task corroborates the idea that, while multisensory processes are fully automatic and once the association is optimally established it cannot be reversed, attentional processes are more top-down and likely need to be continuously trained in order to be maintained.

It should be acknowledged that, for practical issues, the experimenters who conducted the training were not blind to the purpose of the study. Even though the experimenters were highly trained and were explicitly told that any outcome would have contributed to the knowledge about brain plasticity in children, we cannot completely exclude a slight bias.

In conclusion, these training-induced, long-term plastic changes in the developing brain suggest that, first, multisensory integration is not under maturational constraints, that is, it is not the growing body - which has to recalibrate all the sensory signals in the first years of life – that imposes limits to multisensory development. It rather appears that multisensory integration is protracted throughout late childhood because the environment does not typically impose high demands on the child's brain. However, if the child's brain is given the opportunity to systematically train an ability, it rapidly shows adult-like patterns of multimodal cue integration.

The fact that children can rapidly learn to recalibrate their sensory systems has dramatic implications for all those conditions in which multisensory integration appears pathologically impaired. This is particularly true for autistic (Brandwein et al., 2012), but also for dyslexic children, who have recently been documented to have impaired multisensory integration capabilities (Harrar, Tammam, Pérez-Bellido, Stein & Spence, 2014). Thus, training with action-like mini games could prove a useful, fast and fun way to promote long-lasting plastic multisensory and attentive changes in these populations.

### Data Availability Statement

The data that support the findings of this study are available on request from the corresponding author (elena.nava@unimib.it). Please note that in the Supplementary Figure, all data from each participants in the Multisensory Task are plotted as a function of Time (Pre- and Post-Training) and Condition (Visual, Tactile and Bimodal).

## References

- Bahrick, L. E., & Lickliter, R. (2000). Intersensory redundancy guides attentional selectivity and perceptual learning in infancy. *Developmental Psychology*, *36*, 190.
- Bavelier, D., Green, C. S., Pouget, A., and Schrater, P. (2012). Brain plasticity through the life span: learning to learn and action video games. *Annual Review of Neuroscience*, *35*, 391-416.
- Bediou, B., Adams, D. M., Mayer, R. E., Tipton, E., Green, C. S., & Bavelier, D. (2018). Meta-analysis of action video game impact on perceptual, attentional, and cognitive skills. *Psychological Bulletin*, *144*, 77-110.
- Brandwein, A. B., Foxe, J. J., Butler, J. S., Russo, N. N., Altschuler, T. S., Gomes, H., & Molholm, S. (2012). The development of multisensory integration in high-functioning autism: high-density electrical mapping and psychophysical measures reveal impairments in the processing of audiovisual inputs. *Cerebral Cortex*, *23*, 1329-1341.
- Brown, L. G. (1996). Additional rules for the transformed up-down method in psychophysics. *Perception & Psychophysics*, *58*, 959-962.
- Burr, D., & Gori, M. (2012). *Multisensory integration develops late in humans*. In *The Neural Bases of Multisensory Processes*, Eds. Murray, M.M., & Wallace, M.T. Boca Raton (FL): CRC Press/Taylor & Francis.
- Deneve, S., & Pouget, A. (2004). Bayesian multisensory integration and cross-modal spatial links. *Journal of Physiology-Paris*, *98*, 249-258.
- Dye, M. W., & Bavelier, D. (2004). Playing video games enhances visual attention in children. *Journal of Vision*, *4*, 40.
- Franceschini, S., Gori, S., Ruffino, M., Viola, S., Molteni, M., and Facoetti, A. (2013). Action video games make dyslexic children read better. *Current Biology*, *23*, 462-466.
- Gambacorta, C., Huang, S., Vedamurthy, I., Nahum, M., Bayliss, J., Bavelier, D., & Levi, D. (2014). Action Video Games as a Treatment of Amblyopia in Children: A Pilot Study of a novel, child-friendly action game. *Journal of Vision*, *14*, 665-665.
- Gori, M., Del Viva, M., Sandini, G., and Burr, D.C. (2008). Young children do not integrate visual and haptic form information. *Current Biology*, *18*, 694-698.
- Gori, M., Sandini, G., & Burr, D. (2012). Development of visuo-auditory integration in space and time. *Frontiers in Integrative Neuroscience*, *6*, 77.

- Gori, M. (2015). Multisensory integration and calibration in children and adults with and without sensory and motor disabilities. *Multisensory Research*, 28, 71-99.
- Green, C. S., and Bavelier, D. (2003). Action video game modifies visual selective attention. *Nature*, 423, 534-537.
- Green, C.S., Bavelier, D. (2006). Enumeration versus multiple object tracking: the case of action video game players. *Cognition*, 101, 217-245.
- Green, C. S., Li, R., & Bavelier, D. (2010). Perceptual learning during action video game playing. *Topics in Cognitive Science*, 2, 202-216.
- Green, C. S., & Bavelier, D. (2012). Learning, attentional control, and action video games. *Current Biology*, 22, 197-206.
- Green, C. S., Pouget, A., & Bavelier, D. (2010). Improved probabilistic inference as a general learning mechanism with action video games. *Current Biology*, 20, 1573-1579.
- Harrar, V., Tammam, J., Pérez-Bellido, A., Pitt, A., Stein, J., & Spence, C. (2014). Multisensory integration and attention in developmental dyslexia. *Current Biology*, 24, 531-535.
- Nardini, M., Jones, P., Bedford, R., & Braddick, O. (2008). Development of cue integration in human navigation. *Current Biology*, 18, 689-693.
- Parise, C. V., Spence, C., and Ernst, M.O. (2012). When correlation implies causation in multisensory integration. *Current Biololy*, 22, 46-49.
- Petrini, K., Remark, A., Smith, L., and Nardini, M. (2014). When vision is not an option: children's integration of auditory and haptic information is suboptimal. *Developmental Science*, 17, 376-387.
- Talsma, D., Senkowski, D., Soto-Faraco, S., and Woldorff, M. G. (2010). The multifaceted interplay between attention and multisensory integration. *Trends in Cognitive Science*, 14, 400-410.
- Welch, R. B. (1999). *Meaning, attention, and the "unity assumption" in the intersensory bias of spatial and temporal perceptions*. In *Advances in Psychology*, 129, 371-387. North-Holland.

## Figure Captions

Table 1. Details of the participants: age, training group assigned and level of familiarity with video games.

Fig. 1. Graphical representation of a child performing the multisensory task.

Fig. 2 Stimuli and schematic procedure of the MOT task.

Fig. 3. Results of the multisensory (a) and MOT task (b). The upper panel shows the results of the three groups of children that underwent action video game (AVG), and non-action video game (NAVG) training,

## Optimal multisensory integration after video game training

as well as the controls (CTRL). Only the AVG group showed optimal multisensory integration following the training, as well as enhanced visuo-spatial abilities.

The lower panel shows the long-term effects of the AVG training in a 3 month-follow-up in both the multisensory and MOT task.

Fig. 4. Psychometric functions depicting the cumulative performance of the three groups before (red line) and after (grey line) the training.