



Contents lists available at ScienceDirect

## International Journal of Infectious Diseases

journal homepage: [www.elsevier.com/locate/ijid](http://www.elsevier.com/locate/ijid)

## Pharmacogenetic considerations in the treatment of co-infections with HIV/AIDS, tuberculosis and malaria in Congolese populations of Central Africa



Srinivas Reddy Pallerla<sup>a,b,1</sup>, Darrel Ornelle Elion Assiana<sup>c,d,1</sup>, Le Thi Kieu Linh<sup>a,b</sup>, Frederick Nchang Cho<sup>a,e</sup>, Christian G. Meyer<sup>a,b,f</sup>, Kaossarath Adédjokè Fagbemi<sup>a,g</sup>, Ayola Akim Adegniko<sup>a,h</sup>, Véronique Penlap Beng<sup>i</sup>, Eric A. Achidi<sup>e</sup>, Gauthier Mesia Kahunu<sup>j</sup>, Mathew Bates<sup>k</sup>, Martin P. Grobusch<sup>a,h,1</sup>, Peter G. Kremsner<sup>a,h</sup>, Francine Ntoumi<sup>a,c,d</sup>, Thirumalaisamy P. Velavan<sup>a,b,c,f,\*</sup>

<sup>a</sup> Institute of Tropical Medicine, University of Tübingen, Wilhelmstrasse 27, 72074 Tübingen, Germany

<sup>b</sup> Vietnamese-German Center for Medical Research, VG-CARE, Hanoi, Viet Nam

<sup>c</sup> Fondation Congolaise pour la Recherche Médicale (FCRM), Brazzaville, Congo

<sup>d</sup> Faculty of Sciences and Technology, University Marien Ngouabi, Brazzaville, Congo

<sup>e</sup> Department of Biochemistry and Molecular Biology, Faculty of Science, Laboratory of Infectious Diseases, Faculty of Health Sciences, University of Buea, Buea, Cameroon

<sup>f</sup> Faculty of Medicine, Duy Tan University, Da Nang, Viet Nam

<sup>g</sup> Department of Biomedical Sciences, Laboratory of Cytogenetics and Medical Genetics, Faculty of Health Sciences, University of Abomey-Calavi, Benin

<sup>h</sup> Centre de Recherches Médicales de Lambarene, Lambarene, Gabon

<sup>i</sup> Department of Biochemistry, Faculty of Science, University of Yaoundé 1, Yaoundé, Cameroon

<sup>j</sup> Unit of Clinical Pharmacology and Pharmacovigilance, University of Kinshasa, Kinshasa, Democratic Republic of the Congo

<sup>k</sup> School of Life Sciences, University of Lincoln, Lincoln, United Kingdom

<sup>1</sup> Center for Tropical Medicine and Travel Medicine, Department of Infectious Diseases, Amsterdam University Medical Centers, University of Amsterdam, The Netherlands

## ARTICLE INFO

## Article history:

Received 16 November 2020

Received in revised form 30 November 2020

Accepted 4 December 2020

## Keywords:

Malaria

Tuberculosis

HIV

Pharmacogenetics

Republic of Congo

## ABSTRACT

**Background:** HIV-infection, tuberculosis and malaria are the big three communicable diseases that plague sub-Saharan Africa. If these diseases occur as co-morbidities they require polypharmacy, which may lead to severe drug–drug–gene interactions and variation in adverse drug reactions, but also in treatment outcomes. Polymorphisms in genes encoding drug-metabolizing enzymes are the major cause of these variations, but such polymorphisms may support the prediction of drug efficacy and toxicity. There is little information on allele frequencies of pharmacogenetic variants of enzymes involved in the metabolism of drugs used to treat HIV-infection, TB and malaria in the Republic of Congo (ROC). The aim of this study was therefore to investigate the occurrence and allele frequencies of 32 pharmacogenetic variants localized in absorption distribution, metabolism and excretion (ADME) and non-ADME genes and to compare the frequencies with population data of Africans and non-Africans derived from the 1000 Genomes Project.

**Results:** We found significant differences in the allele frequencies of many of the variants when comparing the findings from ROC with those of non-African populations. On the other hand, only a few variants showed significant differences in their allele frequencies when comparing ROC with other African populations. In addition, considerable differences in the allele frequencies of the pharmacogenetic variants among the African populations were observed.

**Conclusions:** The findings contribute to the understanding of pharmacogenetic variants involved in the metabolism of drugs used to treat HIV-infection, TB and malaria in ROC and their diversity in different populations. Such knowledge helps to predict drug efficacy, toxicity and ADRs and to inform individual and population-based decisions.

© 2020 The Author(s). Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

\* Corresponding author at: Institute of Tropical Medicine, University of Tübingen, Wilhelmstrasse 27 72074 Tübingen Germany.  
E-mail address: [velavan@medizin.uni-tuebingen.de](mailto:velavan@medizin.uni-tuebingen.de) (T.P. Velavan).

<sup>1</sup> Equal contribution.

## Background

HIV/AIDS, tuberculosis (TB) and malaria, the big three killers, are diseases with high prevalences and considerable mortality in many developing countries. The world-wide prevalence of these diseases is over 250 million, with sub-Saharan Africa accounting for 71%, 25% and 92% of HIV/AIDS, TB and malaria, respectively (Chaudhry et al., 2016). As in many other sub-Saharan African countries, transmission of the malaria-causing parasite *Plasmodium falciparum* is high in the Republic of Congo (ROC) (WHO, 2015b), together with a high burden of TB and HIV-infection (Linguissi et al., 2017). These diseases overlap almost completely, leading to a high rate of concurrent occurrence of HIV/TB, TB/malaria and HIV/TB/malaria; however, incidences vary widely between individuals and communities. Comorbidities require a polypharmaceutical approach, which may lead to drug–drug–gene interactions and drug-induced pathology, variations in adverse drug reactions (ADR) and therapeutic efficacy as well as the development of drug resistances (Alomar, 2014; Bassi et al., 2017; Rattan et al., 1998; Turner and Wainberg, 2006). It has recently been reported that the 25 most common drugs causing ADR in Africa are drugs used to treat TB, malaria and HIV-infections (Rajman et al., 2017).

Treatment of TB involves combinations of drugs to be taken over an extended period of time and in HIV-infection, the respective drug combinations are taken lifelong. The first-line treatment of HIV-infection includes highly active antiretroviral (HAART) drugs, with dolutegravir, efavirenz, raltegravir and nevirapine being recommended by WHO (2019). Standard first-line TB treatment consists of isoniazid, rifampicin, pyrazinamide and ethambutol for two months, followed by isoniazid and rifampicin for another four months. For the treatment of uncomplicated malaria, artemisinin combination therapies (ACTs) such as artesunate-amodiaquine, artemether-lumefantrine and other combinations are the most commonly used drugs used in sub-Saharan Africa (WHO, 2015a). When multiple drugs are administered, the treatment outcomes, in particular the efficacy, ADRs and the degree of emergence of drug resistance vary at the individual and population level. These variations are influenced by the age, sex and weight of patients, dosage, comorbidities and genetic factors. It is known that up to 95% of variations in treatment outcomes can be explained by genetic factors alone (Ross et al., 2012). Genetic studies at the population level and the 1000 Genome Project Phase 3 indicate that African ethnic groups are the most genetically diverse populations (Tishkoff et al., 2009). Regional allele frequencies of African genomes also show a high degree of variation (Gurdasani et al., 2015; Ramos et al., 2014).

Genetic variations in drug absorption, distribution, metabolism and excretion (ADME) genes both within and between populations can influence the pharmacokinetics and pharmacodynamics of drugs (Ma et al., 2002; Wilson et al., 2001) and lead to differences in treatment outcomes, drug resistance and increased toxicity (Li et al., 2011). Pharmacogenomic studies of drugs used to treat the three diseases have been extensively conducted and variants in ADME and non-ADME genes have been found associated with treatment outcomes and toxicity. Some of the common genes include *CYP2B6*, *CYP3A5*, *CYP3A4*, *CYP2A6*, *ABCB1*, *NR1I3*, *UGT2B7*, *NAT2*, *CYP2E1*, *GSTMI*, *GSTTI*, *SLCO1B1* and *UGT1A9* (Bains, 2013; Chaudhry et al., 2016). While translation of pharmacogenetic information into clinical practice has been slow, pharmacogenomic drug labels are increasingly being approved by the US Food and Drug Administration (FDA), the European Medicines Agency (EMA), Swiss Agency of Therapeutic Products (Swissmedic) and the Pharmaceuticals and Medical Devices Agency, Japan (PMDA) (Whirl-Carrillo et al., 2012).

There are very few studies (Peko et al., 2019a, b) available so far on the allele frequencies of pharmacogenetic variants associated with the outcomes of the drugs used to treat the three big diseases in the Republic of Congo (ROC). The main objective of the present study is to identify allele frequencies of recognized pharmacogenetic variants that are associated with treatment outcomes of HIV/AIDS, TB and malaria in the ROC and to compare the frequencies with those in other African and non-African populations. Such knowledge helps to predict drug efficacy, toxicity and ADRs and inform individual and population-based decisions.

## Materials and methods

### Study population

Ethical approval was obtained from the Institutional Ethics Committee of the Fondation Congolaise pour la Recherche Médicale in Brazzaville, ROC. Following written informed consent from parents/guardians, a total of 67 children aged between one and ten years with fever were included randomly out of a cohort which was recruited for a previous study at the Marien Ngouabi Hospital in the north of Brazzaville (Etoka-Beka et al., 2016; Gampio Gueye et al., 2019). Two millilitres of venous blood were available in EDTA tubes stored at  $-20^{\circ}\text{C}$ .

### SNP variants and selection criteria

SNPs with functional and clinical significance for drugs used to treat HIV-infection, TB and malaria were selected from the Pharmacogenomics Knowledge Base (PharmGKB; [www.pharmgkb.org](http://www.pharmgkb.org)). SNPs with allele frequencies  $>10\%$  known to occur in African populations based on the data from 1000 Genome Project were considered in our study.

The custom ordered TaqMan™ SNP panel which was designed and applied included 32 SNPs located in 18 ADME and non-ADME genes; (Table 1). ADME genes included those encoding ATP Binding Cassette Subfamily B Member 1 (*ABCB1*), cytochrome P450 family (*CYPs*) *CYP2A6*, *CYP2B6*, *CYP2D6*, *CYP2C19*, *CYP2C8*, *CYP3A4*, *CYP3A5*, N-acetyltransferase 2 (*NAT2*), solute carrier organic anion transporter family member 1B1 (*SLCO1B1*), UDP-glucuronosyltransferase family (*UGT1A1*). Non-ADME genes included those encoding glucose-6-phosphate dehydrogenase (*G6PD*), nuclear factor Kappa B subunit 1 (*NF-κB1*), tumor necrosis factor (*TNF-α*), nuclear receptor subfamily 1, group 1, member 2 (*NR1I2*), cut like homeobox 2 (*CUX2*) and ATP/GTP binding protein like 4 (*AGBL4*).

### DNA isolation, genotyping and quality control

Genomic DNA was extracted from the blood samples using the QIAamp DNA Minikit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. The quantity and quality of DNA was determined with the NanoDrop™ (Thermo Fisher Scientific Inc.) and Qubit™ 4 Fluorometer using the Qubit™ dsDNA BR Assay Kit (Thermo Fisher Scientific, Waltham, MA, USA). DNA concentrations ranged between 1–40 ng/μL. Genotyping of the 32 SNPs (Thermo Fisher assay IDs; Supplementary Table 1) were performed by allele discrimination with TaqMan® SNP Genotyping Assays on a Thermo QuantStudio5™ Real-Time 384 well PCR system (Thermo Fisher Scientific, Life Technologies, Carlsbad, USA) according to the manufacturer's protocol. 5 μL final reaction volume was used with a recommended DNA concentration of 0.2 ng/μL. Data obtained was loaded into the TaqMan® Genotyper Software (Thermo Fisher, Life Technologies, Carlsbad, USA) to automatically retrieve the genotyping data. Hardy–Weinberg Equilibrium (HWE) was calculated and only SNPs that passed underwent statistical analysis.

**Table 1**

ADME and non-ADME gene variants with clinical significance in drugs used to treat malaria, tuberculosis and HIV-infection.

Gene	SNP ID	Defining SNP	Phenotype/Association	HIV, TB and Malaria drug substrates	Refs
Absorption, distribution, metabolism and excretion (ADME)					
<i>ABCB1</i>	rs1045642	<i>ABCB1</i> *6_3435C>T (I1145I)	Null	Rifampicin, Efavirenz, Nevirapine, Nelfinavir, Atazanavir and Ritonavir	Ciccacci et al. (2010); Park et al. (2010); Yimer et al. (2011)
	rs3842	<i>ABCB1</i> _c193A>G (3'UTR)	Not defined	Efavirenz	Elens et al. (2010)
<i>CYP2A6</i>	rs28399433	<i>CYP2A6</i> *9_-48T>G (Promoter)	Reduced function	Efavirenz	Shahi et al. (1990); Soeria-Atmadja et al. (2017)
<i>CYP2B6</i> <sup>a</sup>	rs28399499	<i>CYP2B6</i> *18_c.983T>C (I328T)	Reduced function	Efavirenz and Nevirapine	Ciccacci et al. (2013); Gandhi et al. (2012)
	rs3745274	<i>CYP2B6</i> *6_c.516G>T (Q172H)	Reduced function	Efavirenz and Nevirapine	Cusato et al. (2016); Sinxadi et al. (2015)
<i>CYP2D6</i> <sup>a</sup>	rs1065852	<i>CYP2D6</i> *10_c.100T>C (S34P)	Intermediate function	Hydroxychloroquine, Tafenoquine and Primaquine	Bennett et al. (2013); Saito et al. (2018); St Jean et al. (2016)
<i>CYP2C19</i> <sup>a</sup>	rs12768009 <sup>b</sup>	<i>CYP2C19</i> _c.168+3235G>A (Intron)	Reduced function	Nevirapine	Lehr et al. (2011)
	rs4244285	<i>CYP2C19</i> *2_19154G>A (P227P)	Reduced function	Nelfinavir	Haas et al. (2005); Kattel et al. (2015)
<i>CYP2C8</i>	rs11572103	<i>CYP2C8</i> *2_c.805A>T (I269F)	Reduced function	Amodiaquine	Parikh et al. (2007)
<i>CYP3A5</i> <sup>a</sup>	rs776746	<i>CYP3A5</i> *3_6986A>G (Splice defect)	Loss of function	Lumefantrine, Atazanavir, Ritonavir and Nevirapine	Brown et al. (2012); Castillo-Mancilla et al. (2016); Mutagonda et al. (2017)
	rs10264272	<i>CYP3A5</i> *6_14690G>A (K208K)	Reduced function		
	rs41303343	<i>CYP3A5</i> *7_27132insA (-346S)	Reduced function		
<i>CYP3A4</i> <sup>a</sup>	rs2740574	<i>CYP3A4</i> *1B_-392A>G (Intergenic)	Reduced function	Efavirenz, Atazanavir, Lumefantrine and Indinavir	Bertrand et al. (2009); Haas et al. (2005); Kile et al. (2012); Mutagonda et al. (2017)
<i>NAT2</i> <sup>a</sup>	rs1801280	<i>NAT2</i> *5_341T>C (I114T)	Slow acetylator function	Isoniazid/TB treatment	Ben Mahmoud et al. (2012); Chan et al. (2017); Shi et al. (2015); Singla et al. (2014)
	rs1799930	<i>NAT2</i> *6_590G>A (R197Q)	Slow acetylator function		
	rs1799929	<i>NAT2</i> *11_481C>T (L161L)	Association		
	rs1208	<i>NAT2</i> *12_803A>G (K268R)	Fast acetylator function		
	rs1041983	<i>NAT2</i> *13_282C>T (Y94Y)	Fast acetylation function		
	rs1495741	<i>NAT2</i> _18415371G>A (Tag SNP)	Association	Isoniazid/TB treatment	Chamorro et al. (2016); Ho et al. (2013)
<i>SLCO1B1</i>	rs2306283	<i>SLCO1B1</i> *1b_388A>G (N130D)	Enhanced function	Rifampicin and Efavirenz	Kwara et al. (2014)
	rs4149032	<i>SLCO1B1</i> *5_c.85-7793C>T (Intron)	Reduced function	Rifampicin and Isoniazid	Calcagno et al. (2019); Chigutsa et al. (2011)
<i>UGT2B7</i>	rs7439366 <sup>b</sup>	<i>UGT2B7</i> *2_2100C>T (Y268H)	Reduced/Enhanced function (Substrate dependent)	Efavirenz	Cusato et al. (2016); Kwara et al. (2009)
<i>UGT1A1</i> <sup>a</sup>	rs3064744	<i>UGT1A1</i> *28_ TA (Indel)	Reduced function	Atazanavir, Dolutegravir, Ritonavir and Raltegravir	Johnson et al. (2014); Vardhanabhuti et al. (2015)
	rs10929303	<i>UGT1A1</i> _1813C>T (3'UTR)	Reduced function	Atazanavir and Ritonavir	Nishijima et al. (2014)
Non-ADME					
<i>G6PD</i> <sup>a</sup>	rs1050828	<i>G6PD</i> _c.202G>A (V68M)	Loss of function	Chlorproguanil, Dapsone, Mefloquine, Amodiaquine, Pyrimethamine and Sulfadoxine	Fanello et al. (2008); Sicard et al. (1978)
	rs1050829	<i>G6PD</i> _c.376A>G (N126D)			
<i>NF-κB1</i>	rs4647992	<i>NF-κB1</i> _c.159+305C>T (Intron)	Association	TB treatment	Zhang et al. (2019)
<i>TNF-α</i>	rs1800629	<i>TNF-α</i> _ -308G>A (Regulatory)	Association	Ethambutol, Isoniazid, Pyrazinamide and Rifampicin	Kim et al. (2012)
<i>NR112</i>	rs2472677	<i>NR112</i> _63396C>T (Intron)	Association	Atazanavir, TB and HIV drugs, and Isoniazid	Calcagno et al. (2019); Schipani et al. (2010); Siccardi et al. (2008)
<i>CUX2</i>	rs7958375	<i>CUX2</i> _c.64-11987G>A (Intron)	Association (GWAS)	Rifampicin	Petros et al. (2016)
<i>AGBL4</i>	rs319952	<i>AGBL4</i> _c.206+5387T>C (Intron)			
	rs320003	<i>AGBL4</i> _c.91+2046C>T (Intron)			

Note: All SNPs are reported in PharmGKB except rs4647992.

GWAS, genome wide association study.

<sup>a</sup> Pharmacogenetic labelling for TB/HIV/Malaria drugs approved by US Food and Drug Administration (FDA)/European Medicines Agency (EMA).<sup>b</sup> LD SNPs used for genotyping (rs12768009 - rs4986894, rs7439366 - rs7438135, rs3064744 - rs887829).

## Data analysis

Five SNPs failed in the assessment of Hardy–Weinberg Equilibrium (HWE). The allele frequencies of the remaining 27 SNPs were compared with those of non-African and African populations derived from data of the 1000 genome phase 3 project ([www.ensembl.org](http://www.ensembl.org)). This database includes data of non-African populations, namely of mixed American (AMR), East Asian (EAS), European (EUR) and South Asian (SAS) groups and of African (AFR) populations. The latter include Yoruba in Ibadan, Nigeria (YRI), Luhya in Webuye, Kenya (LWK), Gambian in the West Coast Region, The Gambia (GWD), Mende in Sierra Leone (MSL), Esan in Nigeria (ESN), Americans of African descent in the South-West of the USA (ASW) and people of African-Caribbean descent from Barbados

(ACB). Pairwise comparisons of genotypes among the populations were done using Fisher's exact test, taking into account a p-value of <0.05.

## Results

Using a custom-designed panel of SNPs we genotyped in 67 samples 32 SNPs localized in 12 ADME and 6 non-ADME genes which are known to have pharmacokinetic and pharmacodynamic relevance for drugs used to treat TB, HIV/AIDS and malaria. Details of all SNPs, genes, drug substrates and their pharmacogenetic relevance are given in Table 1. The allele frequencies of the 32 variants are shown in Table 2. Five out of the 32 SNPs failed in the assessment of HWE (*CYP2B6*\*6, *CYP2D6*\*10, rs1050828 and

rs1050829 [G6PD], rs4647992 [NF-κB1]) and were excluded from further statistical analysis. Failure of SNPs passing HWE could be due to the small sample size or errors in genotyping.

Allele frequencies in the ROC and non-African populations

Non-African populations included mixed American (AMR), East Asian (EAS), European (EUR) and South Asian (SAS) groups.

A pairwise comparison of the allele frequencies observed in ROC with these four non-African populations revealed significant differences in allele frequencies for the majority of SNPs tested (Table 2; p-values in Supplementary Table 2). 18 polymorphisms in the AMR group and 21 polymorphisms in the EAS, EUR and SAS populations showed significant differences. In the comparison of ROC with all non-African populations, the frequencies of 10 polymorphisms were significantly different, namely those of ABCB1\*6, CYP2B6\*18, CYP2C8\*2, CYP3A5\*3, CYP3A5\*6, CYP3A5\*7, CYP3A4\*1B, SLC01B1\*5, UGT1A1\_1813C>T and rs7958375-CUX2. Significant differences of allele frequencies applied also to 16 polymorphisms in some of the non-African populations only, namely rs3842-ABCB1 (AMR, EUR), CYP2A6\*9 (EAS, EUR), rs12768009-CYP2C19 (AMR, EAS, SAS), CYP2C19\*2 (AMR, EAS, SAS), NAT2\*5 (EAS), NAT2\*6 (EAS, EUR, SAS), NAT2\*11 (EAS, EUR), NAT2\*12 (EAS, SAS), NAT2\_18415371G>A (EAS, EUR, SAS), SLC01B1\*1b (AMR, EUR, SAS), UGT2B7\*2 (EUR, SAS), UGT1A1\*28 (AMR, EAS, EUR), rs1800629-TNF-α (AMR, EAS, SAS), rs2472677-NR1I2 (EAS, EUR, SAS), rs319952-AGBL4 (AMR, EUR, SAS) and rs320003-AGBL4 (AMR, EUR, SAS). This is in concordance with published literature (Rajman et al., 2017; Xing et al., 2010), which

describes significant genetic differences between non-African and African populations.

Allele frequencies in the ROC and other African populations

Other African populations with which comparisons were made included Yoruba in Ibadan, Nigeria (YRI), Luhya in Webuye, Kenya (LWK), Gambians in the West Coast Region, The Gambia (GWD), Mende in Sierra Leone (MSL), Esan in Nigeria (ESN), Americans of African descent in the South-West of the USA (ASW) and people of African-Caribbean descent from Barbados (ACB).

Pairwise comparisons of the allele frequencies of the 27 SNPs in the ROC population with the other African populations were made. Significant differences were found for only a few of the SNPs (Table 2; p-values in Supplementary Table 2). In detail, 5 polymorphisms in the ACB group, 3 polymorphisms in ASW, 5 polymorphisms in ESN, 2 polymorphisms in GWD, 5 polymorphisms in LWK, 5 polymorphisms in MSL and 4 polymorphisms in the YRI population showed significant differences of allele frequencies. Polymorphisms exhibiting significant differences in comparisons of the ROC with other African populations are rs3842-ABCB1 (ACB, GWD), CYP2A6\*9 (ACB, GWD), CYP2B6\*18 (ACB, ESN, LWK), CYP3A5\*3 (ASW), CYP3A5\*6 (LWK), NAT2\*5 (MSL, YRI), NAT2\*6 (ACB, ASW, ESN, LWK), NAT2\*11 (ESN, MSL, YRI) NAT2\*13 (ACB, ESN, MSL, YRI), SLC01B1\*5 (ESN), UGT1A1\_1813C>T (LWK, MSL), rs1800629-TNF-α (ACB, GWD), and rs2472677-NR1I2 (MSL, YRI). As expected, the number of SNPs with frequency differences in the ROC compared to other African populations was very small. Despite the low degree of differences, there are enormous

Table 2 Allele frequencies of 32 SNPs in Republic of Congo genotyped and allele frequencies in non-African and African populations from the 1000 Genome Project phase 3 data.

Gene	SNP ID	Risk Allele	Non-African populations				African populations							ROC (Genotyped)	HWE (P-values)	
			AMR	EAS	EUR	SAS	AFR	ACB	ASW	ESN	GWD	LWK	MSL			YRI
ABCB1	rs1045642	T	<b>43</b>	<b>40</b>	<b>52</b>	<b>58</b>	12–19	15	19	12	19	14	15	13	17	0.4
	rs3842	G	<b>16</b>	<b>30</b>	<b>14</b>	<b>18</b>	4–25	<b>4</b>	17	17	<b>12</b>	25	16	15	24	0.06
CYP2A6	rs28399433	G	<b>10</b>	<b>24</b>	<b>7</b>	15	6–11	<b>6</b>	11	10	<b>6</b>	9	7	10	14	0.51
CYP2B6	rs28399499	C	<b>1</b>	<b>0</b>	<b>0</b>	<b>0</b>	6–12	<b>6</b>	10	<b>7</b>	<b>10</b>	<b>6</b>	8	12	16	0.74
	rs3745274	T	<b>37</b>	<b>22</b>	<b>24</b>	<b>38</b>	35–41	<b>38</b>	35	41	35	<b>36</b>	35	40	36	0.02 <sup>a</sup>
CYP2D6	rs1065852	T	15	57	20	17	4–16	15	16	9	12	4	17	11	8	0.01 <sup>a</sup>
CYP2C19	rs12768009	A	<b>10</b>	<b>37</b>	15	<b>37</b>	14–23	16	14	22	14	23	19	17	19	0.79
	rs4244285	A	<b>11</b>	<b>31</b>	15	<b>36</b>	13–21	15	14	21	13	21	18	17	19	0.79
CYP2C8	rs11572103	A	<b>1</b>	<b>0</b>	<b>0</b>	<b>1</b>	14–24	21	15	20	24	14	17	20	21	0.13
CYP3A5	rs776746	G	<b>20</b>	<b>29</b>	<b>6</b>	<b>33</b>	69–89	75	<b>69</b>	89	77	88	88	83	84	0.86
	rs10264272	A	<b>2</b>	<b>0</b>	<b>0</b>	<b>0</b>	5–24	12	5	14	16	<b>24</b>	16	17	11	0.15
	rs41303343	AA	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	9–14	11	12	9	14	<b>12</b>	14	12	12	0.27
CYP3A4	rs2740574	G	<b>10</b>	<b>0</b>	<b>3</b>	<b>4</b>	66–83	66	67	77	79	83	83	76	75	0.96
NAT2	rs1801280	C	36	<b>4</b>	45	35	24–36	28	31	27	34	36	<b>24</b>	<b>24</b>	36	0.75
	rs1799930	A	17	<b>26</b>	<b>28</b>	<b>36</b>	19–29	<b>26</b>	<b>29</b>	<b>26</b>	19	<b>27</b>	<b>22</b>	<b>20</b>	16	0.47
	rs1799929	T	34	<b>4</b>	<b>44</b>	<b>32</b>	17–33	<b>25</b>	<b>26</b>	<b>19</b>	28	<b>33</b>	<b>19</b>	<b>17</b>	29	0.69
	rs1208	G	37	<b>4</b>	44	<b>36</b>	35–46	38	36	<b>39</b>	44	46	<b>35</b>	<b>37</b>	46	0.51
	rs1041983	T	29	44	31	43	39–53	<b>50</b>	45	<b>53</b>	39	43	<b>50</b>	<b>50</b>	36	0.75
	rs1495741	A	65	<b>48</b>	<b>76</b>	<b>78</b>	51–67	59	67	<b>56</b>	57	66	54	51	63	0.39
	rs2306283	A	<b>53</b>	<b>24</b>	<b>60</b>	<b>45</b>	12–25	21	25	12	19	16	19	19	17	0.45
SLCO1B1	rs4149032	C	<b>61</b>	<b>40</b>	<b>66</b>	<b>49</b>	19–36	30	36	<b>19</b>	23	28	27	28	24	0.82
UGT2B7	rs7439366	T	<b>32</b>	<b>28</b>	<b>49</b>	<b>40</b>	18–28	28	25	<b>24</b>	19	25	18	21	25	0.4
UGT1A1	rs3064744	(TA) <sub>8</sub>	<b>36</b>	<b>13</b>	<b>29</b>	<b>40</b>	38–46	43	42	40	38	46	41	46	47	0.44
	rs10929303	T	<b>22</b>	<b>13</b>	<b>23</b>	<b>17</b>	32–50	46	39	50	47	<b>32</b>	<b>33</b>	45	46	0.3
G6PD	rs1050828	A	1	0	0	0	4–21	13	17	16	4	18	7	21	19	<.0001 <sup>a</sup>
	rs1050829	G	3	0	0	0	28–38	33	29	35	36	34	28	38	40	<.0001 <sup>a</sup>
NF-κB1	rs4647992	T	3	4	5	4	8–24	10	9	10	8	24	8	10	9	0.03 <sup>a</sup>
TNF-α	rs1800629	A	7	6	13	5	7–14	14	7	13	14	9	16	10	18	0.34
NR1I2	rs2472677	T	48	<b>62</b>	<b>66</b>	<b>57</b>	28–42	42	35	36	38	40	<b>28</b>	<b>36</b>	48	1
CUX2	rs7958375	G	<b>98</b>	<b>100</b>	<b>100</b>	<b>99</b>	76–86	83	86	85	84	79	76	81	80	0.19
AGBL4	rs319952	T	<b>51</b>	67	<b>40</b>	<b>46</b>	63–80	70	63	71	80	67	78	74	73	0.47
	rs320003	C	<b>51</b>	67	<b>40</b>	<b>46</b>	64–80	69	64	75	80	74	80	74	73	0.6

Bold and underlined: statistically significant (P < 0.05, Fisher exact test) pairwise comparison with congo population (P-values in Supplementary Table 2). African (AFR), Yoruba in Ibadan, Nigeria (YRI), Luhya in Webuye, Kenya (LWK), Gambians in the West Coast Region, The Gambia (GWD), Mende in Sierra Leone (MSL), Esan in Nigeria (ESN), Americans of African Ancestry in SW USA (ASW), African Caribbeans in Barbados (ACB), Mixed American (AMR), East Asian (EAS), European (EUR), South Asian (SAS) and Republic of Congo (ROC).

<sup>a</sup> Failed Hardy–Weinberg equilibrium (HWE).

differences in allele frequencies between all African populations as can be seen in the AFR column of Table 2, which shows the lowest and highest allele frequencies for all genetic variants. For example, the allele frequency of rs3842 SNP ranges from 4% to 25%. For some SNPs the allele frequencies among African populations do not differ significantly however, due to the high burden of disease this may lead to considerable differences in treatment efficacy and drug toxicity.

## Discussion

In this study, we performed Taqman genotyping assays of SNPs located in 18 genes in 67 samples of febrile children recruited in ROC (Etoka-Beka et al., 2016; Gampio Gueye et al., 2019).

### CYP variations

Cytochromes P450 (CYPs) are a well characterized family of phase 1 drug metabolising enzymes involved in the metabolism of 70–80% of drugs. Polymorphisms in these genes are associated with altering the pharmacokinetics of several drugs and leading to ADRs (Phillips et al., 2001) or suboptimal clinical outcome. *CYP2B6\*6* and *CYP2B6\*18* alleles exhibit reduced enzymatic activity and increased plasma concentrations of efavirenz and nevirapine, which may lead to ADRs (Ciccacci et al., 2013; Cusato et al., 2016; Gandhi et al., 2012). Due to its clinical significance, *CYP2B6\*6* is approved for efavirenz drug labelling by FDA/EMA/Swissmedic/PMDA (Whirl-Carrillo et al., 2012). Studies have shown that in patients with the *CYP2B6\*6* homozygous allele, reduction in the daily dose of efavirenz from 600 to 200 mg/day was sufficient to maintain therapeutic levels and reduce ADRs (Dhoro et al., 2015). The frequency of the *CYP2B6\*6* allele in the ROC population is 36%. In this study, this particular allele *CYP2B6\*6* failed to pass HWE assessment. A previous study showed an allele frequency of 55% (Peko et al., 2019a), which is the highest compared to other African and to non-African populations.

*CYP2C8\*2* reduces the enzyme activity and alters the bioavailability of the antimalarial drug amodiaquine, leading to ADRs and possibly the emergence of drug resistance (Parikh et al., 2007). Amodiaquine is known to be the drug most commonly associated with ADRs in Africa (Cavaco et al., 2005; Kudzi et al., 2009). The allele frequency of this variant in ROC is 21%, and in other African populations the frequency ranges from 14 to 24%, while it is virtually non-existent in non-African populations. Since this variant occurs almost exclusively in African populations, respective genotyping in cases of ADRs is recommended for ROC as well as African populations treated with amodiaquine. *CYP3A5\*3*, *-\*6* and *-\*7* reduce enzyme function and increase plasma concentrations of quinine, lumefantrine, nevirapine and atazanavir (Brown et al., 2012; Castillo-Mancilla et al., 2016; Mutagonda et al., 2017).

The *CYP2A6\*9* variant reduces enzyme activity and increases plasma levels of efavirenz (Soeria-Atmadja et al., 2017). This observation failed to be replicated in some studies. The allele frequency of this variant in ROC is 14%; the frequencies show significant differences in other African (6–11%) and non-African populations (7–24%). The allele frequency in the ROC is highest among all African populations, suggesting a higher risk of efavirenz-induced toxicity, and further studies are needed to strengthen the associations. The *CYP2C19\*2* and *CYP2C19\_G>A* variants, which are in linkage disequilibrium, decrease enzyme function and increase plasma concentrations of nevirapine (Lehr et al., 2011). Only a few studies have been conducted so far and further studies are recommended. The allele frequency in ROC is 19% and significant differences among African and non-African

populations were observed. *CYP3A4\*1B* reduces the enzyme function and increases the plasma concentrations of lumefantrine, nevirapine atazanavir, and efavirenz (Bertrand et al., 2009; Haas et al., 2005; Kile et al., 2012; Mutagonda et al., 2017). The frequency of the *CYP3A4\*1B* allele in ROC is 75% and also high allele frequencies are seen in other African populations (66–83%) compared to non-African populations (0–10%). Since African populations have the highest risk of increasing plasma concentrations of drugs metabolized by *CYP3A4* and subsequent toxicity; genotyping and dose adjustment could reduce toxicity and improve compliance.

### NAT and UGT variations

N-acetyltransferase (NAT) and UDP-glucuronosyltransferase (UGT) are phase II metabolizing enzymes which play a key role in detoxification of drugs and xenobiotics. Isoniazid is metabolized by NAT2, and the slow acetylator variants are associated with anti-TB drug-induced hepatotoxicity (Ben Mahmoud et al., 2012; Chan et al., 2017; Shi et al., 2015; Singla et al., 2014). Due to its clinical importance it is approved for pharmacogenetic labelling by FDA. The NAT2 variants assessed in this study include the slow acetylation variants *NAT2\*5* and *-\*6*, the fast acetylation variants *NAT2\*12* and *-\*13*, and drug-induced liver injury associated variants *NAT2\_tag* SNP and *NAT2\*11*. The allele frequency of the slow acetylator *NAT2\*5* and *-\*6* variants in ROC is 36% and 17%, respectively, and their frequencies vary considerably across populations. *UGT2B7\*2*, *UGT1A1\*28* and *-\*76* cause variable drug concentrations of efavirenz, atazanavir and ritanovir and toxicity (Johnson et al., 2014; Nishijima et al., 2014; Vardhanabhuti et al., 2015). Due to their clinical significance, FDA labelling has been approved for atazanavir (Du et al., 2019). The allele frequencies of *UGT2B7\*2*, *UGT1A1\*28* and *UGT1A1\_1813C>T* variants in ROC are 25%, 47% and 46% respectively. The frequencies vary significantly in other African populations, ranging from 18 to 28%, 38 to 46% and 32 to 50%, respectively and also differ significantly in non-African populations. The *UGT2B7\*2* allele frequency is generally lower in African than in non-African populations. The *UGT1A1\*28* and *UGT1A1\_1813C>T* allele frequencies are higher in African than in non-African populations.

### ABCB1 and SLCO1B1 variations

The P-glycoprotein (P-gp), encoded by *ABCB1* and the solute carrier organic anion transporter family member 1B1 protein, encoded by the *SLCO1B1* gene, are transporter proteins which facilitate transfer of drugs across cell membranes. Variants in these genes lead to toxicity or inefficacy and drug resistance.

The *ABCB1\*6*, *ABCB1\_c193A>G*, *SLCO1B1\*1b* and *SLCO1B1\*5* variants are associated with altered efficacy, pharmacokinetics and toxicity of rifampicin, isoniazid, efavirenz, nevirapine, atazanavir and ritonavir (Calcagno et al., 2019; Ciccacci et al., 2010; Elens et al., 2010; Kwara et al., 2014; Park et al., 2010; Yimer et al., 2011). The allele frequency of *ABCB1\*6* and *c193A>G* in ROC is 17% and 24% respectively, and the frequencies differ significantly between all populations. The most extensively studied variant *ABCB1\*6* is associated with altered efficacy and toxicity in the treatment with anti-HIV and anti-TB drugs.

The frequencies of *SLCO1B1\*1b* and *-\*5* in ROC are 17% and 24%, respectively. These variants occur significantly more frequently in non-African populations compared to African populations, indicating the relative risk for elevation of rifampicin and efavirenz plasma concentrations and isoniazid drug-induced toxicity. The *SLCO1B1* polymorphisms are common in South Africa and often associated with increased rifampicin clearance, leading to the recommendation of dose adaptation (Chigutsa et al., 2011).

### Non-ADME gene variations

The non-absorption, distribution, metabolism and excretion (non-ADME) variants of *G6PD*, *NF-κB1*, *TNF-α*, *NR112*, *CUX2* and *AGBL4* are associated with toxicity and the pharmacokinetics of anti-HIV, TB and malaria drugs (Calcagno et al., 2019; Fanello et al., 2008; Kim et al., 2012; Petros et al., 2016; Schipani et al., 2010; Siccardi et al., 2008).

TNF-α is an inflammatory cytokine, which plays an important role in drug-induced immune reactions. NR112 (PXR) is a transcription factor that activates transcription of several ADME genes. *CUX2* and *AGBL4* gene variants have been found to be associated with anti-TB drug-induced hepatotoxicity; however, the causal mechanisms have not been identified so far. The variants rs1800629 (*TNF-α*), rs4647992 (*NF-κB1*), rs2472677 (*NR112*), rs7958375 (*CUX2*), rs319952 and rs320003 (*AGBL4*) are associated with anti-TB treatment-induced ADRs.

Study limitations include the small sample size, which needed to be kept small because of cost considerations. Also, variants with less than 10% allele frequency were not included despite their clinical significance. This high cutoff was kept, keeping in mind that most of the studies have low sample sizes and identifying statistical differences becomes difficult.

### Conclusions

We have designed a custom SNP panel and successfully genotyped distinct pharmacogenetic variants in ROC. When comparing the allele-frequencies of African with non-African populations, we found variations in allele frequencies among all populations. This study helps in predicting efficacy and ADRs at the population and individual level and underlines the importance of understanding the role and function of pharmacogenetic variants.

### Ethics approval and consent to participate

Ethical approval was obtained from the Institutional Ethics Committee of the Fondation Congolaise pour la Recherche Médicale in Brazzaville, ROC. Following written informed consent from parents/guardians, a total of 67 children aged between one and ten years with fever were included.

### Consent for publication

All authors agreed with the results and conclusions. All authors consented to this version of the manuscript to be published.

### Availability of data and materials

All related data supporting the results reported in the article is available within the manuscript and in supplementary data file.

### Conflict of interest

None declare.

### Funding

This work has been supported under the grant agreement CANTAM2 (EDCTP/RegNet2015-1045).

### Author contributions

TPV designed and supervised the study, and wrote the manuscript. AAA, VBP, EAA, GM, MB, MPG, FN and PGK were involved in the study design. FN and PGK contributed to the study

materials. SRP designed the SNP panel. SRP, DEA analyzed and wrote the first draft of the manuscript. CGM edited the manuscript. SRP, LTKL, CFN and KAF performed the experiments.

### Acknowledgements

The authors acknowledge the support of field workers and consented participants of the study.

### Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.ijid.2020.12.009>.

### References

- Alomar MJ. Factors affecting the development of adverse drug reactions (Review article). *Saudi Pharm J* 2014;22(2):83–94.
- Bains RK. African variation at Cytochrome P450 genes: evolutionary aspects and the implications for the treatment of infectious diseases. *Evol Med Public Health* 2013;2013(1):118–34.
- Bassi PU, Osakwe AI, Ogar CK, Elagbaje C, Nwankwo BB, Balogun ST, et al. Impact of comorbidity on adverse drug reaction profile in a cohort of patients treated with Artemisinin combination therapies for uncomplicated malaria in Nigeria. *Pharmacol Res Perspect* 2017;5(2):e00302.
- Ben Mahmoud L, Ghozzi H, Kamoun A, Hakim A, Hachicha H, Hammami S, et al. Polymorphism of the N-acetyltransferase 2 gene as a susceptibility risk factor for antituberculosis drug-induced hepatotoxicity in Tunisian patients with tuberculosis. *Pathol Biol (Paris)* 2012;60(5):324–30.
- Bennett JW, Pybus BS, Yadava A, Tosh D, Sousa JC, McCarthy WF, et al. Primaquine failure and cytochrome P-450 2D6 in Plasmodium vivax malaria. *N Engl J Med* 2013;369(14):1381–2.
- Bertrand J, Treluyer JM, Panhard X, Tran A, Auleley S, Rey E, et al. Influence of pharmacogenetics on indinavir disposition and short-term response in HIV patients initiating HAART. *Eur J Clin Pharmacol* 2009;65(7):667–78.
- Brown KC, Hosseini-pour MC, Hoskins JM, Thirumaran RK, Tien HC, Weigel R, et al. Exploration of CYP450 and drug transporter genotypes and correlations with nevirapine exposure in Malawians. *Pharmacogenomics* 2012;13(1):113–21.
- Calcagno A, Cusato J, Sekaggya-Wiltshire C, von Braun A, Motta I, Turyasingura G, et al. The influence of pharmacogenetic variants in HIV/Tuberculosis coinfecting patients in Uganda in the SOUTH study. *Clin Pharmacol Ther* 2019;106(2):450–7.
- Castillo-Mancilla JR, Aquilante CL, Wempe MF, Smeaton LM, Firnhaber C, LaRosa AM, et al. Pharmacogenetics of unboosted atazanavir in HIV-infected individuals in resource-limited settings: a sub-study of the AIDS Clinical Trials Group (ACTG) PEARLS study (NWCS 342). *J Antimicrob Chemother* 2016;71(6):1609–18.
- Cavaco I, Stromberg-Norklit J, Kaneko A, Msellem MI, Dahoma M, Ribeiro VL, et al. CYP2C8 polymorphism frequencies among malaria patients in Zanzibar. *Eur J Clin Pharmacol* 2005;61(1):15–8.
- Chamorro JG, Castagnino JP, Musella RM, Noguera M, Frias A, Visca M, et al. tagSNP rs1495741 as a useful molecular marker to predict antituberculosis drug-induced hepatotoxicity. *Pharmacogenet Genomics* 2016;26(7):357–61.
- Chan SL, Chua APG, Aminkeng F, Chee CBE, Jin S, Loh M, et al. Association and clinical utility of NAT2 in the prediction of isoniazid-induced liver injury in Singaporean patients. *PLoS One* 2017;12(10):e0186200.
- Chaudhry M, Alessandrini M, Pepper MS. Pharmacogenomics for infectious diseases in sub-Saharan Africa: successes and opportunities. *Appl Transl Genom* 2016;9:3–5.
- Chigutsa E, Visser ME, Swart EC, Denti P, Pushpakom S, Egan D, et al. The SLCO1B1 rs4149032 polymorphism is highly prevalent in South Africans and is associated with reduced rifampin concentrations: dosing implications. *Antimicrob Agents Chemother* 2011;55(9):4122–7.
- Ciccacci C, Borgiani P, Ceffa S, Sirianni E, Marazzi MC, Altan AM, et al. Nevirapine-induced hepatotoxicity and pharmacogenetics: a retrospective study in a population from Mozambique. *Pharmacogenomics* 2010;11(1):23–31.
- Ciccacci C, Di Fusco D, Marazzi MC, Zimba I, Erba F, Novelli G, et al. Association between CYP2B6 polymorphisms and Nevirapine-induced SJS/TEN: a pharmacogenetics study. *Eur J Clin Pharmacol* 2013;69(11):1909–16.
- Cusato J, Tomasello C, Simiele M, Calcagno A, Bonora S, Marinaro L, et al. Efavirenz pharmacogenetics in a cohort of Italian patients. *Int J Antimicrob Agents* 2016;47(2):117–23.
- Dhoro M, Zvada S, Ngara B, Nhachi C, Kadzirange G, Chonzi P, et al. CYP2B6\*6, CYP2B6\*18, Body weight and sex are predictors of efavirenz pharmacokinetics and treatment response: population pharmacokinetic modeling in an HIV/AIDS and TB cohort in Zimbabwe. *BMC Pharmacol Toxicol* 2015;16:4.
- Du P, Wang A, Ma Y, Li X. Association between the UGT1A1\*28 allele and hyperbilirubinemia in HIV-positive patients receiving atazanavir: a meta-analysis. *Biosci Rep* 2019;39(5).
- Elsens L, Vandercam B, Yombi JC, Lison D, Wallemacq P, Haufroid V. Influence of host genetic factors on efavirenz plasma and intracellular pharmacokinetics in HIV-1-infected patients. *Pharmacogenomics* 2010;11(9):1223–34.

- Etoka-Beka MK, Ntoumi F, Kombo M, Deibert J, Poulain P, Vouvongui C, et al. Plasmodium falciparum infection in febrile Congolese children: prevalence of clinical malaria 10 years after introduction of artemisinin-combination therapies. *Trop Med Int Health* 2016;21(12):1496–503.
- Fanello CI, Karema C, Avellino P, Bancone G, Uwimana A, Lee SJ, et al. High risk of severe anaemia after chlorproguanil-dapsone+artesunate antimalarial treatment in patients with G6PD (A-) deficiency. *PLoS One* 2008;3(12):e4031.
- Gampio Gueye NS, Peko SM, Nderu D, Koukouikila-Koussounda F, Vouvongui C, Kobawila SC, et al. An update on glucose-6-phosphate dehydrogenase deficiency in children from Brazzaville, Republic of Congo. *Malar J* 2019;18(1):57.
- Gandhi M, Greenblatt RM, Bacchetti P, Jin C, Huang Y, Anastos K, et al. A single-nucleotide polymorphism in CYP2B6 leads to &3-fold increases in efavirenz concentrations in plasma and hair among HIV-infected women. *J Infect Dis* 2012;206(9):1453–61.
- Gurdasani D, Carstensen T, Tekola-Ayele F, Pagani L, Tachmazidou I, Hatzikotoulas K, et al. The African genome variation project shapes medical genetics in Africa. *Nature* 2015;517(7534):327–32.
- Haas DW, Smeaton LM, Shafer RW, Robbins GK, Morse GD, Labbe L, et al. Pharmacogenetics of long-term responses to antiretroviral regimens containing Efavirenz and/or Nelfinavir: an Adult Aids Clinical Trials Group Study. *J Infect Dis* 2005;192(11):1931–42.
- Ho HT, Wang TH, Hsiung CH, Perng WC, Wang NC, Huang TY, et al. The NAT2 tag SNP rs1495741 correlates with the susceptibility of antituberculosis drug-induced hepatotoxicity. *Pharmacogenet Genomics* 2013;23(4):200–7.
- Johnson DH, Venuto C, Ritchie MD, Morse GD, Daar ES, McLaren PJ, et al. Genomewide association study of atazanavir pharmacokinetics and hyperbilirubinemia in AIDS Clinical Trials Group protocol A5202. *Pharmacogenet Genomics* 2014;24(4):195–203.
- Kattel K, Evande R, Tan C, Mondal G, Grem JL, Mahato RI. Impact of CYP2C19 polymorphism on the pharmacokinetics of nelfinavir in patients with pancreatic cancer. *Br J Clin Pharmacol* 2015;80(2):267–75.
- Kile DA, MaWhinney S, Aquilante CL, Rower JE, Castillo-Mancilla JR, Anderson PL. A population pharmacokinetic-pharmacogenetic analysis of atazanavir. *AIDS Res Hum Retroviruses* 2012;28(10):1227–34.
- Kim SH, Kim SH, Yoon HJ, Shin DH, Park SS, Kim YS, et al. TNF-alpha genetic polymorphism -308G/A and antituberculosis drug-induced hepatitis. *Liver Int* 2012;32(5):809–14.
- Kudzi W, Dodoo AN, Mills JJ. Characterisation of CYP2C8, CYP2C9 and CYP2C19 polymorphisms in a Ghanaian population. *BMC Med Genet* 2009;10:124.
- Kwara A, Lartey M, Sagoe KW, Kenu E, Court MH. CYP2B6, CYP2A6 and UGT2B7 genetic polymorphisms are predictors of efavirenz mid-dose concentration in HIV-infected patients. *AIDS* 2009;23(16):2101–6.
- Kwara A, Cao L, Yang H, Poethke P, Kurpewski J, Tashima KT, et al. Factors associated with variability in rifampin plasma pharmacokinetics and the relationship between rifampin concentrations and induction of efavirenz clearance. *Pharmacotherapy* 2014;34(3):265–71.
- Lehr T, Yuan J, Hall D, Zimdahl-Gelling H, Schaefer HG, Staab A, et al. Integration of absorption, distribution, metabolism, and elimination genotyping data into a population pharmacokinetic analysis of nevirapine. *Pharmacogenet Genomics* 2011;21(11):721–30.
- Li J, Zhang L, Zhou H, Stoneking M, Tang K. Global patterns of genetic diversity and signals of natural selection for human ADME genes. *Hum Mol Genet* 2011;20(3):528–40.
- Linguissi LS, Gwom LC, Nkenfou CN, Bates M, Petersen E, Zumla A, et al. Health systems in the Republic of Congo: challenges and opportunities for implementing tuberculosis and HIV collaborative service, research, and training activities. *Int J Infect Dis* 2017;56:62–7.
- Ma MK, Woo MH, McLeod HL. Genetic basis of drug metabolism. *Am J Health Syst Pharm* 2002;59(21):2061–9.
- Mutagonda RF, Kamuhabwa AAR, Minzi OMS, Massawe SN, Asghar M, Homann MV, et al. Effect of pharmacogenetics on plasma lumefantrine pharmacokinetics and malaria treatment outcome in pregnant women. *Malar J* 2017;16(1):267.
- Nishijima T, Tsuchiya K, Tanaka N, Joya A, Hamada Y, Mizushima D, et al. Single-nucleotide polymorphisms in the UDP-glucuronosyltransferase 1A-3' untranslated region are associated with atazanavir-induced nephrolithiasis in patients with HIV-1 infection: a pharmacogenetic study. *J Antimicrob Chemother* 2014;69(12):3320–8.
- Parikh S, Ouedraogo JB, Goldstein JA, Rosenthal PJ, Kroetz DL. Amodiaquine metabolism is impaired by common polymorphisms in CYP2C8: implications for malaria treatment in Africa. *Clin Pharmacol Ther* 2007;82(2):197–203.
- Park WB, Choe PG, Song KH, Jeon JH, Park SW, Kim HB, et al. Genetic factors influencing severe atazanavir-associated hyperbilirubinemia in a population with low UDP-glucuronosyltransferase 1A1\*28 allele frequency. *Clin Infect Dis* 2010;51(1):101–6.
- Peko SM, Gueye NSG, Vouvongui C, Koukouikila-Koussounda F, Kobawila SC, Nderu D, et al. Cytochrome P450 CYP2B6\*6 distribution among Congolese individuals with HIV, Tuberculosis and Malaria infection. *Int J Infect Dis* 2019a;82:111–6.
- Peko SM, Ntoumi F, Vouvongui C, Nderu D, Kobawila SC, Velavan TP, et al. Distribution of the cytochrome P450 CYP2C8\*2 allele in Brazzaville, Republic of Congo. *Int J Infect Dis* 2019b;85:49–53.
- Petros Z, Lee MM, Takahashi A, Zhang Y, Yimer G, Habtewold A, et al. Genome-wide association and replication study of anti-tuberculosis drugs-induced liver toxicity. *BMC Genomics* 2016;17(1):755.
- Phillips KA, Veenstra DL, Oren E, Lee JK, Sadee W. Potential role of pharmacogenomics in reducing adverse drug reactions: a systematic review. *JAMA* 2001;286(18):2270–9.
- Rajman I, Knapp L, Morgan T, Masimirembwa C. African genetic diversity: implications for cytochrome P450-mediated drug metabolism and drug development. *EBioMedicine* 2017;17:67–74.
- Ramos E, Doumatey A, Elkhoulou AG, Shriner D, Huang H, Chen G, et al. Pharmacogenomics, ancestry and clinical decision making for global populations. *Pharmacogenomics J* 2014;14(3):217–22.
- Rattan A, Kalia A, Ahmad N. Multidrug-resistant Mycobacterium tuberculosis: molecular perspectives. *Emerg Infect Dis* 1998;4(2):195–209.
- Ross S, Anand SS, Joseph P, Pare G. Promises and challenges of pharmacogenetics: an overview of study design, methodological and statistical issues. *JRSM Cardiovasc Dis* 2012;1(1).
- Saito T, Gutierrez Rico EM, Kikuchi A, Kaneko A, Kumondai M, Akai F, et al. Functional characterization of 50 CYP2D6 allelic variants by assessing primaquine 5-hydroxylation. *Drug Metab Pharmacokinet* 2018;33(6):250–7.
- Schipani A, Siccardi M, D'Avolio A, Baietto L, Simiele M, Bonora S, et al. Population pharmacokinetic modeling of the association between 63396C-&T pregnane X receptor polymorphism and unboosted atazanavir clearance. *Antimicrob Agents Chemother* 2010;54(12):5242–50.
- Shahi GS, Mochhala SM, Das NP, Sunamoto J, Tajima K. The neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) induces changes in the heme spin state of microsomal cytochrome P-450. *Biochem Int* 1990;22(5):895–901.
- Shi J, Xie M, Wang J, Xu Y, Liu X. Susceptibility of N-acetyltransferase 2 slow acetylators to antituberculosis drug-induced liver injury: a meta-analysis. *Pharmacogenomics* 2015;16(18):2083–97.
- Sicard D, Kaplan JC, Labie D. Haemoglobinopathies and G.-6-P.D. deficiency in Laos. *Lancet* 1978;2(8089):571–2.
- Siccardi M, D'Avolio A, Baietto L, Gibbons S, Sciandra M, Colucci D, et al. Association of a single-nucleotide polymorphism in the pregnane X receptor (PXR 63396C-&T) with reduced concentrations of unboosted atazanavir. *Clin Infect Dis* 2008;47(9):1222–5.
- Singla N, Gupta D, Birbian N, Singh J. Association of NAT2, GST and CYP2E1 polymorphisms and anti-tuberculosis drug-induced hepatotoxicity. *Tuberculosis (Edinb)* 2014;94(3):293–8.
- Sinxadi PZ, Leger PD, McIlerron HM, Smith PJ, Dave JA, Levitt NS, et al. Pharmacogenetics of plasma efavirenz exposure in HIV-infected adults and children in South Africa. *Br J Clin Pharmacol* 2015;80(1):146–56.
- Soeria-Atmadja S, Osterberg E, Gustafsson LL, Dahl ML, Eriksen J, Rubin J, et al. Genetic variants in CYP2B6 and CYP2A6 explain interindividual variation in efavirenz plasma concentrations of HIV-infected children with diverse ethnic origin. *PLoS One* 2017;12(9):e0181316.
- St Jean PL, Xue Z, Carter N, Koh GC, Duparc S, Taylor M, et al. Tafenoquine treatment of Plasmodium vivax malaria: suggestive evidence that CYP2D6 reduced metabolism is not associated with relapse in the Phase 2b DETECTIVE trial. *Malar J* 2016;15:97.
- Tishkoff SA, Reed FA, Friedlaender FR, Ehret C, Ranciaro A, Froment A, et al. The genetic structure and history of Africans and African Americans. *Science* 2009;324(5930):1035–44.
- Turner D, Wainberg MA. HIV transmission and primary drug resistance. *AIDS Rev* 2006;8(1):17–23.
- Vardhanabuti S, Ribaldo HJ, Landovitz RJ, Ofotokun I, Lennox JL, Currier JS, et al. Screening for UGT1A1 genotype in study A5257 would have markedly reduced premature discontinuation of atazanavir for hyperbilirubinemia. *Open Forum Infect Dis* 2015;2(3):ofv085.
- Whirl-Carrillo M, McDonagh EM, Hebert JM, Gong L, Sangkuhl K, Thorn CF, et al. Pharmacogenomics knowledge for personalized medicine. *Clin Pharmacol Ther* 2012;92(4):414–7.
- WHO. Guidelines for the Treatment of Malaria – 3rd edition. 2015.
- WHO. World Malaria Report 2015. Geneva: World Health Organization; 2015.
- WHO. Update of Recommendations on First- and Second-Line Antiretroviral Regimens. Geneva, Switzerland: World Health Organization; 2019 2019 (WHO/CDS/HIV/19.15).
- Wilson JF, Weale ME, Smith AC, Gratrix F, Fletcher B, Thomas MG, et al. Population genetic structure of variable drug response. *Nat Genet* 2001;29(3):265–9.
- Xing J, Watkins WS, Shlien A, Walker E, Huff CD, Witherspoon DJ, et al. Toward a more uniform sampling of human genetic diversity: a survey of worldwide populations by high-density genotyping. *Genomics* 2010;96(4):199–210.
- Yimer G, Ueda N, Habtewold A, Amogne W, Suda A, Riedel KD, et al. Pharmacogenetic & pharmacokinetic biomarker for efavirenz based ARV and rifampicin based anti-TB drug induced liver injury in TB-HIV infected patients. *PLoS One* 2011;6(12):e27810.
- Zhang J, Zhao Z, Bai H, Wang M, Jiao L, Peng W, et al. Genetic polymorphisms in PXR and NF-kappaB1 influence susceptibility to anti-tuberculosis drug-induced liver injury. *PLoS One* 2019;14(9):e0222033.