

RESEARCH ARTICLE

Ecological and life-history correlates of erythrocyte size and shape in Lepidosauria

Zachary Penman | D. Charles Deeming  | Carl D. Soulsbury 

School of Life Sciences and Environmental Sciences, University of Lincoln, Lincoln, UK

Correspondence

Carl D. Soulsbury, School of Life Sciences and Environmental Sciences, University of Lincoln, Brayford Pool, Lincoln LN6 7TS, UK.

Email: csoulsbury@lincoln.ac.uk

Abstract

Blood oxygen-carrying capacity is shaped both by the ambient oxygen availability as well as species-specific oxygen demand. Erythrocytes are a critical part of oxygen transport and both their size and shape can change in relation to species-specific life-history, behavioural or ecological conditions. Here, we test whether components of the environment (altitude), life history (reproductive mode, body temperature) and behaviour (diving, foraging mode) drive erythrocyte size variation in the Lepidosauria (lizards, snakes and rhynchocephalians). We collected data on erythrocyte size (area) and shape (L/W: elongation ratio) from Lepidosauria across the globe ($N = 235$ species). Our analyses show the importance of oxygen requirements as a driver of erythrocyte size. Smaller erythrocytes were associated with the need for faster delivery (active foragers, high-altitude species, warmer body temperatures), whereas species with greater oxygen demands (diving species, viviparous species) had larger erythrocytes. Erythrocyte size shows considerable cross-species variation, with a range of factors linked to the oxygen delivery requirements being major drivers of these differences. A key future aspect for study would include within-individual plasticity and how changing states, for example, pregnancy, perhaps alter the size and shape of erythrocytes in Lepidosauria.

KEYWORDS

blood oxygen-carrying capacity, erythrocyte, life history, oviparity, viviparity

1 | INTRODUCTION

There is considerable variation in erythrocyte size between- and within-taxa across all vertebrates (Gregory, 2000; Hawkey et al., 1991; Lay & Baldwin, 1999). There are a number of key drivers of erythrocyte size such as a positive relationship between genome size and cell size across vertebrates (Gregory, 2000, 2001, 2002; Mueller et al., 2008). For vertebrates with nucleated erythrocytes, erythrocyte size is directly related to nucleus size (Gregory, 2002). There are also broad differences between ectotherms and

endotherms with reptiles, amphibians and fish all having larger cells than mammals and birds (Gregory, 2001; Hawkey et al., 1991). In addition, nucleated avian erythrocytes are larger than those of non-nucleated mammals of an equivalent body mass (Gregory, 2002). The smaller erythrocytes of endotherms (bird, mammals) in comparison to ectotherms (reptiles, fish, amphibia) are thought to derive from lower mass-specific resting metabolic rates (RMR) and lower oxygen demands of ectotherms (Hawkey et al., 1991), something that also contributes to within-taxon patterns (Gregory, 2002; Kozłowski et al., 2020). In contrast, body mass does not necessarily

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2022 The Authors. *Journal of Evolutionary Biology* published by John Wiley & Sons Ltd on behalf of European Society for Evolutionary Biology.

correlate with erythrocyte size. Positive interspecific relationships between erythrocyte size and body mass have been demonstrated in geckos (Starostová et al., 2005) and birds (Kostecka-Myrcha & Chołostiakow-Gromek, 2001; Soulsbury et al., 2022). In mammals and teleost fish, there is no relationship between erythrocyte size and body mass or size (mammals: Promislow, 1991; Savage et al., 2007; teleosts: Martins et al., 2021), although for mammals there is a significant positive correlation when analysed with independent contrasts (Promislow, 1991).

Erythrocyte size can also change throughout an organism's life. In some species, erythrocytes can increase in size throughout ontogeny, as seen in monitor lizards (Frýdlová et al., 2013), 10 species of teleost (Lahnsteiner, 2021) and three of five species of eyelid geckos (Starostová et al., 2013). In contrast, erythrocyte size did not change in mole salamanders (Davies, 2008) or two of five species of eyelid geckos (Starostová et al., 2013). Such plasticity is undoubtedly underpinned by between-species differences in oxygen demand as well as changing mass-specific metabolic rate (Pis, 2008).

In other cases, some behavioural and ecological variables may impact erythrocyte sizes. For example, differences in oxygen availability, for example, altitude, can drive variation in blood oxygen-carrying capacity (González-Morales et al., 2015; Lu et al., 2015; Monge & Leon-Velarde, 1991; Navas & Chauí-Berlinck, 2007). Across-species, erythrocytes of anurans are smaller at higher altitudes (Navas & Chauí-Berlinck, 2007), whereas comparisons within species of lizards show increasing erythrocyte size with higher altitude (González-Morales et al., 2017), largest sizes at intermediate altitudes (González-Morales et al., 2017) or decreasing erythrocyte size with higher altitude (Guadarrama et al., 2020). Similarly, species with seasonal altitudinal migration can show plastic size changes in erythrocyte size; alpine accentors (*Prunella collaris*), for example, have smaller erythrocytes when living at high altitude (Haas & Janiga, 2020; Janiga et al., 2017). Other sources of variation in erythrocyte size are less well studied, but again thought to be related to oxygen demand. For example, in tropical reef teleosts, species with a higher aerobic capacity and greater activity levels have smaller erythrocytes (Well & Baldwin, 1990), but more broadly across teleosts, erythrocyte shape was unrelated to habitat or body size (Martins et al., 2021). Similarly, birds with greater energetic demands (e.g. long distance migration) had smaller erythrocytes, whereas diving birds had larger erythrocytes (Soulsbury et al., 2022). These studies demonstrate that components of species ecology and life history are important for determining erythrocyte size. Finally, erythrocyte shape can also vary, being rounder (smaller length to width ratio) or elongated (greater length to width ratio). Erythrocyte shape is poorly studied, but is known to vary plastically within-species, for example, elongated erythrocytes were reported in alpine accentors breeding at high altitudes (Haas & Janiga, 2020; Janiga et al., 2017), whereas across bird species, erythrocytes were more elongated at higher altitudes and for species migrating long distances (Soulsbury et al., 2022). In contrast, cell shape (PCA of cell shape components) also varies extensively in teleosts, but did not vary with habitat or body size (Martins et al., 2021). To date,

erythrocyte shape is relatively unexplored and key drivers of shape variation are uncertain.

Lepidosauria (lizards, snakes and tuatara) are excellent models to study the relationships between ecology and physiology, because of their widespread global distribution and their diversification in a range of ecological niches (Meiri, 2018). As such, cross-species analyses of Lepidosauria allow the study of key patterns in evolutionary ecology at a global scale (Meiri et al., 2020a; Slavenko et al., 2019). A number of key studies have examined macroecological variation and ecological drivers in key traits in squamates (lizards and snakes). These traits include body size (Slavenko et al., 2019), body plan (Wiens et al., 2006), reproductive mode (Ma et al., 2018; Pincheira-Donoso et al., 2017; Rafferty et al., 2013) and life history (Costa et al., 2008; Meiri et al., 2020a, 2020b; Scharf et al., 2015). Studies that examine environmental physiology on a global scale are generally rarer and have mostly focused on thermal ecology (Andrews & Schwarzkopf, 2012; Garcia-Porta et al., 2019).

In this study, we take a first step in understanding the relationships between ecology and physiology by compiling a unique dataset to analyse whether a variety of life-history and ecological traits have driven selection on erythrocyte size in Lepidosauria. We built on recent analyses in birds (Soulsbury et al., 2022), and predict that components of Lepidosauria ecology and behaviour will play a major role in shaping erythrocyte sizes. Smaller erythrocytes may be predicted to occur where the ability and speed to exchange gases between pulmonary and tissues are critical. Hence, we predict that where tissue-oxygen demands are likely to be increased (higher body temperatures, active foraging) or there is a need for enhanced gas exchange (altitude), species will have smaller erythrocytes. Conversely, where there is a need for greater within-blood oxygen-carriage (viviparity vs. oviparity) and storage (diving), there will be increases in erythrocyte size. Finally, we test the same set of factors in relation to erythrocyte shape. However, too few studies have previously tested this parameter, hence the predictions may be too speculative.

2 | MATERIALS AND METHODS

2.1 | Data collection: Erythrocyte size

We gathered data from the published literature. Specifically, we focussed on erythrocyte area (μm^2), rather than mean corpuscular volume (MCV). MCV is usually derived by multiplying a volume of blood by the proportion of blood that is cellular (the haematocrit), and dividing that product by the number of erythrocytes (red blood cells) in that volume. This misses key information on cellular shape. We instead collected length of the long axis (L , in μm) and width of the short axis (W , in μm), and area, which was calculated as the area based on length \times width measurements of an ellipsoid ($\text{area} = \pi \cdot L/2 \cdot W/2$ in μm^2). We calculated the elongation ratio of the cell using the length and width of the erythrocytes (L/W). Elongation ratio is a dimensionless measure, but captures whether

the erythrocyte is round ($L/W = 1$), oval shaped ($L/W = 1.4$ – 1.6) or flatter ellipses ($L/Q = >1.8$). Greater elongation ratios give greater surface areas for a given volume (see example in bacteria: Ojick et al., 2019), which may impact gas exchange rates.

2.2 | Data collection: Species ecology and natural history

The data set comes from populations across the five continents even though there is a bias for some reptile biodiversity hotspots, particularly in south eastern Europe (Figure S1). For each species, we extracted the location of capture (where possible). Altitude was included as reported in the articles or as calculated from <http://www.twcc.fr/> using the coordinates given in the original publication. The exception was one species of sea snake, which had no capture location but was collected at sea level.

Few studies had any information on study-specific body masses, ages or sexes of sampled animals. To include body mass, we used published datasets to assign body mass (Feldman et al., 2016). We also included the reproductive mode (oviparous or viviparous) (Pyrone & Burbrink, 2014), but although there is a continuum of maternal-foetal contact (Albergotti & Guillet, 2011), we lacked enough data to assign these to individual species. Any ovo-viviparous species in our data set were considered viviparous in this context because the mother carries the young until full term.

We also included information on whether the species regularly dives, that is, whether the species was aquatic or semi-aquatic, or not (yes/no). Where information was available, we included information on foraging mode. We used the nomenclature of Meiri (2018) who categorized Sauria (lizards) as ambush predators ('sit and wait'), active foragers ('active foraging') or using a mixed strategy ('mixed'). We extended these data to include snakes, some additional lizards and the tuatara using additional datasets, for example, Lourdaï et al. (2014) and Glaudas et al. (2019), and natural history descriptions. Our distribution of foraging strategies had a limited number of 'mixed' strategies. Instead, we simplified categories to those that were active foragers or those using ambush foraging some or all of the time, that is, combining 'ambush' and 'mixed' categories. Finally, we used existing sources to include species-specific mean body temperature ($^{\circ}\text{C}$) where measured; for ectotherms, most behavioural and physiological processes depend directly on body temperature (see dataset for sources).

2.3 | Data processing and statistical analysis

From our initial data set, we carried out four different analyses covering differing information per study/species (Table S1). Initially, we built the allometric relationships between \log_{10} -transformed body size and erythrocyte size and shape. Second, we analysed a model that tested the effects of body mass, altitude, diving behaviour and

reproductive mode. Initially, we considered interactions between variables, specifically reproductive mode and altitude. However, interactions were not significant, and the model was simplified. The third model included the additional variable of foraging mode. In this analysis, we note a clear outlier during exploratory plotting – the rock python *Python sebae*. In our data set, the erythrocyte size for this species was taken from a study using small ($\sim 2.7\text{kg}$) snakes (Jegade et al., 2020) and body masses of adults are $\sim 50\text{kg}$ (Feldman et al., 2016). There may be age- or size-specific changes in foraging, so we chose to exclude this single data point as we were not sure how to categorize such small snakes. Our final model tested the additional effect of mean body temperature to the second model. The use of these four models maximized coverage of the data but meant that there were differing numbers of species and samples from each species for each of the four different models.

Analysis was carried using phylogenetic mixed models (PMM) using the MCMCglmm package (Hadfield, 2010), with the inclusion of a phylogenetic covariance matrix, with species retained as a second random effect within the models ($\text{random} = \sim \text{animal}$). The phylogenetic covariance matrix was created using a time-calibrated Lepidosauria phylogeny (Tonini et al., 2016), and pruned to fit the species in each model subset. For the PMM models, we set uninformative priors (R -inverse.gamma($V = 1$, $\text{nu} = 0.002$); G -inverse.gamma($V = 1$, $\text{nu} = 0.002$)), used a single independent chain of 500,000 iterations, with sampling taking place every 500 iterations after a 10,000 burn in. The phylogenetic signal (λ) was estimated by dividing the variance explained by the phylogeny, by the sum of all variance components. The phylogenetic signal varies between 0 and 1, where 0 represents no evolutionary signal (no covariance in the residuals due to shared ancestry), and 1 indicates that the observed covariance in residuals follows that expected under a Brownian motion model of trait evolution (Freckleton et al., 2002). Convergence of the MCMC chain was checked using Heidelberger and Welch's convergence diagnostic test for stationarity (implemented in the R package *coda* 0.19–3; Plummer et al., 2006), and the level of autocorrelation was checked to ensure adequate ($>1,000$) effective sample size for each estimated parameter. All models were run in R version 4.0.3 (R Core Team, 2020).

3 | RESULTS

3.1 | General patterns of erythrocyte size

Erythrocyte size varied considerably between species (Figure 1). The lizard with the smallest erythrocyte areas was the eastern smooth-throated lizard *Liolaemus wiegmannii* ($47.12 \mu\text{m}^2$) and the snake with the smallest erythrocyte was the Egyptian cobra *Naja haje* ($52.77 \mu\text{m}^2$). The species with the three largest erythrocyte areas were marine file snake *Acrochordus granulatus* ($287.11 \mu\text{m}^2$), the Komodo dragon *Varanus komodoensis* ($252.90 \mu\text{m}^2$) and the tuatara *Sphenodon punctatus* ($257.62 \mu\text{m}^2$; Figure 1). Most species had




FIGURE 1 Mean erythrocyte area (μm^2) and mean elongation ratio for individual species across the phylogenetic relationship of Lepidosauria. For illustration, the approximate location of representative taxa (top to bottom: *Sphenodon punctatus*; *Gehyra mutilata*; *Eumeces fasciatus*; *Lacerta agilis*; *Varanus komodoensis*; *Stellagama stellio*; *Boa constrictor*; *Vipera latastei*; *Notoechis ater*; *Elaphe quadrivirgata*) are included as tip labels were too small to be visible. Species that are underlined are not in the phylogeny but represent closely related species. All images are taken from Phylopic (<http://phylopic.org>) under Public Domain Dedication 1.0 license (<http://creativecommons.org/publicdomain/zero/1.0/>)

an erythrocyte area lying within the range of 50–200 μm^2 (95.4%, 272/285 species). For elongation ratios, the species with roundest cells were snakes, such as the eastern ribbon snake *Thamnophis saurita* [sauritus] (ratio = 1.22) and the Montpellier snake *Malpolon monspessulanus* (ratio = 1.23; Figure 1), with the most elongated cells found in eastern smooth-throated lizard *Liolaemus wiegmannii* (ratio = 2.40) and the coastal viper *Montivipera xanthina* (ratio = 2.37; Figure 1).

Irrespective of species, larger Lepidosauria had significantly larger erythrocytes in terms of area (Posterior mean = 11.43, upper and lower bounds of the 95% Credible Intervals (CI) = 5.48–17.43, pMCMC <0.001; λ = 0.70 (0.54/0.83)), Figure 2; N = 285 from 235 species). In contrast, there was no effect of body mass on elongation ratio (Posterior mean = 0.01, 95% CI = -0.03–0.04, pMCMC = 0.926; λ = 0.42 (0.14/0.67); N = 266 from 227 species).

3.2 | Effect of diving, altitude and reproductive mode on erythrocyte size

Species that dived had significantly larger erythrocytes (Table 1a; Figure 3a), but no change in elongation ratio (Table 1a). Viviparous species also had erythrocytes with a larger area (Figure 3b), but elongation ratio did not differ between oviparous and viviparous species (Table 1a). Finally, altitude had a significant negative effect on erythrocyte area (Table 1a; Figure 3c), and had a positive relationship with elongation ratio, that is, more elongated cells (Table 1a; Figure 4). Interestingly, viviparous species had larger erythrocytes than oviparous species irrespective of altitude (Figure 3c).




FIGURE 2 Relationships between body mass (\log_{10} , g) and erythrocyte area (μm^2). Raw data points are shown, with regression line (+95CI) drawn through predicted values derived from the phylogenetic MCMC model

3.3 | Foraging, body temperature and erythrocyte size

Across species, those that were active foragers as opposed to those using mixed/sit-and-wait strategies had significantly smaller erythrocytes (Table 1c; Figure 5a). Similarly, those species with higher mean body temperatures had smaller erythrocytes (Table 1c; Figure 5b). In contrast, neither foraging strategy or mean body temperature affected elongation ratio (Table 1c).

4 | DISCUSSION

In Lepidosauria, there were significant positive relationships between body mass and erythrocyte size but no significant effect on shape, as measured by elongation ratio. Our analysis found that while controlling for body mass there were complex patterns of changes in Lepidosaurian erythrocyte size and shape. Some drivers were linked to environmental conditions, including altitude and climate, whereas others were related to life-history and behaviour factors such as reproductive mode, body temperature, foraging or diving.

4.1 | Body mass effects

We found a significant positive interspecific relationship between body mass and erythrocyte size in Lepidosauria, which has also been found within species (Frýdlová et al., 2013; Starostová et al., 2005). Increased body size is also associated with larger erythrocytes in amphibians (Gregory, 2001; Hawkey et al., 1991) and birds (Kostecka-Myrcha & Chołostiakow-Gromek, 2001; Soulsbury et al., 2022). The increase in genome size with body size (Gregory, 2000, 2001, 2002; Mueller et al., 2008), leads to larger nucleus sizes; larger erythrocytes are, therefore, needed to house the larger nucleus. Hence, the anucleate erythrocytes of mammals do not exhibit any relationship with body size, although many nucleated cell types do exhibit such a relationship (Savage et al., 2007). Erythrocyte size is larger in fish than in amphibians, which are larger than in reptiles (Hartman & Lesser, 1964), and this reflects broad changes in the pulmonary-circulatory system (Snyder & Sheafor, 1999). Having demonstrated that body size is important in Lepidosauria, further analysis is required to investigate how other blood parameters that are important for oxygen transport are affected by environmental and life-history variables.

4.2 | Altitude and climatic effects on erythrocyte size

Vertebrates that live at high altitudes (above 2000 m) are subjected to hypoxic conditions that challenge aerobic metabolism. They usually exhibit functional and structural modifications that allow them to cope with the concomitant decrease in O_2 tension that potentially

TABLE 1 Phylogenetic mixed model analysis outputs of erythrocyte area and elongation ratio in relation to (a) body mass, altitude, diving behaviour and reproductive mode and (b) foraging mode (mixed or active foraging) and (c) maximum body temperature

	Variable	Posterior mean	lower 95% CI	upper 95% CI	pMCMC ^a
(a) Altitude, diving and reproductive mode					
Area $\lambda = 0.24$ (0.00/0.58) N = 151 (129 species)	Body mass	10.45	3.01	17.26	0.006
	Altitude	-0.01	-0.02	-0.01	0.001
	Diving	32.31	10.66	55.78	0.006
	Reproductive mode	21.44	6.41	37.64	0.005
Elongation ratio $\lambda = 0.38$ (0.03/0.72) N = 132 (118 species)	Altitude	8.21×10^5	2.45×10^5	1.43×10^4	0.005
	Diving	-0.04	-0.13	0.20	0.621
	Reproductive mode	0.04	-0.07	0.16	0.478
(b) Foraging mode					
Area $\lambda = 0.70$ (0.50/0.90) N = 177 (143 species)	Body mass	7.51	-0.90	15.25	0.065
	Diving	38.85	19.33	57.58	0.001
	Reproductive mode	14.43	-0.72	30.17	0.065
	Foraging mode	16.98	2.70	32.66	0.031
Elongation ratio $\lambda = 0.44$ (0.13/0.73) N = 168 (138 species)	Diving	-0.01	-0.14	0.11	0.802
	Reproductive mode	-0.04	-0.14	0.07	0.523
	Foraging mode	0.02	-0.08	0.13	0.634
(c) Body temperature					
Area $\lambda = 0.41$ (0.00/0.65) N = 162 (124 species)	Body mass	13.46	7.50	20.02	0.001
	Diving	15.63	-2.81	33.43	0.094
	Reproductive mode	7.73	-3.87	21.73	0.220
	Mean body temp	-1.48	-2.79	-0.02	0.037
Elongation ratio $\lambda = 0.54$ (0.26/0.78) N = 143 (116 species)	Diving	0.05	-0.08	0.18	0.455
	Reproductive mode	-0.02	-0.13	0.07	0.684
	Mean body temp	0.006	-0.004	0.02	0.244

^aParticle Markov chain Monte Carlo (pMCMC) is the proportion of coefficients in the posterior distribution estimates that are $\neq 0$, multiplied by two (for a two-tailed test), and is analogous to a frequentist p -value.

constrains aerobic life in such environments (González-Morales et al., 2015, 2017). Some of these adaptations include changes in the size or morphology of respiratory surfaces, increased heart size, elevated haemoglobin concentration, higher haematocrit, increased capillary density and elevated myoglobin levels (Monge & Leon-Velarde, 1991; Samaja et al., 2003; Weber, 2007). We show that at higher altitudes, Lepidosauria in general had smaller erythrocytes. Similarly, across several species of anurans, species from higher altitudes also had smaller erythrocytes (Navas & Chauí-Berlinck, 2007), whereas in lizards, within-species comparisons showed mixed effects of altitude on erythrocyte size (González-Morales et al., 2015, 2017; Guadarrama et al., 2020). Size and shape affect surface area to volume ratio, meaning that small erythrocytes will exchange oxygen more efficiently than larger cells, and ellipsoid cells are more efficient than round cells (Hartman & Lesser, 1963). Another haematological parameter, haematocrit, that is, the ratio of the volume of red blood cells to the total volume of blood, tends to be increased at higher altitudes (Lu et al., 2015). Increased haematocrit allows greater oxygen-carrying capacity, but at the same time increases blood viscosity (Mehri et al., 2018). Smaller erythrocytes allow more rapid oxygenation and deoxygenation of haemoglobin (Holland &

Forster, 1966; Jones, 1979) without perhaps the deleterious increase in viscosity. Although not part of the current study, an understanding of the relationship between cell size and haematocrit would help determine whether high altitude, especially in species where erythrocytes size is larger at higher altitudes (e.g. González-Morales et al., 2015).

Altitude was also associated with a change in shape with cells becoming more elongated with increasing altitudinal hypoxia. In birds, those found at higher altitudes also had more elongated cells (Soulsbury et al., 2022), suggesting a general benefit of elongated cells in low-oxygen environments. Narrower, more elongated cells may pass through capillary beds more easily and so maximize the rate of blood flow directly or through a reduction in blood viscosity (e.g. alpine accentors: Janiga et al., 2017), although the exact benefits deserve further investigation.

4.3 | Life history and behaviour

Our results had three clear patterns for factors associated with life history and behaviour. First, energetic activities, that is, diving and




FIGURE 3 Boxplots showing the effect of (a) diving and (b) reproductive mode on erythrocyte area (μm^2) and scatterplot showing the effect of (c) altitude and reproductive mode (oviparity = grey circles, viviparity = black circles) on erythrocyte area (μm^2). Raw data points are shown and both boxplots and regression lines (+95CI) are drawn through predicted values derived from the phylogenetic MCMC models




FIGURE 4 The relationship between elongation ratio of erythrocytes and altitude (m). Raw data points are shown, with regression line (+95CI) drawn through predicted values derived from the phylogenetic MCMC model

active foraging in our analysis, had contrasting effects on erythrocyte size. During diving in birds and mammals, the blood is an important oxygen store and haemoglobin concentration is elevated (Minias, 2020). Larger erythrocytes may contribute significantly to binding more oxygen and is, therefore, critical for oxygen storage capacity (Soulsbury et al., 2022). For example, in diving mammals, MCV and erythrocyte sizes are all significantly larger than in non-diving counterparts (Davis, 2014; Wickham et al., 1989). Similarly, reptiles use a range of physiological mechanisms, including bradycardia, prolonged apnoea, cardiac shunts and Bohr effects, to support diving behaviour (Anderson, 1966; Millard & Johansen, 1974; Seymour & Webster, 1975). In diving turtles, oxygen is stored both in the lungs and blood (Arango et al., 2021; Lutz & Bentley, 1985), and

blood oxygen stores are thought to be important in some squamates (Ferguson & Thornton, 1984). Marine snakes have significantly higher haematocrit levels (Brischoux et al., 2011; Feder, 1980), with the file snake *Acrochordus granulatus*, the species with the largest recorded erythrocytes, having the largest blood oxygen stores of any reptile (Feder, 1980). Our work demonstrates that along with other blood oxygen components, diving species have larger erythrocytes which allow greater quantities of oxygen to be stored in the blood. Larger erythrocytes are also found in diving birds (Soulsbury et al., 2022), suggesting a widespread adaptation to diving across vertebrates.

Our second result was the finding that Lepidosauria that gave birth to live young (viviparous and ovo-viviparous) had larger erythrocytes. Producing live offspring places a metabolic cost on the gravid females, leading to higher oxygen consumption especially towards the end of pregnancy (Beuchat & Vleck, 1990; Jackson et al., 2015; Robert & Thompson, 2002; Thompson & Speake, 2006). No study has looked specifically at erythrocyte size in Lepidosauria, and the only similar study showed that erythrocyte size was largest in mid-gestation gravid fish (Ingermann & Terwilliger, 1982), representing a plastic response. Our analysis shows that live birth has led to the selection for larger erythrocytes, perhaps as a result of higher oxygen demands of the female during gestation (Schultz et al., 2008; see also Munns & Daniels, 2007) but it could also be due to greater oxygen requirements of the developing offspring (Foucart et al., 2014). Previously, low oxygen was thought to be a constraint on the evolution of viviparity (Andrews, 2002), but this is unlikely to be the case (see Pincheria-Donoso et al., 2017). Indeed, the number of viviparous snakes and lizards found at higher altitudes suggests the reverse (Feldman et al., 2015; Pincheria-Donoso et al., 2017) and there is a need to consider the interaction between oxygen demand, altitude and the evolution of viviparity (Watson & Cox, 2021). It should be pointed out that our analysis did not take into account whether viviparous species had placental maternal transfer. This could be important because ovo-viviparity involves the retention of embryos *in*




FIGURE 5 The relationship between erythrocyte area (μm^2) and (a) foraging mode (active or ambush foraging) and (b) mean recorded body temperature (degrees Celsius) across Lepidosauria

utero but there is no development of a placental-like structure that facilitates transfer of nutrition; this may also affect the transfer of respiratory gases (Blackburn & Stewart, 2011). Although not directly measured, it is inferred that viviparous species with placentation exchange respiratory gases and this is facilitated by changes in maternal blood chemistry that enhances oxygen exchange (Ingerman et al., 1991). To fully understand the link between oxygen demand, oxygen transport and reproductive mode, examples of within-species (Rechnagel et al., 2021) and within-individual variation (Laird et al., 2019), that is, showing both oviparous and viviparous reproductive modes, as well as cross-species comparisons (Foster et al., 2020) all provide ideal opportunities to link evolutionary form to function. Our

analysis suggests that changing blood oxygen-carrying capacity by altering size and/or shape of erythrocytes is one mechanism to avoid constraints of low oxygen and may allow viviparous Lepidosauria to exploit higher altitude areas. In addition, future work should include the diversity of placenta types, and hence the degree of maternal-foetal contact, as an important modifier of oxygen transport and delivery (Stewart & Blackburn, 1988; Thompson & Speake, 2006).

Finally, we found that active foraging species of Lepidosauria and those with higher mean body temperatures had smaller erythrocytes. In line with this, a handful of studies have found that erythrocyte size negatively correlates with activity, for example, aerobic swimming ability in teleosts (Lay & Baldwin, 1999) and migration distance in birds (Soulsbury et al., 2022). Differences in activity also alter erythrocyte size in some anurans (Atatür et al., 1999). Our work supports the idea that rapid oxygenation and deoxygenation of haemoglobin (Holland & Forster, 1966; Jones, 1979), facilitated by having smaller erythrocytes, is a key component of an active lifestyle. In line with this, species with higher mean body temperatures had smaller erythrocytes. High temperatures lead to higher metabolic rates and oxygen consumption (e.g. Clark et al., 2006) and in turn, across and within species, higher metabolic rates are associated with smaller erythrocyte size (Bury et al., 2019; Czarnoleski et al., 2017; Goodman & Heath, 2010; Gregory, 2002; Starostová et al., 2009). This further emphasizes the importance of mode of life and environmental conditions as important drivers of erythrocyte size.

5 | CONCLUSIONS

This study has demonstrated that across Lepidosauria erythrocyte size and shape can be affected by a range of biotic and abiotic factors and are linked to overall metabolism. Although limited to cell dimensions, the analysis demonstrates that this could be one of the key factors affecting changes in blood oxygenation required by different life histories in different habitats. It would be interesting to extend this analysis to determine whether factors, such as altitude, diving or sustained activity, also affect other key aspects of Lepidosaurian haematology, for example, haematocrit and haemoglobin concentration.

ACKNOWLEDGEMENTS

We are grateful to the two reviewers (Diogo Provete & Shai Meiri) whose valuable comments really benefitted the manuscript.

CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

AUTHOR CONTRIBUTIONS

CDS conceived the ideas and analysed the data; ZP and CDS collected the data; CDS and DCD led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/jeb.14004>.

OPEN RESEARCH BADGES



This article has been awarded Open Materials, Open Data, Badges. All materials and data are publicly accessible via the Open Science Framework at <https://doi.org/10.6084/m9.figshare.19328945.v1>.

DATA AVAILABILITY STATEMENT

Data and R code are available at FigShare (<https://doi.org/10.6084/m9.figshare.19328945.v1>). All data are taken from available sources.

ORCID

D. Charles Deeming  <https://orcid.org/0000-0002-9587-6149>

Carl D. Soulsbury  <https://orcid.org/0000-0001-8808-5210>

REFERENCES

- Albergotti, L. C., & Guillelte, L. J. Jr (2011). Viviparity in reptiles: Evolution and reproductive endocrinology. In D. O. Norris, & K. H. Lopez *Hormones and reproduction of vertebrates* Vol 3, (pp. 247–275). Academic Press.
- Anderson, H. (1966). Physiological adaptations to diving in vertebrates. *Physiological Reviews*, 46, 212–243.
- Andrews, R. M. (2002). Low oxygen: a constraint on the evolution of viviparity in reptiles. *Physiological and Biochemical Zoology*, 75, 145–154.
- Andrews, R. M., & Schwarzkopf, L. (2012). Thermal performance of squamate embryos with respect to climate, adult life history, and phylogeny. *Biological Journal of the Linnean Society*, 106, 851–864.
- Arango, B. G., Harfush-Meléndez, M., Marmolejo-Valencia, J. A., Merchant-Larios, H., & Crocker, D. E. (2021). Blood oxygen stores of olive ridley sea turtles, *Lepidochelys olivacea* are highly variable among individuals during Arribada nesting. *Journal of Comparative Physiology B*, 191, 185–194.
- Atatür, M. K., Arikan, H., & Çevik, I. E. (1999). Erythrocyte sizes of some anurans from Turkey. *Turkish Journal of Zoology*, 23, 111–114.
- Beuchat, C. A., & Vleck, D. (1990). Metabolic consequences of viviparity in a lizard, *Sceloporus jarrovi*. *Physiological Zoology*, 63, 555–570.
- Blackburn, D. G., & Stewart, J. R. Jr (2011). Viviparity and placentation in snakes. In D. Sever, & R. Aldridge (Eds.), *Reproductive biology and phylogeny of snakes* (pp. 119–181). CRC Press.
- Brischoux, F., Gartner, G. E., Garland, T. Jr, & Bonnet, X. (2011). Is aquatic life correlated with an increased hematocrit in snakes? *PLoS One*, 6(2), e17077. <https://doi.org/10.1371/journal.pone.0017077>
- Bury, S., Bury, A., Sadowska, E. T., Cichoń, M., & Bauchinger, U. (2019). More than just the numbers—contrasting response of snake erythrocytes to thermal acclimation. *Science Nature*, 106, 24.
- Clark, T. D., Butler, P. J., & Frappell, P. B. (2006). Factors influencing the prediction of metabolic rate in a reptile. *Functional Ecology*, 20, 105–113.
- Costa, G. C., Vitt, L. J., Pianka, E. R., Mesquita, D. O., & Colli, G. R. (2008). Optimal foraging constrains macroecological patterns: body size and dietary niche breadth in lizards. *Global Ecology and Biogeography*, 17, 670–677.
- Czarnoleski, M., Labecka, A. M., Starostová, Z., Sikorska, A., Bonda-Ostaszewska, E., Woch, K., Kubička, L., Kratochvíl, L., & Kozłowski, J. (2017). Not all cells are equal: effects of temperature and sex on the size of different cell types in the Madagascar ground gecko *Paroedura picta*. *Biology Open*, 6, 1149–1154.
- Davis, A. K. (2008). Ontogenetic changes in erythrocyte morphology in larval mole salamanders, *Ambystoma talpoideum*, measured with image analysis. *Comparative Clinical Pathology*, 17, 23–28.
- Davis, R. W. (2014). A review of the multi-level adaptations for maximizing aerobic dive duration in marine mammals: from biochemistry to behavior. *Journal of Comparative Physiology B*, 184, 23–53.
- Feder, M. E. (1980). Blood oxygen stores in the file snake, *Acrochordus granulatus*, and in other marine snakes. *Physiological Zoology*, 53, 394–401.
- Feldman, A., Bauer, A. M., Castro-Herrera, F., Chirio, L., Das, I., Doan, T. M., Maza, E., Meirte, D., de Campos Nogueira, C., Nagy, Z. T., & Torres-Carvajal, O. (2015). The geography of snake reproductive mode: a global analysis of the evolution of snake viviparity. *Global Ecology and Biogeography*, 24, 1433–1442.
- Feldman, A., Sabath, N., Pyron, R. A., Mayrose, I., & Meiri, S. (2016). Body sizes and diversification rates of lizards, snakes, amphibians and the tuatara. *Global Ecology and Biogeography*, 25, 187–197.
- Ferguson, J. H., & Thornton, R. M. (1984). Oxygen storage capacity and tolerance of submergence of a non-aquatic reptile and an aquatic reptile. *Comparative Biochemistry and Physiology Part A, Physiology*, 77, 183–187.
- Foster, C. S., Thompson, M. B., Van Dyke, J. U., Brandley, M. C., & Whittington, C. M. (2020). Emergence of an evolutionary innovation: gene expression differences associated with the transition between oviparity and viviparity. *Molecular Ecology*, 29, 1315–1327. <https://doi.org/10.1111/mec.15409>
- Foucart, T., Lourdais, O., DeNardo, D. F., & Heulin, B. (2014). Influence of reproductive mode on metabolic costs of reproduction: insight from the bimodal lizard *Zootoca vivipara*. *Journal of Experimental Biology*, 217, 4049–4056.
- Freckleton, R. P., Harvey, P. H., & Pagel, M. (2002). Phylogenetic analysis and comparative data: a test and review of evidence. *The American Naturalist*, 160, 712–726.
- Frydlová, P., Hnízdo, J., Chylíková, L., Šimková, O., Cikánová, V., Velenský, P., & Daniel Frynta, D. (2013). Morphological characteristics of blood cells in monitor lizards: is erythrocyte size linked to actual body size? *Integrative Zoology*, 8, 39–45. <https://doi.org/10.1111/j.1749-4877.2012.00295.x>
- García-Porta, J., Irisarri, I., Kirchner, M., Rodríguez, A., Kirchoff, S., Brown, J. L., MacLeod, A., Turner, A. P., Ahmadzadeh, F., Albaladejo, G., & Crnobrnja-Isailovic, J. (2019). Environmental temperatures shape thermal physiology as well as diversification and genome-wide substitution rates in lizards. *Nature Communications*, 10, 1–12.
- Glaudas, X., Glennon, K. L., Martins, M., Luiselli, L., Fearn, S., Trembath, D. F., Jelić, D., & Alexander, G. J. (2019). Foraging mode, relative prey size and diet breadth: A phylogenetically explicit analysis of snake feeding ecology. *Journal of Animal Ecology*, 88, 757–767. <https://doi.org/10.1111/1365-2656.12972>
- González-Morales, J. C., Quintana, E., Díaz-Albiter, H., Guevara-Fiore, P., & Fajardo, V. (2015). Is erythrocyte size a strategy to avoid hypoxia in Wiegmann's torquate lizards (*Sceloporus torquatus*)? Field evidence. *Canadian Journal of Zoology*, 93, 377–382.
- González-Morales, J. C., Beamonte-Barrientos, R., Bastiaans, E., Guevara-Fiore, P., Quintana, E., & Fajardo, V. (2017). A mountain or a plateau? hematological traits vary nonlinearly with altitude in a highland lizard. *Physiological and Biochemical Zoology*, 90, 638–645.
- Goodman, R. M., & Heah, T. P. (2010). Temperature-induced plasticity at cellular and organismal levels in the lizard *Anolis carolinensis*. *International Zoology*, 5, 208–217.
- Gregory, T. R. (2000). Nucleotypic effects without nuclei: genome size and erythrocyte size in mammals. *Genome*, 43, 895–901. <https://doi.org/10.1139/g00-069>

- Gregory, T. R. (2001). The bigger the C-value, the larger the cell: genome size and red blood cell size in vertebrates. *Blood Cells, Molecules, and Diseases*, 27, 830–843.
- Gregory, T. R. (2002). A bird's-eye view of the C-value enigma: genome size, cell size, and metabolic rate in the class Aves. *Evolution*, 56, 121–130.
- Guadarrama, S. S., Domínguez-Vega, H., Díaz-Albiter, H. M., Quijano, A., Bastiaans, E., Carrillo-Castilla, P., Manjarrez, J., Gómez-Ortíz, Y., & Fajardo, V. (2020). Hypoxia by altitude and welfare of captive beaded lizards (*Heloderma horridum*) in Mexico: Hematological approaches. *Journal of Applied Animal Welfare Science*, 23, 74–82.
- Haas, M., & Janiga, M. (2020). Variation in erythrocyte morphology in alpine accentors (*Prunella collaris* Scop.) from Tian Shan, Rila and the High Tatra mountains and effects of molting. *European Zoology Journal*, 87, 475–488.
- Hadfield, J. D. (2010). MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. *The Journal of Statistical Software*, 33, 1–22.
- Hartman, F. A., & Lesser, M. A. (1963). Erythrocyte size in birds. *The Auk*, 80, 467–473.
- Hartman, F. A., & Lesser, M. A. (1964). Erythrocyte measurements in fishes, amphibia, and reptiles. *Biological Bulletin*, 126, 83–88.
- Hawkey, C. M., Bennett, P. M., Gascoyne, S. C., Hart, M. G., & Kirkwood, J. K. (1991). Erythrocyte size, number and haemoglobin content in vertebrates. *The British Journal of Haematology*, 77, 392–397.
- Holland, R. A. B., & Forster, R. E. (1966). The effect of size of red cells on the kinetics of their oxygen uptake. *Journal of General Physiology*, 49, 727–742.
- Ingermann, R. I., Berner, N. J., & Ragsdale, F. R. (1991). effect of pregnancy and temperature on red cell oxygen-affinity in the viviparous snake *Thamnophis elegans*. *Journal of Experimental Biology*, 156, 399–406.
- Ingermann, R. L., & Terwilliger, R. C. (1982). Blood parameters and facilitation of maternal-fetal oxygen transfer in a viviparous fish (*Embiotoca lateralis*). *Comparative Biochemistry and Physiology Part A, Physiology*, 73, 497–501.
- Jackson, A. G., Leu, S. Y., Ford, N. B., & Hicks, J. W. (2015). Patterns of oxygen consumption during simultaneously occurring elevated metabolic states in the viviparous snake *Thamnophis marcianus*. *Journal of Experimental Biology*, 218, 3570–3579.
- Janiga, M., Haas, M., & Kufelová, M. (2017). Age, sex and seasonal variation in the shape and size of erythrocytes of the alpine accentor, *Prunella collaris* (Passeriformes: Prunellidae). *European Zoology Journal*, 84, 583–590.
- Jegede, H. O., Olaifa, F. H., Adah, A. S., Adah, A. D., & Omobowale, T. O. (2020). Erythrocyte osmotic fragility and erythrocyte sizes for captive African rock pythons (*Python sebae*). *Comparative Clinical Pathology*, 29, 317–320. <https://doi.org/10.1007/s00580-019-03064-0>
- Jones, D. A. (1979). The importance of surface area/volume ratio to the uptake of oxygen by red cells. *Journal of General Physiology*, 74, 643–646.
- Kostelecka-Myrcha, A., & Chołostiakow-Gromek, J. (2001). Body mass dependence of the haemoglobin content to surface area ratio of avian erythrocytes. *Acta Ornithologica*, 36, 123–128.
- Kozłowski, J., Konarzewski, M., & Czarnoleski, M. (2020). Coevolution of body size and metabolic rate in vertebrates: a life-history perspective. *Biological Reviews*, 95, 1393–1417.
- Lahnsteiner, F. (2021). Erythrocyte morphometry in teleost fish—Species-specific, inter-individual and environmental-related differences. *Acta Zoology*, 102, 237–247.
- Laird, M. K., Thompson, M. B., & Whittington, C. M. (2019). Facultative oviparity in a viviparous skink (*Saiphos equalis*). *Biology Letters*, 15, 20180827.
- Lay, P. A., & Baldwin, J. (1999). What determines the size of teleost erythrocytes? Correlations with oxygen transport and nuclear volume. *Fish Physiology and Biochemistry*, 20, 31–35.
- Lourdais, O., Gartner, G. E., & Brischoux, F. (2014). Ambush or active life: foraging mode influences haematocrit levels in snakes. *Biological Journal of the Linnean Society*, 111, 636–645.
- Lu, S., Xin, Y., Tang, X., Yue, F., Wang, H., Bai, Y., Niu, Y., & Chen, Q. (2015). Differences in hematological traits between high- and low-altitude lizards (genus *Phrynocephalus*). *PLoS One*, 10, e0125751.
- Lutz, P. L., & Bentley, T. B. (1985). Respiratory physiology of diving in the sea turtle. *Copeia*, 1985, 671–679.
- Ma, L., Buckley, L. B., Huey, R. B., & Du, W. G. (2018). A global test of the cold-climate hypothesis for the evolution of viviparity of squamate reptiles. *Global Ecology and Biogeography*, 27, 679–689.
- Martins, B. O., Franco-Belussi, L., Siqueira, M. S., Fernandes, C. E., & Provet, D. B. (2021). The evolution of red blood cell shape in fishes. *Journal of Evolutionary Biology*, 34, 537–548. <https://doi.org/10.1111/jeb.13757>
- Mehri, R., Mavriplis, C., & Fenech, M. (2018). Red blood cell aggregates and their effect on non-Newtonian blood viscosity at low hematocrit in a two-fluid low shear rate microfluidic system. *PLoS One*, 13(7), e0199911. <https://doi.org/10.1371/journal.pone.0199911>
- Meiri, S. (2018). Traits of lizards of the world: variation around a successful evolutionary design. *Global Ecology and Biogeography*, 27, 1168–1172.
- Meiri, S., Avila, L., Bauer, A. M., Chapple, D. G., Das, I., Doan, T. M., Doughty, P., Ellis, R., Grismer, L., Kraus, F., & Morando, M. (2020a). The global diversity and distribution of lizard clutch sizes. *Global Ecology and Biogeography*, 29, 1515–1530.
- Meiri, S., Feldman, A., Schwarz, R., & Shine, R. (2020b). Viviparity does not affect the numbers and sizes of reptile offspring. *Journal of Animal Ecology*, 89, 360–369.
- Millard, R. W., & Johansen, K. (1974). Ventricular outflow dynamics in a lizard, *Varanus niloticus* - Responses to hypoxia, hypercarbia and diving. *Journal of Experimental Biology*, 60, 871–880.
- Minias, P. (2020). Ecology and evolution of blood oxygen-carrying capacity in birds. *The American Naturalist*, 195, 788–801.
- Monge, C., & Leon-Velarde, F. (1991). Physiological adaptation to high altitude: Oxygen transport in mammals and birds. *Physiological Reviews*, 71, 1135–1172.
- Mueller, R. L., Gregory, T. R., Gregory, S. M., Hsieh, A., & Boore, J. L. (2008). Genome size, cell size, and the evolution of enucleated erythrocytes in attenuate salamanders. *Zoology (Jena)*, 111, 218–230.
- Munns, S., & Daniels, C. (2007). Breathing with big babies: ventilation and oxygen consumption during pregnancy in the lizard *Tiliqua rugosa*. *Physiological and Biochemical Zoology*, 80, 35–45.
- Navas, C. A., & Chaui-Berlinck, J. G. (2007). Respiratory physiology of high-altitude anurans: 55 years of research on altitude and oxygen. *Respiratory Physiology & Neurobiology*, 158, 307–313.
- Ojkic, N., Serbanescu, D., & Banerjee, S. (2019). Surface-to-volume scaling and aspect ratio preservation in rod-shaped bacteria. *Elife*, 8, e47033. <https://doi.org/10.7554/eLife.47033>
- Pincheira-Donoso, D., Jara, M., Reaney, A., García-Roa, R., Saldarriaga-Córdoba, M., & Hodgson, D. J. (2017). Hypoxia and hypothermia as rival agents of selection driving the evolution of viviparity in lizards. *Global Ecology and Biogeography*, 26, 1238–1246.
- Pis, T. (2008). Resting metabolic rate and erythrocyte morphology in early development of thermoregulation in the precocial grey partridge (*Perdix perdix*). *Comparative Biochemistry and Physiology Part A, Physiology*, 151, 211–218.
- Plummer, M., Best, N., Cowles, K., & Vines, K. (2006). CODA: convergence diagnosis and output analysis for MCMC. *R News*, 6, 7–11.
- Promislow, D. E. (1991). The evolution of mammalian blood parameters: patterns and their interpretation. *Physiological Zoology*, 64, 393–431.
- Pyron, R. A., & Burbrink, F. T. (2014). Early origin of viviparity and multiple reversions to oviparity in squamate reptiles. *Ecology Letters*, 17, 13–21.

- R Core Team (2020). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Rafferty, A. R., Evans, R. G., Scheelings, T. F., & Reina, R. D. (2013). Limited oxygen availability in utero may constrain the evolution of live birth in reptiles. *American Naturalist*, *181*, 245–253.
- Recknagel, H., Carruthers, M., Yurchenko, A. A., Nokhbatolfigohahai, M., Kamenos, N. A., Bain, M. M., & Elmer, K. R. (2021). The functional genetic architecture of egg-laying and live-bearing reproduction in common lizards. *Nature Ecology: Evolution*, *5*, 1546–1556. <https://doi.org/10.1038/s41559-021-01555-4>
- Robert, K. A., & Thompson, M. B. (2000). Energy consumption by embryos of a viviparous lizard, *Eulamprus tympanum*, during development. *Comparative Biochemistry and Physiology*, *127*, 481–486.
- Samaja, M., Crespi, T., Guazzi, M., & Vandergriff, K. D. (2003). Oxygen transport in blood at high altitude: Role of the hemoglobin-oxygen affinity and impact of the phenomena related to hemoglobin allosterism and red cell function. *European Journal of Applied Physiology*, *90*, 351–359. <https://doi.org/10.1007/s00421-003-0954-8>
- Savage, V. M., Allen, A. P., Brown, J. H., Gillooly, J. F., Herman, A. B., Woodruff, W. H., & West, G. B. (2007). Scaling of number, size, and metabolic rate of cells with body size in mammals. *Proceedings of the National Academy of Sciences of the United States of America*, *104*, 4718–4723.
- Scharf, I., Feldman, A., Novosolov, M., Pincheira-Donoso, D., Das, I., Böhm, M., Uetz, P., Torres-Carvajal, O., Bauer, A., Roll, U., & Meiri, S. (2015). Late bloomers and baby boomers: ecological drivers of longevity in squamates and the tuatara. *Global Ecol. Biogeog.*, *24*, 396–405.
- Schultz, T. J., Webb, J. K., & Christian, K. A. (2008). The physiological cost of pregnancy in a tropical viviparous snake. *Copeia*, *2008*, 637–642.
- Seymour, R. S., & Webster, M. E. D. (1975). Gas transport and blood acid-base balance in diving sea snakes. *Journal of Experimental Zoology*, *191*, 169–181.
- Slavenko, A., Feldman, A., Allison, A., Bauer, A. M., Böhm, M., Chirio, L., Colli, G. R., Das, I., Doan, T. M., LeBreton, M., & Martins, M. (2019). Global patterns of body size evolution in squamate reptiles are not driven by climate. *Global Ecology and Biogeography*, *28*, 471–483.
- Snyder, G. K., & Sheafor, B. A. (1999). Red blood cells: centerpiece in the evolution of the vertebrate circulatory system. *American Zoologist*, *39*, 189–198.
- Soulsbury, C. D., Dobson, J., Deeming, D. C., & Minias, P. (2022). Energetic lifestyle drives size and shape of avian erythrocytes. *Integrative and Comparative Biology*, *62*. <https://doi.org/10.1093/icb/icab195>
- Starostová, Z., Konarzewski, M., Kozłowski, J., & Kratochvíl, L. (2013). Ontogeny of metabolic rate and red blood cell size in eyelid geckos: species follow different paths. *PLoS One*, *8*, e64715. <https://doi.org/10.1371/journal.pone.0064715>
- Starostová, Z., Kratochvíl, L., & Frynta, D. (2005). Dwarf and giant geckos from the cellular perspective: the bigger the animal, the bigger its erythrocytes? *Functional Ecology*, *19*, 744–749.
- Starostová, Z., Kubička, L., Konarzewski, M., Kozłowski, J., & Kratochvíl, L. (2009). Cell size but not genome size affects scaling of metabolic rate in eyelid geckos. *American Naturalist*, *174*, E100–E105.
- Stewart, J. R., & Blackburn, D. G. (1988). Reptilian placentation: structural diversity and terminology. *Copeia*, *1988*, 839–852.
- Thompson, M. B., & Speake, B. K. (2006). A review of the evolution of viviparity in lizards: structure, function and physiology of the placenta. *Journal of Comparative Physiology B*, *176*, 179–189.
- Tonini, J. F. R., Beard, K. H., Ferreira, R. B., Jetz, W., & Pyron, R. A. (2016). Fully-sampled phylogenies of squamates reveal evolutionary patterns in threat status. *Biological Conservation*, *204*, 23–31.
- Watson, C. M., & Cox, C. L. (2021). Elevation, oxygen, and the origins of viviparity. *The Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*, *336*, 457–469.
- Weber, R. E. (2007). High-altitude adaptations in vertebrate hemoglobins. *Respiratory Physiology & Neurobiology*, *158*, 132–142.
- Wells, R. M. G., & Baldwin, J. (1990). Oxygen transport potential in tropical reef fish with special reference to blood viscosity and haematocrit. *Journal of Experimental Marine Biology and Ecology*, *41*, 131–143.
- Wickham, L. L., Elsner, R., White, F. C., & Cornell, L. H. (1989). Blood viscosity in phocid seals: possible adaptations to diving. *Journal of Comparative Physiology B*, *159*, 153–158.
- Wiens, J. J., Brandley, M. C., & Reeder, T. W. (2006). Why does a trait evolve multiple times within a clade? Repeated evolution of snake-like body form in squamate reptiles. *Evolution*, *60*, 123–141. <https://doi.org/10.1554/05-328.1>

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Penman, Z., Deeming, D. C., & Soulsbury, C. D. (2022). Ecological and life-history correlates of erythrocyte size and shape in Lepidosauria. *Journal of Evolutionary Biology*, *00*, 1–11. <https://doi.org/10.1111/jeb.14004>